

CS23331-DAA-2024-CSE / 5-Pair with Difference- $O(n^2)$ Time Complexity, $O(1)$ Space Complexity


5-Pair with Difference- $O(n^2)$ Time Complexity, $O(1)$ Space Complexity

Started on	Wednesday, 5 November 2025, 6:56 AM
State	Finished
Completed on	Wednesday, 5 November 2025, 6:56 AM
Time taken	36 secs
Marks	1.00/1.00
Grade	4.00 out of 4.00 (100%)

Question 1 | Correct Mark 1.00 out of 1.00 [Flag question](#)

Given an array A of sorted integers and another non negative integer k, find if there exists 2 indices i and j such that $A[j] - A[i] = k$, $i \neq j$.

Input Format:

First Line n - Number of elements in an array

Next n Lines - N elements in the array

k - Non - Negative Integer

Output Format:

1 - If pair exists

0 - If no pair exists

Explanation for the given Sample Testcase:

YES as $5 - 1 = 4$

So Return 1.

For example:

Input	Result
3 1 3 5 4	1

Answer: (penalty regime: 0 %)

```
1 #include <stdio.h>
2
3 int main() {
4     int n;
5     scanf("%d", &n);
6     int arr[n];
7     for (int i = 0; i < n; i++)
8         scanf("%d", &arr[i]);
9
10    int k;
11    scanf("%d", &k);
12
13    int i = 0, j = 1;
14    while (i < n && j < n) {
15        int diff = arr[j] - arr[i];
16        if (diff == k && i != j) {
17            printf("1\n");
18            return 0;
19        } else if (diff < k) {
20            j++;
21        } else {
22            i++;
23        }
24    }
25    printf("0\n");
26    return 0;
27 }
28
```

	Input	Expected	Got	
✓	3 1 3 5 4	1	1	✓
✓	10 1 4 6 8 12 14 15 20 21 25	1	1	✓

	1			
✓	10 1 2 3 5 11 14 16 24 28 29 0	0	0	✓
✓	10 0 2 3 7 13 14 15 20 24 25 10	1	1	✓

Passed all tests! ✓

Correct

Marks for this submission: 1.00/1.00.

[Finish review](#)

[Back to Course](#)