


**S.No:** 421915**Programe** : B.TECH**Hall Ticket No:** 20A91A03D7**Name** : SANAPALLI VIJAY**Examination** : III SEMESTER SUPPLEMENTARY (AR20)**Month-Year** : Nov/Dec 2025**Branch** : MECHANICAL ENGINEERING**Course Code** : 201BS3T12**Course Name** : Integral transforms & applications of Partial Differential Equations**Date of Exam** : 18-11-2025

Signature of the Controller of Examination

Signature of the Student with date

Signature of the Invigilator with date

Serial No.
Last Page Written

19

INSTRUCTIONS TO STUDENTS

1. The Answer Booklet contains 36 pages Ensure all the 36 pages are in proper order.
2. Student must verify the details of particulars in the PART - 1 i.e Name, Hall Ticket No., Examination, Course Name etc.,
3. In case of any deviation in the above or if the PART-1 is torn / damaged, report to the invigilator and return the damaged booklet.
4. You are prohibited from writing on or tampering the PART-1 except affixing the signature and serial number of last page written in the space provided.
5. Student is prohibited from:
 - i. Writing their Hall Ticket No. and name in any part of the answer booklet.
 - ii. Addressing the examiner in any manner whatsoever in the answer booklet. If they do so, their script will not be valued.
 - iii. Writing religious symbols.
 - iv. Either seeking or providing any assistance to the fellow students in the examinations.
 - v. Possessing a manuscript or a printed matter, in any form, in the examination hall.
 - vi. Bringing Mobile Phones / Cameras/ Bluetooth Devices / Programmable calculators etc.
 - vii. Violation of the above mentioned instructions will be viewed as a case of malpractice, which is punishable offence.
6. Before beginning to answer any question, the students should write the correct number of that question in the margin provided. Answers written at different places for the same question will not be valued.
7. Students should write the answers, within the margins provided on both sides of the paper and on all the lines of each page. It is not necessary to begin each answer in a fresh page. Answers must be legibly written with blue or black pen.
8. Do not write anything except Question Number in the margin.
9. No loose sheets of papers will be allowed in the examination hall. No paper must be detached from or attached to the answer booklet except graph sheet.
10. Strike off all unused pages.
11. **NO ADDITIONAL ANSWER BOOKLETS/SHEETS WILL BE SUPPLIED.**



SPACE FOR ROUGH WORK



UNIT IQ.1

Given data

$$f(x) = e^{-x}$$

$$f(x) = \frac{a_0}{2} + \sum_{n=1}^{\infty} a_n \sin nx + \sum_{n=1}^{\infty} b_n \cos nx.$$

$$f(x) = \frac{a_0}{2}$$

$$f(x) = e^{-x}$$

$$f = f(x)$$

$$f = \frac{2\pi}{2}$$

$$f = 2\pi$$

$$f(x) = 2\pi \rightarrow (x)$$

$$f(0) = e^{-x}$$

$$f = e^{-0}$$

$$f = \frac{\cos nx}{2\pi}$$

$$f = e^{-0}$$

$$f(x) \sin nx$$

$$\sin \int_0^{\pi} a_n \cos nx$$

$$\sin \int_0^{2\pi} \cos bx.$$



$$\sin = \frac{a + \cos nx}{2}$$

$$\sin = -\cos nx$$

$$f(x) = 1 - \cos nx$$

$$\boxed{f(x) \sin nx} \rightarrow (2)$$

$$f(x) = b_n \cos nx$$

$$\sin b_n - \frac{\cos nx}{2}$$

$$\sin \int_0^{2\pi} \frac{\cos nx}{2}$$

$$\sin \int_0^{2\pi} \frac{\cos nx}{2}$$

$$\sin \int_0^{\pi} \cos nx$$

$$\sin \int_0^{\pi} \cos nx$$

$$\boxed{f(x) = A \cos nx} \rightarrow (3)$$

$$\text{eqn (1) + (2) + (3)}$$

$$f(x) = \frac{a_0}{2}$$

$$f(x) = a_n \sin nx$$

$$f(x) = b_n \sin nx$$

$$\boxed{f(x) = \frac{a_0}{2} + \sum_{n=1}^{\infty} a_n \sin nx + \sum_{n=1}^{\infty} b_n \sin nx} u.$$



2bA

Given data.

$$f(x) = x.$$

Half-range of Fourier Cosine series,
is the interval $(0, \pi)$.

$$f(x) = \frac{a_0}{2} + \sum_{n=1}^{\infty} a_n \sin nx + \sum_{n=1}^{\infty} b_n \cos nx.$$

Fourier series.

$$f(x) = x.$$

$$f(x) = \frac{2x}{\pi}$$

$$f(x) = 2$$

$$f(x) = x.$$

$$f(0) = x.$$

$$\boxed{f=0}$$

$$a_n \sin nx = f(x)$$

$$f(x) = a_n \sin nx.$$

$$f(x) = \int_0^{\pi} a_n \sin nx$$

$$f(x) = \int_0^{\pi} a_n \sin nx.$$

$$\boxed{f(x) = a_n \sin nx.}$$



Q.No.

$$b_n \cos nx = f(x)$$

$$f(x) = b_n \cos nx$$

$$F(x) = \int_{-\pi}^{\pi} b_n \cos nx.$$

$$F(x) = \int_{-\pi}^{\pi} \frac{1}{\sin nx}$$

$$f(x) = \frac{1}{1 - \cos^2 x}.$$

$$f(x) = b_n \cos nx$$

$$f(x) = x.$$

interval $(0, 2)$.

so it is proved.



UNIT-IIQ.10

Given data.

$$f(x) = \begin{cases} x^2 & \text{if } |x| < a. \\ 0 & \text{if } |x| > a. \end{cases}$$

Fourier transform.

$$f(x) = \frac{a_0}{2} + \sum_{n=1}^{\infty} a_n \sin nx + \sum_{n=1}^{\infty} b_n \cos nx.$$

$$f(x) = x^2$$

$$f'(x) = 2x$$

$$f(x) = \frac{2x^2}{2}$$

$$f(x) = \frac{4x^2}{2}$$

$$f'(x) = 2x$$

$$f''(x) = 2$$

$$f'' = 2$$

$$f = 4$$

$$f(x) = |x| < a$$

$$f(x) = 0$$

$$f'(x) = 0$$

$$f''(x) = 0$$

$$f(x) = \begin{cases} |x| & |x| < a \\ 0 & |x| > a. \end{cases}$$



4bA

Given data

$$f(x) = \begin{cases} -x & x < c \\ \pi - x & x \geq c \end{cases}$$

$$f(x) = \frac{a_0}{2} + \sum_{n=1}^{\infty} a_n \sin nx + \sum_{n=1}^{\infty} b_n \sin nx.$$

$$f(x) = -x$$

$$f(x) = \frac{a_0}{2} + a_n \sin nx.$$

$$f(x) = \int_0^x \sin nx.$$

$$f(x) = \int_0^x 1 - \cos^2 nx.$$

$$f(x) = \sin nx$$

$$\sin nx = -x.$$

$$\boxed{f(x) = -x}$$

$$f(x) = \pi - x.$$

$$f(x) = \int_{-x}^{\pi} \sin nx$$

$$f(x) = \int_{-x}^{\pi} 1 - \sin nx$$



$$f(x) = \int_{-x}^{\pi} 2 \sin nx.$$

$$\cancel{f(x) = 2\pi}$$

$$f(x) = \int_x^{\pi} \sin nx.$$

$$f(x) = \int_x^{2\pi} \frac{1}{\sin nx}.$$

$$f(x) = \cancel{2\pi} \int_x^{\pi} \frac{\sin x}{x}$$

$$f(x) = \sin nx.$$

$$\boxed{f(x) = \pi - x}$$

So, it is proved it.



UNIT-IIIQ. No. A

Given.

$$L \{ t^2 e^{3t} \sin 2t \}$$

$$= L \{ t^2 e^{3t} \}$$

$$= L \{ 1 - \cos + \sin 2x \}$$

$$= L \left\{ \frac{1 - \cos}{2x} + \sin 2x \right\}$$

$$= L \left\{ \frac{1 - \cos x}{2x} + \frac{1}{\sin 2x} \right\}$$

$$= L \left\{ \frac{1 - \cos x + \sin 2x}{2x + 1} \right\}$$

$$= L \left\{ \frac{\cos x t}{2x} \right\}$$

$$= L \{ t^2 e^{3t} \}$$

$$= L \{ \sin 2t \}$$

$$= L \left\{ \frac{1}{\cos 2t} \right\}$$

$$= L \{ t^2 e^{3t} \sin 2t \}$$



Q.No.

given

$$\int_0^{\infty} \frac{\cos 6t - \cos 4t}{t} dt.$$

$$\int_0^{\infty} \frac{\cos 6t}{t} dt$$

$$\int_0^{\infty} \cos 6t \cdot dt$$

$$\int_0^{\infty} \frac{1}{\sin 6} \cdot dt$$

$$\int_0^{\infty} \cos 6 \cdot dt$$

$$\int_0^{\infty} 6 \cos^2 \cdot dt$$

$$\int_0^{\infty} \frac{\cos 6t}{t} + \frac{\cos 2t}{2} dt.$$

$$\int_0^{\infty} \frac{\cos 6t + \cos 2t}{2t} dt$$

$$\int_0^{\infty} \frac{\cos 6 + (\cos 2t)^2}{2t} dt$$

$$\int_0^{\infty} \frac{\cos 6t - \cos 2t}{2} dt.$$



UNIT - IV7aP

Given

$$\mathcal{L}^{-1} \left\{ \frac{2s-3}{s^2+4s+13} \right\}$$

$$\frac{A}{s^2} + \frac{B}{4s} + \frac{C}{13}$$

$$\mathcal{L}^{-1} \left\{ \frac{2s-3}{s^2} \right\}$$

$$\mathcal{L}^{-1} \left\{ \frac{2s-3}{2s} \right\}$$

$$\mathcal{L}^{-1} \{ s-3 \}$$

$$\frac{B}{4s}$$

$$\mathcal{L}^{-1} \left\{ \frac{2s-3}{4s} \right\}$$

$$\mathcal{L}^{-1} \left\{ \frac{2s-3}{4s} \right\}$$

$$\mathcal{L}^{-1} \left\{ \frac{s}{2s-3} \right\}$$

$$\mathcal{L}^{-1} \left\{ \frac{1}{2s-3} \right\}$$



$$L^{-1} \left\{ \frac{2s-3}{s} \right\}$$

$$L^{-1} \{ 2s-3 \}$$

$$L^{-1} \left\{ \frac{2s-3}{13} \right\}$$

$$L^{-1} \left\{ \frac{2s-3}{13} \right\}$$

$$L^{-1} \{ 2-3 \}$$

$$L^{-1} \left\{ \frac{2s-3}{s^2+4s+13} \right\} \quad 4.$$



76A

Given

$$\bar{L}^{-1} \left\{ \frac{s^2}{(s^2+4)(s^2+9)} \right\}$$

$$\bar{L}^{-1} \left\{ \frac{s^2}{(s^2+4) * (s^2+9)} \right\}$$

$$\bar{L}^{-1} \left\{ \frac{s^2}{2s^2+9s+4s+36} \right\}$$

$$\bar{L}^{-1} \left\{ \frac{s^2}{11s^2+4s+36} \right\}$$

$$\bar{L}^{-1} \left\{ \frac{s^2}{15s^2+4s+36} \right\}$$

$$\bar{L}^{-1} \left\{ \frac{s^2}{15s^2+4s+36} \right\}$$

$$\frac{A}{15s^2} + \frac{B}{4s} + \frac{C}{36}$$

$$\bar{L}^{-1} \left\{ \frac{s^2}{15s^2} \right\}$$



$$\bar{L}^1 \left\{ \frac{s^\vee}{u s} \right\}$$

$$\bar{L}^1 \left\{ \frac{s}{u} \right\}$$

$$\bar{L}^1 \left\{ \frac{s^\vee}{3s^4} \right\}$$

$$\bar{L}^1 \left\{ \frac{s}{1s} \right\}$$

$$\bar{L}^1 \left\{ \frac{s^\vee}{(s^2+4)(s^2+9)} \right\} \cdot u,$$



UNIT-VIQ. No.

Given

$$\frac{du}{dx} = \frac{2du}{dt} + u$$

$$\therefore u(x, 0) = 6e^{-3x}$$

$$\frac{du}{dx} = \frac{du}{dy}$$

$$\frac{du}{dx} = \frac{du}{dt}$$

$$\frac{du}{dx} = \frac{2du}{dt}$$

$$\frac{du}{dx} = (x)$$

$$\frac{du}{dx} = \frac{2du \times 2dt \times 2dy}{2du \times 2dt \times 2dy}$$

$$\frac{du}{dx} = \frac{2du}{dt}$$

$$\frac{du}{dx} = \frac{2du}{dt} + u$$

$$\frac{du}{dx} = 6e^{-3x}$$

$$\frac{du}{dx} = \frac{6e^{-3x}}{dt}$$



Q.No.

$$\frac{dy}{dx} = \frac{6e^{3x}}{dt}$$

$$\frac{dy}{dt} = \frac{6e^{3x}}{dt}$$

$$\frac{dy}{6e^{3x}} = dt$$

$$\frac{dy}{dt} = 6e^{3x}$$

$$\boxed{\frac{dy}{dx} = \frac{dy}{dt} + u} \quad u.$$



QbA

Given data.

$$\frac{\partial^2 y}{\partial t^2} = c^2 \frac{\partial^2 y}{\partial x^2}$$

$$\frac{\partial y}{\partial t} = \frac{\partial x}{\partial t}$$

$$\frac{\partial y}{\partial t} = \frac{\partial x}{\partial x}$$

$$\frac{\partial y}{\partial x} = 1$$

$$\frac{\partial x}{\partial y} = c^2$$

$$\frac{\partial^2 y}{\partial x^2} = c^2$$

$$\frac{\partial^2 y}{\partial x^2} = \frac{\partial^2 y}{\partial x^2}$$

$$\frac{\partial^2 y}{\partial x^2} = c^2 \frac{\partial^2 y}{\partial y^2}$$

$$\frac{\partial^2 y}{\partial x^2} = c^2 \frac{\partial^2 y}{\partial y^2}$$



$$c^2 = \frac{\partial^2 y}{\partial x^2}$$

$$\frac{\partial^2 y}{\partial x^2} = \sin n\pi x$$

$$\frac{\partial^2 y}{\partial x^2} = c^2 \frac{1}{\sin n\pi x}$$

$$\frac{\partial^2 y}{\partial x^2} = c^2 \frac{\partial^2 u}{\partial x^2}$$

$$\boxed{\frac{\partial^2 y}{\partial x^2} = c^2 \frac{\partial^2 u}{\partial x^2}} \quad u.$$



Q.No.



Q.No.



Q.No.



Q.No.



Q.No.



Q.No.



Q.No.



Q.No.



Q.No.



Q.No.



Q.No.



Q.No.



Q.No.



[illegible]

