



S.No: 421941



Programme : B.TECH

Hall Ticket No: 22A91A0314

Name : GANTHAKARI VENKATESH

Examination : III SEMESTER SUPPLEMENTARY
(AR20(R))

Month-Year : Nov/Dec 2025

Branch : MECHANICAL ENGINEERING

Course Code : 201BS3T12

Course Name : Integral Transforms And Applications Of Partial Differential Equations

Date of Exam : 18-11-2025

J. Pavan.

Signature of the Controller of Examination

G. Venkatesh 18/11/2025

Signature of the Student with date

M. S. Chaitanya 18/11/25

Signature of the Invigilator with date

Serial No.
Last Page Written

17

INSTRUCTIONS TO STUDENTS

1. The Answer Booklet contains 36 pages Ensure all the 36 pages are in proper order.
2. Student must verify the details of particulars in the PART - 1 i.e Name, Hall Ticket No., Examination, Course Name etc.,
3. In case of any deviation in the above or if the PART-1 is torn / damaged, report to the invigilator and return the damaged booklet.
4. You are prohibited from writing on or tampering the PART-1 except affixing the signature and serial number of last page written in the space provided.
5. Student is prohibited from:
 - i. Writing their Hall Ticket No. and name in any part of the answer booklet.
 - ii. Addressing the examiner in any manner whatsoever in the answer booklet. If they do so, their script will not be valued.
 - iii. Writing religious symbols.
 - iv. Either seeking or providing any assistance to the fellow students in the examinations.
 - v. Possessing a manuscript or a printed matter, in any form, in the examination hall.
 - vi. Bringing Mobile Phones / Cameras/ Bluetooth Devices / Programmable calculators etc.
 - vii. Violation of the above mentioned instructions will be viewed as a case of malpractice, which is punishable offence.
6. Before beginning to answer any question, the students should write the correct number of that question in the margin provided. Answers written at different places for the same question will not be valued.
7. Students should write the answers, within the margins provided on both sides of the paper and on all the lines of each page. It is not necessary to begin each answer in a fresh page. Answers must be legibly written with blue or black pen.
8. Do not write anything except Question Number in the margin.
9. No loose sheets of papers will be allowed in the examination hall. No paper must be detached from or attached to the answer booklet except graph sheet.
10. Strike off all unused pages.
11. **NO ADDITIONAL ANSWER BOOKLETS/SHEETS WILL BE SUPPLIED.**



SPACE FOR ROUGH WORK



Unit-V

Q. No. 1) Given

$$\frac{\partial u}{\partial x} = 2 \frac{\partial u}{\partial t} + u \longrightarrow \textcircled{1}$$

given

$$u(x, 0) = 6e^{-3x}$$

Let the

$$u(x, t) = XT$$

$$\frac{\partial u}{\partial x} = x'T, \quad \frac{\partial u}{\partial t} = xT'$$

$$u(x, t) = X(x), T(t)$$

when Eqn ① becomes the

$$x'T = 2xT' + XT$$

$$(x' - x)T = 2xT' \longrightarrow \textcircled{2} \quad K = \text{constant}$$

$$\frac{(x' - x)}{x} = \frac{2T'}{T} \Rightarrow \frac{(x' - x)}{x} = K \quad \& \quad \frac{2T'}{T} = K$$

Eqn

$$\frac{(x' - x)}{x} = K \text{ integration in}$$

$$\int \frac{(x' - x)}{x} dx$$

$$\log x = \log(K+1) + \log C, \Rightarrow \log e = \log(K+1) \log C,$$



$$\log e = C_1 e^{(k+1)}$$

$$x = C_1 e^{(k+1)}$$

$$\frac{2T'}{T} = k \text{ is in integration}$$

$$\log T = \log 2k/T + \log C_2$$

$$= \log e + \log x^{2k/2} + \log C_2$$

$$T = C_2 e^{2k/2}$$

$$\text{Also } u(x, 0) = 6e^{-3x}$$

Compare in xT

$$u = C_1 e^{(k+1)x} C_2 e^{2k/2x}$$

$$u = C_1 C_2 e^{(k+1)x + 2k/2x}$$

$$\therefore C_1 C_2 = 6 \quad k = -4 \quad (k+1) = -4+1 = -3$$

$$\therefore u(x, 0) = 6e^{-3x} e^{-4/2x}$$

$$\text{and the eqn } \left[\frac{\partial u}{\partial x} = 2 \frac{\partial u}{\partial t} + u \quad u(x, 0) = 6e^{-3x} e^{-4/2x} \right]$$



96Sol) Given

$$\frac{\partial^2 y}{\partial t^2} = C^2 \frac{\partial^2 y}{\partial x^2} \quad \text{--- (1)}$$

$$u(x, t) = x \tau$$

$$u = x(x), \tau(t)$$

$$\frac{\partial^2 y}{\partial t^2} = x \tau'' \quad \frac{\partial^2 y}{\partial x^2} = x'' \tau$$

Egⁿ (1) Becomes the

$$x \tau'' = C^2 x'' \tau \quad \text{--- (2)}$$

$$x \tau'' = K \quad ; \quad C^2 x'' \tau = K$$

$\therefore K = \text{Constant}$

Integration on Both Sides

$$\begin{aligned} x \tau'' &= K \\ \int x \tau'' dx &= \int K dx \\ \log x \tau'' &= \log(K+1)x + \log C, \\ \log e &= \log(K+1)x + \log C, \\ \log e &= \log(K+1)x + \log C, \\ &= C_1 e^{(K+1)x} \\ x &= C_1 e^{(K+1)x} \end{aligned}$$

$$\begin{aligned} C^2 x'' \tau &= K \\ \int C^2 x'' \tau dx &= \int K dx \\ \log C^2 x'' \tau &= \log(K+1) + \log C_2 \\ \log e &= \log(K+1)/x + \log C_2 \\ \log e &= C_2 e^{(K+1)/C_2} \\ \tau &= C_2 e^{(K+1)x/C_2} \end{aligned}$$



$$\therefore K = 1, 2, 3, 4, \dots$$

$$\text{At } K=3$$

$$X = C_1 e^{4x}$$

$$T = C_2 e^{4x/c^2}$$

~~Then~~ ~~(1)~~

$$XT = C_1 C_2 e^{4x} e^{4x/c^2}$$

$$C_1 C_2 = 8$$

$$XT = C_1 C_2 e^{4x} e^{4x/c^2}$$

$$u = 8e^{4x} e^{4x/c^2}$$

Then

$$\frac{\partial^2 y}{\partial t^2} = c^2 \frac{\partial^2 y}{\partial x^2}$$

$$c^2 = ? \quad c^2 = 4$$

$$c = 2$$

$$u(x, t) = 8e^{4x} e^{4x/c^2}$$

$$u(x, t) = 8e^{4x} e^{4x/2}$$

$$u(x, t) = 8e^{4x} e^{2x}$$



Unit - I

2aSo) Given

$$f(x) = e^{-x}, (0, 2\pi)$$

$$y = a_0 \sum_{n=1}^{\infty} \int_0^{2\pi} f(x) \cos nx dx + \int_0^{2\pi} f(x) b_n \sin nx dx \quad \text{--- (1)}$$

$$y = a_0 \sum_{n=1}^{\infty} a_n \cos nx dx + b_n \sin nx dx$$

$$a_0 = \frac{1}{\pi} \int_0^{2\pi} f(x) dx$$

$$a_n = \frac{1}{\pi} \int_0^{2\pi} f(x) \cos nx dx$$

$$b_n = \frac{1}{\pi} \int_0^{2\pi} f(x) \sin nx dx$$

$$\frac{x^{n-1}}{n-1}$$

$$a_0 = \frac{1}{\pi} \int_0^{2\pi} e^{-x} dx$$

$$= \frac{1}{\pi} \int_0^{2\pi} (e)^x dx$$

$$= \frac{1}{\pi} [e^x]_0^{2\pi}$$

$$= \frac{1}{\pi} [2\pi - 0]$$

$$a_0 = 2$$



$$a_n = \frac{1}{\pi} \int_0^{2\pi} f(x) \cos nx \, dx$$

$$a_n = \frac{1}{\pi} \int_0^{2\pi} e^{-x} \cos nx \, dx$$

$$a_n = \frac{1}{\pi} [1 - \sin x]_0^{2\pi}$$

$$a_n = \frac{1}{\pi} [(2\pi - 0) - (\sin 2\pi - 0)]$$

$$a_n = \frac{1}{\pi} (2\pi - \sin 2\pi)$$

$$a_n = \frac{2}{\pi} (\pi - \sin \pi)$$

$$= 2 (-1)$$

$$a_n = -2$$

$$b_n = \frac{1}{\pi} \int_0^{2\pi} f(x) \sin nx \, dx$$

$$= \frac{1}{\pi} \int_0^{2\pi} e^{-x} \sin nx \, dx$$

$$= \frac{1}{\pi} [1 + \cos x]_0^{2\pi}$$

$$= \frac{1}{\pi} (2\pi + \cos 2\pi)$$

$$= \frac{1}{\pi} 2 [\pi + \cos \pi]$$

$$= \frac{2}{\pi} \pi (1 + 0) \quad b_n = \frac{2}{\pi}$$



$\therefore$ Eqⁿ ① in a_0, a_n, b_n

$$y = a_0 \sum_{n=1}^{\infty} a_n \cos nx dx + b_n \sin nx dx$$

$$y = 2 \sum_{n=1}^{\infty} -2 \cos nx dx + \frac{2}{\pi} \sin nx dx$$

2b Sol) Given That
 $f(x) = x$

given $(0, 2)$

$$y = a_0 \sum_{n=1}^{\infty} a_n \cos nx dx \longrightarrow \textcircled{1}$$

$$a_0 = \frac{1}{\pi} \int_0^2 f(x) dx \longrightarrow \textcircled{2}$$

$$a_n = \frac{1}{\pi} \int_0^2 f(x) \cos nx dx \longrightarrow \textcircled{3}$$

$$a_0 = \frac{1}{\pi} \int_0^2 x dx$$

$$a_0 = \frac{1}{\pi} [x^2]_0^2 dx$$

$$a_0 = \frac{1}{\pi} [2 - 0]$$

$$a_0 = \frac{2}{\pi}$$



$$a_n = \frac{1}{\pi} \int_0^2 f(x) \cos nx \, dx$$

$$= \frac{1}{\pi} \int_0^2 x \cos nx \, dx$$

$$= \frac{1}{\pi} \left[1 - \sin x \right]_0^2$$

$$= \frac{1}{\pi} [2 - \sin 2x]$$

$$= \frac{2}{\pi} [1 - \sin]$$

$$a_n = \frac{2}{\pi} \cos$$

$$\therefore a_0, a_n \text{ in Eqn. (i)}$$

$$y = a_0 \sum_{n=1}^{\infty} a_n \cos nx \, dx$$

$$= \frac{2}{\pi} \sum_{n=1}^{\infty} \frac{2}{\pi} \cos(\cos nx) \, dx$$

$$\boxed{\sum_{n=1}^{\infty} 2 \cos nx \, dx}$$



Unit - III

Q.68) Given

$$L \{ t^2 e^{-3t} \sin 2t \}$$

$$L \{ t^2 \} \quad L \{ e^{-3t} \sin 2t \}$$

$$L \{ x^n \} = \frac{n!}{(s-a)^{n+1}}$$

$$L \{ e^{-at} \sin bt \} = \frac{s-a}{(s-a)^2 + b^2}$$

$$L \{ t^2 \} = \frac{2!}{(s-a)^3}$$

$$L \{ e^{-3t} \sin 2t \} = \frac{s-3}{(s+3)^2 + 2^2}$$

$$= \frac{2!}{(s-a)^3}$$

$$= \frac{s-3}{s^2 + 6s + 13}$$

$$= \frac{2}{(3+3)^3}$$

$$= \frac{s-3}{s^2 + 6s + 13}$$

$$= \frac{2}{(3)^3}$$

in quadratic eqn

$$s^2 + 6s + 13$$

$$L \{ t^2 \} = 2.4486.5$$

$$= \frac{-b \pm \sqrt{b^2 - 4ac}}{2a}$$

$$= \frac{-36 \pm \sqrt{36 - 4(1)(13)}}{2(1)}$$

$$s = \frac{-36 \pm \sqrt{36 - 4(1)(13)}}{2(1)}$$



$$S = \frac{-36 \pm \sqrt{36 - 52}}{2}$$

$$= \frac{-36 \pm \sqrt{16}}{2} = \frac{-36 \pm 2\sqrt{4}}{2} = \frac{-36 \pm 2 \cdot 2}{2}$$

$$S = 34 \quad / S = -34$$

Ques) Given

$$= \int_0^{\infty} \frac{\cos 6t - \cos 4t}{t} dt$$

$$= \int_0^{\infty} \frac{\cos 6t}{t} dt - \int_0^{\infty} \frac{\cos 4t}{t} dt$$

$$L\{\cos at\} = \frac{s}{s^2 + a^2}$$

$$= \int_0^{\infty} \frac{\cos 6t}{t} dt$$

$$= \left[\frac{-\sin 6t}{t} \right]_0^{\infty} = \frac{-\sin 6[\infty - 0]}{[\infty - 0]}$$



$$= -\sin 6$$

$$\mathcal{L}\{\cos at\} = \frac{s}{s^2 - a^2}$$

$$-\int_0^{\infty} \frac{\cos 4t}{t} dt = \left[\frac{-\sin 4t}{t} \right]_0^{\infty}$$

$$= \frac{-\sin 4(\infty - 0)}{\infty - 0}$$

$$= -\sin 4$$

$$\mathcal{L}\{\cos at\} = \frac{s}{s^2 - a^2}$$

$$\frac{s}{s^2 - 6^2} - \frac{s}{s^2 - 4^2}$$

$$\therefore \left[= \frac{4}{s^2 - 12} \right]$$



Unit - IV8a5a) ~~Given~~

~~$$L^{-1} \left\{ \frac{s}{(s^2 + a^2)^2} \right\}$$~~

~~$$L^{-1} \left\{ \frac{s}{(s^2 + a^2)^2} \right\} = L^{-1} \{ \cosh at \}$$~~

Unit - IV7b5a) ~~Given~~

~~$$L^{-1} \left\{ \frac{s^2}{(s^2 + 4)(s^2 + 9)} \right\}$$~~

~~$$L^{-1} \left\{ \frac{s^2}{(s^2 + 4)} \right\} \quad L^{-1} \left\{ \frac{s^2}{(s^2 + 9)} \right\}$$~~

~~$$L^{-1} \left\{ \frac{s}{(s^2 + a^2)} \right\} = L^{-1} \{ \cosh at \}$$~~

In Convolution theorem

$$= \{ \cosh 2t \cdot \cosh 3t \}$$

$$P(s) \cdot Q(s) = F(s) \cdot G(s)$$



$$\bar{F}(s) = \cosh 2t$$

$$\bar{G}(s) = \cosh 3t$$

$$L^{-1} \left\{ \left[\frac{s^2}{s^2 + 4} \right] \left[\frac{s^2}{s^2 + 9} \right] \right\}$$

$$\bar{f}(t) \cdot \bar{g}(t)$$

$$[\cosh 2t \cdot \cos 3t]$$

7980) Given

$$L^{-1} \left\{ \frac{2s - 3}{s^2 + 4s + 13} \right\}$$

$$L^{-1} \left[\frac{(s-a)}{(s+a)^2 + b^2} \right] \Rightarrow \frac{1}{\sqrt{a^2 + b^2}} e^{-at} \sin bt$$

$$\therefore L^{-1} \left\{ e^{-at} \sin bt \right\}$$



Unit - II

3asol) Given

$$f(x) = \int_0^{\infty} \frac{\cos \lambda x}{1 + \lambda^2} d\lambda = \frac{\pi}{2} e^{-x}, \quad x \geq 0$$

$$y = \frac{a_0}{2} \sum_{n=1}^{\infty} [a_n \cos nx dx + b_n \sin nx dx] \sin nx dx \quad \text{--- (1)}$$

$$= \left[a_0 + \sum_{n=1}^{\infty} a_n \cos nx dx + b_n \sin nx dx \right] \sin nx dx$$

$$a_0 = \frac{1}{\pi} \int_0^x f(x) dx$$

$$a_n = \frac{1}{\pi} \int_0^x f(x) \cos nx dx$$

$$b_n = \frac{1}{\pi} \int_0^x f(x) \sin nx dx$$

$$a_0 = \frac{1}{\pi} \int_0^{\infty} \frac{\cos px}{1 + p^2} dp$$

$$= \frac{1}{\pi} \left[\frac{\cos p(x)}{1 + p^2} \right]_0^{\infty}$$

$$= \frac{1}{\pi} \left[\cos p(x) / 1 + p^2 \right]$$



$$a_0 = \int_0^x f(x) dx \quad \int_0^{\infty} \frac{\cos px}{1+p^2} dp \rightarrow (2)$$

$$\int_0^x \cos px dx = \int_0^x \frac{1}{1+p^2} dx$$

in the form of the

$$a_n = \frac{\pi}{2} \int_0^{\infty} \cos px dx = \frac{\pi}{2} [e]_0^{\infty} = e^{-x}$$

$$a_n = \frac{\pi}{2} e^{-x}$$

$$\therefore p = \lambda$$

in the eqⁿ (2) in $p = \lambda$

$$\int_0^{\infty} \frac{\cos \lambda x}{1+\lambda^2} d\lambda = a_n$$

$$\therefore a_n = \frac{\pi}{2} e^{-x}$$

$$\left[\therefore \int_0^{\infty} \frac{\cos \lambda x}{1+\lambda^2} d\lambda = \frac{\pi}{2} e^{-x} \right]$$



Q.No.



Q.No.



Q.No.



Q.No.



Q.No.



Q.No.



Q.No.



Q.No.



Q.No.



Q.No.



Q.No.



Q.No.



Q.No.



Q.No.



Q.No.



Q.No.



Q.No.

$$\begin{array}{r} 37 \\ 37 \overline{) 219} 9 \\ 132 6 \\ \underline{37} 9103 \\ \underline{3987} 24428.5 \\ \underline{214} 108 \\ \underline{108} 08 \\ \underline{08} 76 \\ \underline{76} 13 \\ \underline{13} 12 \\ \underline{12} 10 \end{array}$$

$$\begin{array}{r} 4 \\ \hline S^2 + 12 \end{array}$$





