

2235. Add Two Integers

Solved

Easy

Topics

Companies

Given two integers `num1` and `num2`, return *the sum of the two integers*.

Example 1:

Input: `num1 = 12, num2 = 5`
Output: `17`
Explanation: `num1` is 12, `num2` is 5, and their sum is `12 + 5 = 17`, so 17 is returned.

Example 2:

Input: `num1 = -10, num2 = 4`
Output: `-6`
Explanation: `num1 + num2 = -6`, so `-6` is returned.

Constraints:

- `-100 <= num1, num2 <= 100`



Seen this question in a real interview before? 1/6

Yes

No

Code

Java Auto

```
1 class Solution {
2     public int sum(int num1, int num2) {
3         return num1+num2;
4     }
5 }
```

Saved

Ln 3, Col 26

Testcase Test Result

Accepted Runtime: 0 ms

Case 1

Case 2

Input

num1 =
12

num2 =
5

Output

Chrome

FileEditViewHistoryBookmarksProfilesTabWindowHelp

▼

Subl ×

Best ×

Onli ×

IS Myn ×

my WeF ×

Vish ×

æ You ×

æ Palin ×

Settl ×

Add ×

Subl ×

Subl ×

+

Ask Gemini

←

→

↻

🔍 leetcode.com/problems/subtract-the-product-and-sum-of-digits-of-an-integer/description/

☆

Error

⋮

🏠

☰ Problem List

⏪

⏩

🔗

🔒

▶

📤 Submit

📄

🌟

🔌

⚙️

🔥 0

🕒

👤

🖱️

Premium

📄 Description

📝 Note ×

📖 Editorial

🧪 Solutions

🕒 Submissions

1281. Subtract the Product and Sum of Digits of an Integer

Solved

Easy

🔖 Topics

🏢 Companies

💡 Hint

Given an integer number `n`, return the difference between the product of its digits and the sum of its digits.

Example 1:

Input: `n = 234`

Output: 15

Explanation:

Product of digits = $2 * 3 * 4 = 24$

Sum of digits = $2 + 3 + 4 = 9$

Result = $24 - 9 = 15$

Example 2:

Input: `n = 4421`

Output: 21

Explanation:

Product of digits = $4 * 4 * 2 * 1 = 32$

Sum of digits = $4 + 4 + 2 + 1 = 11$

Result = $32 - 11 = 21$

Constraints:

- `1 <= n <= 10^5`

</> Code

Java ▼ 🔒 Auto

☰ 📖 {} ↶ ↷ ↵ ↶ ↷

```
1 class Solution {
2     public int subtractProductAndSum(int n) {
3         int sum = 0;
4         int product = 1;
5
6         while (n > 0) {
7             int digit = n % 10;
8             sum += digit;
9             product *= digit;
10            n /= 10;
11        }
12        return product - sum;
13    }
14 }
```

SavedLn 1, Col 1

☑️ Testcase

>_ Test Result

You must run your code first

👍 2.9K

👤

💬 94

☆

🔗

🔍

17 Online

ChromeFileEditViewHistoryBookmarksProfilesTabWindowHelp

SubtracBest ApOnlineIS Myntrammy WeForSVishwalYou seePalindroSettingAdd TwSubtracSubtracFin x+

leetcode.com/problems/find-numbers-with-even-number-of-digits/description/

☆Error

Ask Gemini

Problem List<>↔

🔒▶️📁 Submit📄🌟

🧩⚙️🔥 0🕒👤🌟 Premium

DescriptionAccepted xNote xEditorialSolutionsSubmissions

1295. Find Numbers with Even Number of Digits

Solved✔️

EasyTopicsCompaniesHint

Given an array `nums` of integers, return how many of them contain an **even number** of digits.

Example 1:

Input: `nums = [12,345,2,6,7896]`

Output: 2

Explanation:

12 contains 2 digits (even number of digits).

345 contains 3 digits (odd number of digits).

2 contains 1 digit (odd number of digits).

6 contains 1 digit (odd number of digits).

7896 contains 4 digits (even number of digits).

Therefore only 12 and 7896 contain an even number of digits.

Example 2:

Input: `nums = [555,901,482,1771]`

Output: 1

Explanation:

Only 1771 contains an even number of digits.

Constraints:

- `1 <= nums.length <= 500`
- `1 <= nums[i] <= 105`

</> Code

Java🔒 Auto

```
1 class Solution {
2     public int findNumbers(int[] nums) {
3         int count=0;
4         for(int n:nums){
5             int digits=0;
6             while (n>0){
7                 digits++;
8                 n/=10;
9             }
10            if (digits%2==0){
11                count++;
12            }
13        }
14        return count;
15    }
16 }
```

SavedLn 14, Col 21

☑️ Testcase>_ Test Result

Accepted

Runtime: 0 ms

☑️ Case 1

☑️ Case 2

Input

nums =
[12, 345, 2, 6, 7896]

Output

2

Expected

👍 3K👎🗨 210🌟🔗🔍

25 Online

ChromeFileEditViewHistoryBookmarksProfilesTabWindowHelp

SubtraBest AOnlineIS Myntrmy WeForVishwae You seae PalindSettingsAdd TSubtraSubtraFind NCox

Ask Gemini

leetcode.com/problems/convert-the-temperature/description/

☆Error

Problem List

Submit

Premium

DescriptionAccepted ×Note ×EditorialSolutionsSubmissions

2469. Convert the Temperature

Solved

EasyTopicsCompaniesHint

You are given a non-negative floating point number rounded to two decimal places `celsius`, that denotes the **temperature in Celsius**.

You should convert Celsius into **Kelvin** and **Fahrenheit** and return it as an array `ans = [kelvin, fahrenheit]`.

Return *the array* `ans`. Answers within `10-5` of the actual answer will be accepted.

Note that:

- `Kelvin = Celsius + 273.15`
- `Fahrenheit = Celsius * 1.80 + 32.00`

Example 1:

Input: `celsius = 36.50`

Output: `[309.65000,97.70000]`

Explanation: Temperature at 36.50 Celsius converted in Kelvin is 309.65 and converted in Fahrenheit is 97.70.

Example 2:

Input: `celsius = 122.11`

Output: `[395.26000,251.79800]`

Explanation: Temperature at 122.11 Celsius converted in Kelvin is 395.26 and converted in Fahrenheit is 251.798.

Code

JavaAuto

```
1 class Solution {
2     public double[] convertTemperature(double celsius) {
3         double kelvin=celsius+273.15;
4         double fahrenheit=celsius*1.80+32.00;
5         return new double[]{kelvin,fahrenheit};
6     }
7 }
```

SavedLn 5, Col 47

TestcaseTest Result

Accepted

Runtime: 0 ms

Case 1Case 2

Input

celsius =

36.50

Output

[309.65000,97.70000]

Expected

764104

9 Online

[Description](#) |
 [Note](#) X |
 [Editorial](#) |
 [Solutions](#) |
 [Submissions](#)

1512. Number of Good Pairs

Easy Topics Companies Hint

Given an array of integers `nums`, return *the number of good pairs*.

A pair (i, j) is called *good* if `nums[i] == nums[j]` and $i < j$.

Example 1:

```
Input: nums = [1,2,3,1,1,3]
Output: 4
Explanation: There are 4 good pairs (0,3), (0,4), (3,4), (2,5) 0-indexed.
```

Example 2:

```
Input: nums = [1,1,1,1]
Output: 6
Explanation: Each pair in the array are good.
```

Example 3:

```
Input: nums = [1,2,3]
Output: 0
```

Constraints:

- `1 <= nums.length <= 100`
- `1 <= nums[i] <= 100`

 Code

Java Auto

```

1  class Solution {
2      public int numIdenticalPairs(int[] nums) {
3          int count=0;
4          for(int i=0;i<nums.length;i++){
5              for(int j=i+1;j<nums.length;j++){
6                  if (nums[i]==nums[j]){
7                      count++;
8                  }
9              }
10         }
11         return count;
12     }
13 }

```

Saved

Ln 5, Col 24

☑ Testcase | >_ Test Result

Accepted Runtime: 0 ms

✓ Case 1 ✓ Case 2 ✓ Case 3

Input

```
nums =  
[1,2,3,1,1,3]
```

Output

4

Expected

2520. Count the Digits That Divide a Number

Easy Topics Companies Hint

An integer `val` divides `nums` if `nums % val == 0`.

Input: num = 7

Output: 1

Explanation: 7 divides itself, hence the answer is 1.

Input: num = 121

Output: 2

Explanation: 121 is divisible by 1, but not 2. Since 1 occurs twice as a digit, we return 2.

Input: num = 1248

Output: 4

Explanation: 1248 is divisible by all of its digits, hence the answer is 4.

- `1 <= num <= 109`
- `num` does not contain `0` as one of its digits.

● 8 Online

Java Auto

```
1 class Solution {
2     public int countDigits(int n) {
3         int cnt=0;
4         int div=1;
5         while(div<=n){
6             int ld=(n%(div*10))/div;
7             if(n%ld==0) cnt++;
8             div=div*10;
9         }
10        return cnt;
11    }
12 }
```

Saved

Ln 1, Col 1

☑ Testcase | ➤ Test Result

Accepted Runtime: 0 ms

✓ Case 1 ✓ Case 2 ✓ Case 3

Input

```
num =  
7
```

Output

1

Expected

ChromeFileEditViewHistoryBookmarksProfilesTabWindowHelp

Subtrac xBest Ap xOnline xIS Myntra xmy WeForS xVishwal xYou sea xPalindro xSettings xMaximu x

Ask Gemini

leetcode.com/problems/maximum-69-number/

☆Error

Problem List

Submit

Premium

Description

Note x

Editorial

Solutions

Submissions

1323. Maximum 69 Number

Solved

Easy

Topics

Companies

Hint

You are given a positive integer `num` consisting only of digits `6` and `9`.

Return the maximum number you can get by changing **at most one digit** (`6` becomes `9`, and `9` becomes `6`).

Example 1:

Input: `num = 9669`

Output: `9969`

Explanation:

Changing the first digit results in 6669.

Changing the second digit results in 9969.

Changing the third digit results in 9699.

Changing the fourth digit results in 9666.

The maximum number is 9969.

Example 2:

Input: `num = 9996`

Output: `9999`

Explanation:

Changing the last digit 6 to 9 results in the maximum number.

Example 3:

Input: `num = 9999`

Output: `9999`

Explanation:

It is better not to apply any change.

Code

JavaAuto

```
1 class Solution {
2     public int maximum69Number (int n) {
3         int diff=3,div=1,sum=0;
4         while(div<=n){
5             int ld=(n%(div*10))/div;
6             if(ld==6) sum=diff;
7             diff*=10;
8             div*=10;
9         }
10        return n+sum;
11    }
12 }
```

SavedLn 1, Col 1

TestcaseTest Result

AcceptedRuntime: 0 ms

Case 1

Case 2

Case 3

Input

num =
9669

Output

9969

Expected

3.3K428

4 Online

Easy Topics Companies Hint

- For example, given `num = 2932`, you have the following digits: two `2`'s, one `9` and one `3`. Some of the possible pairs `[new1, new2]` are `[22, 93]`, `[23, 92]`, `[223, 9]` and `[2, 329]`.

Example 1:

Output: 52

Example 2:

Output: 13

Constraints:

👍 1.5K 🗨️ 49 ⭐ 📌 ?

● 5 Online

Java Auto

Saved

Ln 4, Col 30

☑ Testcase | ➤ Test Result

Accepted Runtime: 2 ms

✓ Case 1 ✓ Case 2

Input

```
num =  
2932
```

Output

52

Expected

[Description](#) |
 [Note](#) × |
 [Editorial](#) |
 [Solutions](#) |
 [Submissions](#)

2485. Find the Pivot Integer

Easy Topics Companies Hint

Given a positive integer n , find the **pivot integer** x such that:

- The sum of all elements between 1 and x inclusively equals the sum of all elements between x and n inclusively.

Return *the pivot integer* `x`. If no such integer exists, return `-1`. It is guaranteed that there will be at most one pivot index for the given input.

Example 1:

Input: $n = 8$

Output: 6

Explanation: 6 is the pivot integer since: $1 + 2 + 3 + 4 + 5 + 6 = 6 + 7 + 8 = 21$.

Example 2:

Input: $n = 1$

Output: 1

Explanation: 1 is the pivot integer since: $1 = 1$.

Example 3:

Input: $n = 4$

Output: -1

Explanation: It can be proved that no such integer exist.

Contraintes.

 Code

Java Auto

```
1 class Solution {
2     public int pivotInteger(int n) {
3         int total=n*(n+1)/2;
4         int left=0;
5         for(int x=1;x<=n;x++){
6             left+=x;
7             if(left==total-left+x){
8                 return x;
9             }
10        }
11        return -1;
12    }
13 }
```

Saved

Ln 11, Col 19

☑ Testcase | ➤ Test Result

Accepted Runtime: 0 ms

✓ Case 1 ✓ Case 2 ✓ Case 3

Input

$$n =$$

8

Output

6

Expected