

Sherlock and Cost



In this challenge, you will be given an array B and must determine an array A . There is a special rule: For all i , $A[i] \leq B[i]$. That is, $A[i]$ can be any number you choose such that $1 \leq A[i] \leq B[i]$. Your task is to select a series of $A[i]$ given $B[i]$ such that the sum of the absolute difference of consecutive pairs of A is maximized. This will be the array's *cost*, and will be represented by the variable S below.

The equation can be written:

$$S = \sum_{i=2}^N |A_i - A_{i-1}|$$

For example, if the array $B = [1, 2, 3]$, we know that $1 \leq A[1] \leq 1$, $1 \leq A[2] \leq 2$, and $1 \leq A[3] \leq 3$. Arrays meeting those guidelines are:

```
[1,1,1], [1,1,2], [1,1,3]
[1,2,1], [1,2,2], [1,2,3]
```

Our calculations for the arrays are as follows:

```
|1-1| + |1-1| = 0 |1-1| + |2-1| = 1 |1-1| + |3-1| = 2
|2-1| + |1-2| = 2 |2-1| + |2-2| = 1 |2-1| + |3-2| = 2
```

The maximum value obtained is **2**.

Input Format

The first line contains the integer t , the number of test cases.

Each of the next t pairs of lines is a test case where:

- The first line contains an integer n , the length of B
- The next line contains n space-separated integers $B[i]$

Constraints

- $1 \leq t \leq 20$
- $1 \leq n \leq 10^5$
- $1 \leq B[i] \leq 100$

Output Format

For each test case, print the maximum sum on a separate line.

Sample Input

```
1
5
10 1 10 1 10
```

Sample Output

```
36
```

Explanation

The maximum sum occurs when $A[1]=A[3]=A[5]=10$ and $A[2]=A[4]=1$. That is

$$|1 - 10| + |10 - 1| + |1 - 10| + |10 - 1| = 36.$$