

Phosphorylation of Exo1 modulates homologous recombination repair of DNA double-strand breaks

Emma Bolderson¹, Nozomi Tomimatsu², Derek J. Richard¹, Didier Boucher¹, Rakesh Kumar³, Tej K. Pandita³, Sandeep Burma² and Kum Kum Khanna^{1,*}

¹Signal Transduction Laboratory, Queensland Institute of Medical Research, Brisbane, Queensland 4029, Australia, ²Department of Radiation Oncology, UT Southwestern Medical Center at Dallas, Dallas, TX 75390-9187 and ³Department of Radiation Oncology, Washington University School of Medicine, St Louis, MO 63108, USA

Received October 22, 2009; Revised November 18, 2009; Accepted November 24, 2009

ABSTRACT

DNA double-strand break (DSB) repair via the homologous recombination pathway is a multi-stage process, which results in repair of the DSB without loss of genetic information or fidelity. One essential step in this process is the generation of extended single-stranded DNA (ssDNA) regions at the break site. This ssDNA serves to induce cell cycle checkpoints and is required for Rad51 mediated strand invasion of the sister chromatid. Here, we show that human Exonuclease 1 (Exo1) is required for the normal repair of DSBs by HR. Cells depleted of Exo1 show chromosomal instability and hypersensitivity to ionising radiation (IR) exposure. We find that Exo1 accumulates rapidly at DSBs and is required for the recruitment of RPA and Rad51 to sites of DSBs, suggesting a role for Exo1 in ssDNA generation. Interestingly, the phosphorylation of Exo1 by ATM appears to regulate the activity of Exo1 following resection, allowing optimal Rad51 loading and the completion of HR repair. These data establish a role for Exo1 in resection of DSBs in human cells, highlighting the critical requirement of Exo1 for DSB repair via HR and thus the maintenance of genomic stability.

INTRODUCTION

DNA double-strand breaks (DSBs) can be induced by a variety of factors such as chemotherapeutic agents, ionising radiation (IR) and by the products of cellular metabolism, including replication fork collapse. In order to maintain genomic stability, cells possess a complex network of signalling pathways involved in the detection,

signalling and repair of DNA damage. Defects in these DNA repair pathways can lead to human genomic instability syndromes, with increased cancer susceptibility, neurological syndromes and immunodeficiency. The resection of DSBs to produce 3' single-stranded DNA (ssDNA) tracts is a critical step in the repair of DSBs by homologous recombination (1). The ssDNA at the break site is essential for activation of the ATR signalling cascade which re-enforces ATM-induced cell cycle checkpoints (2). The MRN (MRE11, Rad50 and NBS1) complex, in association with CtIP, has been previously reported to be important for DSB resection (3). However, recent reports clearly indicate that yeast MRX (Mre11, Rad50 and XRS1) is involved only in limited resection at the break sites while extensive resection requires additional, redundant nucleases such as Exonuclease 1 (Exo1) and/or DNA2 (4).

Exo1 was first identified in *Schizosaccharomyces pombe* as a nuclease that is induced during meiosis (5). Exo1 belongs to the RAD2 family of nucleases and possesses 5'–3' nuclease activity and 5'-flap endonuclease activity (6,7). Alternate splicing leads to two isoforms of Exo1 (a and b). The isoforms differ at the C-terminus, with Exo1b having an additional 48 amino acids. Exo1 is known to interact with several other proteins involved in replication and DNA repair including PCNA and mismatch repair (MMR) proteins (8). Exo1 is implicated in several DNA repair pathways including MMR, post-replication repair, meiotic and mitotic recombination (9–11). The involvement of Exo1 in DNA repair pathways including MMR suggests it may also be a target for mutation in tumorigenesis. Consistent with this, a cancer-prone phenotype can be observed in Exo1-deficient mice including increased susceptibility to lymphoma development (12). In addition, patients with atypical human non-polyposis colon cancer and other forms of colorectal cancer have been found to have

*To whom correspondence should be addressed. Tel: +61 7 3362 0338; Fax: +61 7 3362 0105; Email: kumkumk@qimr.edu.au

We have shown here that following DSB-induction Exo1 is initially phosphorylated by an ATM-dependent mechanism, however, at later time points Exo1 phosphorylation is independent of ATM suggesting that it is phosphorylated redundantly, possibly by ATR or DNA-PK. We show that while RPA34 recruitment to foci was normal in cells expressing the non-phosphorylatable mutant of Exo1 (S714A), Rad51 recruitment to foci was defective. This suggests that the phosphorylation event at S714 is not required for the generation of the ssDNA tracks but is possibly required to allow Rad51 loading. Interestingly we observed that cells expressing a phospho-mimic Exo1 (S714E) are deficient in generation of ssDNA/DNA resection as measured by the recruitment of RPA34 and Rad51 to foci. This phospho-mimic mutant also has impaired HR activity. At present we can only speculate as to the specific function of the phosphorylation event, however a recent paper examining four phosphorylation sites in yeast Exo1 suggested that phosphorylation probably inhibits the activity of Exo1 (35).

It is tempting to speculate that S714 phosphorylation is not required for the initiation of Exo1-mediated DNA resection, but rather that the phosphorylation is required to attenuate Exo1 activity and perhaps prevent the generation of substrates unsuitable for HR. Inactivation of a nuclease by phosphorylation after DNA damage is not without precedent. A recent paper has also shown that phosphorylation of Mre11 on SQ/TQ residues decreases its affinity for DNA, causes disassembly from chromatin and thereby results in its inactivation, it is possible that phosphorylation of Exo1 may function in a similar way (40). Our interpretations do not exclude additional functions of phosphorylated Exo1 in recombinational DNA repair. It is also possible that phosphorylation of Exo1 on S714 changes its binding partners. We have observed here that phosphorylation of Exo1 by ATM was dispensable for resection as revealed by normal RPA foci formation in non-phosphorylatable mutant (S714A) expressing cells; however these cells are likely to be defective in recombinational repair due to defective loading of Rad51. Indeed we also found that cells expressing the non-phosphorylatable mutant displayed defective *I-SceI*-induced HR. Taken together, our data suggests that the DNA substrates generated by action of non-phosphorylatable Exo1 cannot be used efficiently by Rad51 to form nucleoprotein filaments. Further studies will be required to address whether this is due to the over-resection by unregulated Exo1.

A recent study has shown that HR of IR-induced DSBs that arise during G2 is reduced in ATM-deficient cells (41). Since we have shown that ATM is required for the timely phosphorylation of Exo1 and that this phosphorylation event is required for the regulation of HR, this may further confirm a role for ATM in controlling recombinational repair of DSBs. However, the role of ATM in Exo1 phosphorylation appears to be redundant suggesting that other proteins such as ATR or DNA-PK can compensate albeit at a later time-point.

DNA DSBs induce a vast array of signaling networks, which can greatly influence DNA repair, cell cycle checkpoint activation and cell viability and therefore it is

understandable that multiple levels of regulation may exist to inactivate destructive proteins such as nucleases. Exo1 activity may be attenuated by ATM-dependent phosphorylation following its role in resection to prevent the destruction of substrates required for HR. Given the importance of accurate recombination in preserving the genome, our findings clearly illustrate the role of Exo1 in maintaining genomic stability.

SUPPLEMENTARY DATA

Supplementary Data are available at NAR Online.

FUNDING

The National Health and Research Council of Australia (442903 to K.K.K.); the National Institutes of Health (grant numbers CA129537 and CA123232 to T.K.P.) and National Aeronautics and Space Administration (grant number NNA05CS97G to S.B.). Funding for open access charge: National Health and Research Council of Australia.

Conflict of interest statement. None declared.

REFERENCES

- Mimitou, E.P. and Symington, L.S. (2009) Nucleases and helicases take center stage in homologous recombination. *Trends Biochem. Sci.*, **34**, 264–272.
- Shiotani, B. and Zou, L. (2009) Single-stranded DNA orchestrates an ATM-to-ATR switch at DNA breaks. *Mol. Cell*, **33**, 547–558.
- Sartori, A.A., Lukas, C., Coates, J., Mistrik, M., Fu, S., Bartek, J., Baer, R., Lukas, J. and Jackson, S.P. (2007) Human CtIP promotes DNA end resection. *Nature*, **450**, 509–514.
- Bernstein, K.A. and Rothstein, R. (2009) At loose ends: resecting a double-strand break. *Cell*, **137**, 807–810.
- Szankasi, P. and Smith, G.R. (1992) A DNA exonuclease induced during meiosis of *Schizosaccharomyces pombe*. *J. Biol. Chem.*, **267**, 3014–3023.
- Lee, B.I., Nguyen, L.H., Barsky, D., Fernandes, M. and Wilson, D.M. III (2002) Molecular interactions of human Exo1 with DNA. *Nucleic Acids Res.*, **30**, 942–949.
- Qiu, J., Qian, Y., Chen, V., Guan, M.X. and Shen, B. (1999) Human exonuclease 1 functionally complements its yeast homologues in DNA recombination, RNA primer removal, and mutation avoidance. *J. Biol. Chem.*, **274**, 17893–17900.
- Lee, S.D. and Alani, E. (2006) Analysis of interactions between mismatch repair initiation factors and the replication processivity factor PCNA. *J. Mol. Biol.*, **355**, 175–184.
- Fiorentini, P., Huang, K.N., Tishkoff, D.X., Kolodner, R.D. and Symington, L.S. (1997) Exonuclease I of *Saccharomyces cerevisiae* functions in mitotic recombination in vivo and in vitro. *Mol. Cell Biol.*, **17**, 2764–2773.
- Kirkpatrick, D.T., Ferguson, J.R., Petes, T.D. and Symington, L.S. (2000) Decreased meiotic intergenic recombination and increased meiosis I nondisjunction in exo1 mutants of *Saccharomyces cerevisiae*. *Genetics*, **156**, 1549–1557.
- Tsubouchi, H. and Ogawa, H. (2000) Exo1 roles for repair of DNA double-strand breaks and meiotic crossing over in *Saccharomyces cerevisiae*. *Mol. Biol. Cell*, **11**, 2221–2233.
- Wei, K., Clark, A.B., Wong, E., Kane, M.F., Mazur, D.J., Parris, T., Kolas, N.K., Russell, R., Hou, H. Jr, Kneitz, B. et al. (2003) Inactivation of Exonuclease 1 in mice results in DNA mismatch repair defects, increased cancer susceptibility, and male and female sterility. *Genes Dev.*, **17**, 603–614.
- Sun, X., Zheng, L. and Shen, B. (2002) Functional alterations of human exonuclease 1 mutants identified in atypical hereditary

- nonpolyposis colorectal cancer syndrome. *Cancer Res.*, **62**, 6026–6030.
14. Wu, Y., Berends, M.J., Post, J.G., Mensink, R.G., Verlind, E., van der Sluis, T., Kempinga, C., Sijmons, R.H., van der Zee, A.G., Hollema, H. *et al.* (2001) Germline mutations of EXO1 gene in patients with hereditary nonpolyposis colorectal cancer (HNPCC) and atypical HNPCC forms. *Gastroenterology*, **120**, 1580–1587.
 15. Schaezel, S., Kodandaramireddy, N.R., Ju, Z., Lechel, A., Stepczynska, A., Lilli, D.R., Clark, A.B., Rudolph, C., Kuhnel, F., Wei, K. *et al.* (2007) Exonuclease-1 deletion impairs DNA damage signaling and prolongs lifespan of telomere-dysfunctional mice. *Cell*, **130**, 863–877.
 16. Bolderson, E., Richard, D.J., Edelman, W. and Khanna, K.K. (2009) Involvement of Exo1b in DNA damage-induced apoptosis. *Nucleic Acids Res.*, **37**, 3452–3463.
 17. Tomimatsu, N., Mukherjee, B. and Burma, S. (2009) Distinct roles of ATR and DNA-PKcs in triggering DNA damage responses in ATM-deficient cells. *EMBO Rep.*, **10**, 629–635.
 18. Richard, D.J., Bolderson, E., Cubeddu, L., Wadsworth, R.I., Savage, K., Sharma, G.G., Nicolette, M.L., Tsvetanov, S., McIlwraith, M.J., Pandita, R.K. *et al.* (2008) Single-stranded DNA-binding protein hSSB1 is critical for genomic stability. *Nature*, **453**, 677–681.
 19. Uematsu, N., Weterings, E., Yano, K., Morotomi-Yano, K., Jakob, B., Taucher-Scholz, G., Mari, P.O., van Gent, D.C., Chen, B.P. and Chen, D.J. (2007) Autophosphorylation of DNA-PKcs regulates its dynamics at DNA double-strand breaks. *J. Cell Biol.*, **177**, 219–229.
 20. Khanna, K.K., Keating, K.E., Kozlov, S., Scott, S., Gatei, M., Hobson, K., Taya, Y., Gabrielli, B., Chan, D., Lees-Miller, S.P. *et al.* (1998) ATM associates with and phosphorylates p53: mapping the region of interaction. *Nat. Genet.*, **20**, 398–400.
 21. Young, D.B., Jonnalagadda, J., Gatei, M., Jans, D.A., Meyn, S. and Khanna, K.K. (2005) Identification of domains of ataxia-telangiectasia mutated required for nuclear localization and chromatin association. *J. Biol. Chem.*, **280**, 27587–27594.
 22. Li, Y., Bolderson, E., Kumar, R., Muniandy, P.A., Xue, Y., Richard, D., Seidman, M., Pandita, T.K., Khanna, K.K. and Wang, W. (2009) hSSB1 and hSSB2 form similar multi-protein complexes that participate in DNA damage response. *J. Biol. Chem.*, **284**, 23525–23531.
 23. Gupta, A., Sharma, G.G., Young, C.S., Agarwal, M., Smith, E.R., Paull, T.T., Lucchesi, J.C., Khanna, K.K., Ludwig, T. and Pandita, T.K. (2005) Involvement of human MOF in ATM function. *Mol. Cell Biol.*, **25**, 5292–5305.
 24. Gravel, S., Chapman, J.R., Magill, C. and Jackson, S.P. (2008) DNA helicases Sgs1 and BLM promote DNA double-strand break resection. *Genes Dev.*, **22**, 2767–2772.
 25. Rogakou, E.P., Pilch, D.R., Orr, A.H., Ivanova, V.S. and Bonner, W.M. (1998) DNA double-stranded breaks induce histone H2AX phosphorylation on serine 139. *J. Biol. Chem.*, **273**, 5858–5868.
 26. Chen, H.T., Bhandoola, A., Difilippantonio, M.J., Zhu, J., Brown, M.J., Tai, X., Rogakou, E.P., Brotz, T.M., Bonner, W.M., Ried, T. *et al.* (2000) Response to RAG-mediated VDJ cleavage by NBS1 and gamma-H2AX. *Science*, **290**, 1962–1965.
 27. Peterson, C.L. (2001) Chromatin: mysteries solved? *Biochem. Cell Biol.*, **79**, 219–225.
 28. Litman, R., Peng, M., Jin, Z., Zhang, F., Zhang, J., Powell, S., Andreassen, P.R. and Cantor, S.B. (2005) BACH1 is critical for homologous recombination and appears to be the Fanconi anemia gene product FANCI. *Cancer Cell*, **8**, 255–265.
 29. Pierce, A.J., Johnson, R.D., Thompson, L.H. and Jasin, M. (1999) XRCC3 promotes homology-directed repair of DNA damage in mammalian cells. *Genes Dev.*, **13**, 2633–2638.
 30. Zhou, B.B. and Elledge, S.J. (2000) The DNA damage response: putting checkpoints in perspective. *Nature*, **408**, 433–439.
 31. Mimitou, E.P. and Symington, L.S. (2008) Sae2, Exo1 and Sgs1 collaborate in DNA double-strand break processing. *Nature*, **455**, 770–774.
 32. Zhu, Z., Chung, W.H., Shim, E.Y., Lee, S.E. and Ira, G. (2008) Sgs1 helicase and two nucleases Dna2 and Exo1 resect DNA double-strand break ends. *Cell*, **134**, 981–994.
 33. Bakkenist, C.J. and Kastan, M.B. (2003) DNA damage activates ATM through intermolecular autophosphorylation and dimer dissociation. *Nature*, **421**, 499–506.
 34. Buis, J., Wu, Y., Deng, Y., Leddon, J., Westfield, G., Eckersdorff, M., Sekiguchi, J.M., Chang, S. and Ferguson, D.O. (2008) Mre11 nuclease activity has essential roles in DNA repair and genomic stability distinct from ATM activation. *Cell*, **135**, 85–96.
 35. Morin, I., Ngo, H.P., Greenall, A., Zubko, M.K., Morrice, N. and Lydall, D. (2008) Checkpoint-dependent phosphorylation of Exo1 modulates the DNA damage response. *EMBO J.*, **27**, 2400–2410.
 36. Matsuoka, S., Ballif, B.A., Smogorzewska, A., McDonald, E.R. III, Hurov, K.E., Luo, J., Bakalarski, C.E., Zhao, Z., Solimini, N., Lerenthal, Y. *et al.* (2007) ATM and ATR substrate analysis reveals extensive protein networks responsive to DNA damage. *Science*, **316**, 1160–1166.
 37. van der Horst, A. and Khanna, K.K. (2009) The peptidyl-prolyl isomerase Pin1 regulates cytokinesis through Cep55. *Cancer Res.*, **69**, 6651–6659.
 38. Nimmonkar, A.V., Ozsoy, A.Z., Genschel, J., Modrich, P. and Kowalczykowski, S.C. (2008) Human exonuclease 1 and BLM helicase interact to resect DNA and initiate DNA repair. *Proc. Natl Acad. Sci. USA*, **105**, 16906–16911.
 39. El-Shemerly, M., Hess, D., Pyakurel, A.K., Moselhy, S. and Ferrari, S. (2008) ATR-dependent pathways control hEXO1 stability in response to stalled forks. *Nucleic Acids Res.*, **36**, 511–519.
 40. Di Virgilio, M., Ying, C.Y. and Gautier, J. (2009) PIKK-dependent phosphorylation of Mre11 induces MRN complex inactivation by disassembly from chromatin. *DNA Repair*, **8**, 1311–1320.
 41. Beucher, A., Birraux, J., Tchouandong, L., Barton, O., Shibata, A., Conrad, S., Goodarzi, A.A., Kremler, A., Jeggo, P.A. and Lobrich, M. (2009) ATM and Artemis promote homologous recombination of radiation-induced DNA double-strand breaks in G2. *EMBO J.*, **28**, 3413–3427.