

Regulation of Mitochondrial Poly(ADP-Ribose) Polymerase Activation by the β -Adrenoceptor/cAMP/Protein Kinase A Axis during Oxidative Stress

Attila Brunyanszki, Gabor Olah, Ciro Coletta, Bartosz Szczesny, and Csaba Szabo

Department of Anesthesiology, University of Texas Medical Branch, Galveston, Texas

Received June 18, 2014; accepted July 28, 2014

ABSTRACT

We investigated the regulation of mitochondrial poly(ADP-ribose) polymerase 1 (PARP1) by the cyclic adenosine monophosphate (cAMP)/protein kinase A (PKA) system during oxidative stress in U937 monocytes. Oxidative stress induced an early (10 minutes) mitochondrial DNA damage, and concomitant activation of PARP1 in the mitochondria. These early events were followed by a progressive mitochondrial oxidant production and nuclear PARP1 activation (by 6 hours). These processes led to a functional impairment of mitochondria, culminating in cell death of mixed (necrotic/apoptotic) type. β -Adrenoceptor blockade with propranolol or inhibition of its downstream cAMP/PKA signaling attenuated, while β -adrenoceptor agonists and cAMP/PKA

activators enhanced, the oxidant-mediated PARP1 activation. In the presence of cAMP, recombinant PKA directly phosphorylated recombinant PARP1 on serines 465 (in the automodification domain) and 782 and 785 (both in the catalytic domain). Inhibition of the β -adrenergic receptor/cAMP/PKA axis protected against the oxidant-mediated cell injury. Propranolol also suppressed PARP1 activation in peripheral blood leukocytes during bacterial lipopolysaccharide (LPS)-induced systemic inflammation in mice. We conclude that the activation of mitochondrial PARP1 is an early, active participant in oxidant-induced cell death, which is under the control of β -adrenoceptor/cAMP/PKA axis through the regulation of PARP1 activity by PARP1 phosphorylation.

Introduction

Poly(ADP-ribose) polymerase 1 (PARP1), the major form of the PARP superfamily, is generally viewed as a constitutive nuclear enzyme with physiologic roles in the regulation of DNA repair, chromatin remodeling, and gene transcription (Tulin et al., 2003; Oei et al., 2005; Hottiger et al., 2011; DeVos et al., 2012). Its pathophysiologic overactivation, as a result of oxidative DNA injury, has been implicated in the pathogenesis of several diseases, including neuroinjury, inflammation, and ischemia/reperfusion, and various forms of critical illness (Tulin et al., 2003; Jagtap and Szabo, 2005; DeVos et al., 2012; Szanto et al., 2012; Curtin and Szabo, 2013; Burkle and Virag, 2013). In addition to the nuclear isoform of PARP1 several sets of studies have identified PARP1 in the mitochondrial compartment as well; it appears that, in addition to nuclear PARP1, mitochondrial PARP1 also plays roles in the regulation of various cellular functions—at least in some cell types (Masmoudi et al., 1988; Du et al., 2003; Rossi et al., 2009, reviewed in Burkle and Virag, 2013).

Protein kinase A (PKA), a cyclic adenosine monophosphate (cAMP)-dependent serine/threonine kinase, is one of the major effector proteins of cAMP and a regulator of cellular function in health and disease (Gancedo, 2013; Taylor et al., 2013; Valsecchi et al., 2013). The purpose of the current study was to determine whether PARP1 activity is regulated by the β -adrenoceptor/cAMP/PKA system during oxidative stress. The results demonstrate an early activation of the mitochondrial form of PARP1, unveil its regulation via phosphorylation by PKA through β -adrenoceptor/cAMP signaling and show the role of these processes in oxidant-induced cell death.

Materials and Methods

All chemicals were obtained from Sigma-Aldrich (St. Louis, MO) unless stated otherwise.

Cell Culture. U937 human monocyte histiocytic lymphoma cells, and C2C12 mouse myoblast cells were obtained from ATCC.

Western Blotting Analysis. Western blotting analysis was carried out as previously described (Gerö et al., 2013a) using anti-PARP1 antibody (Cell Signaling Technology, Beverly, MA), anti- β -receptor antibody (Abcam, Cambridge, MA), anti-protein kinase A (R&D Systems, Minneapolis, MN), anti- β -actin-horseradish peroxidase (HRP) conjugate (Santa Cruz Biotechnology, Inc., Dallas, TX), anti-rabbit-IgG (Cell Signaling Technology), and anti-mouse-IgG (Cell Signaling Technology). PARP1 activity was quantified by detection of its product, poly

This work was supported by the National Institutes of Health [Grant GM060338] (to C.S.); the Shriners Hospitals for Children [Grant 85800] (to C.S.); and the American Heart Association [Grant 12POST12060643] (to C.C.).
dx.doi.org/10.1124/mol.114.094318.

ABBREVIATIONS: cAMP, cyclic adenosine monophosphate; DDA, 2,5-dideoxy-adenosine; FSK, forskolin; H₂O₂, hydrogen peroxide; HRP, horseradish peroxidase; ISP, isoproterenol; LC, liquid chromatography; LDH, lactate dehydrogenase; LPS, bacterial lipopolysaccharide; NAD⁺, nicotinamide adenine dinucleotide; PAR, poly(ADP-ribose); PARG, poly(ADP-ribose) glycohydrolase; PARP1, poly(ADP-ribose) polymerase 1; PCR, polymerase chain reaction; PJ34, *N*-(6-oxo-5,6-dihydrophenanthridin-2-yl)-*N,N*-dimethylaminoacetamide hydrochloride; PKA, protein kinase A; PLA, proximity ligation assay; PP, propranolol; siRNA, small interfering RNA.

subsequent cell death response (Tann et al., 2011; Guicciardi et al., 2013; Virag et al., 2013; Dorn, 2013). In the current experimental model, early oxidant-mediated mitochondrial PARP1 activation, as well as other mitochondrial events, such as a progressive mitochondrial oxidant production (Fig. 2A) and mitochondrial electron transport defects (Fig. 11), probably contribute to the cell death. The results of the current research demonstrate that these processes are attenuated by blocking the early-onset PKA activation and the subsequent PARP1 activation. The working model outlined in Fig. 12 shows one possible interpretation of the findings of the current report: the cAMP/PKA axis is stimulated by β -adrenoceptor agonists (such as isoproterenol), cAMP analogs (such as 8-bromo-cAMP), and by adenylyl cyclase activators (such as forskolin), leading to mitochondrial PARP1 phosphorylation and increased PARP activity. Moreover, the cAMP/PKA axis is inhibited by β -adrenoceptor antagonists (such as propranolol), by adenylyl cyclase inhibitor (such as DDA) and by inhibition/downregulation of PKA (by the pharmacological inhibitor Rp-cAMP or by genetic silencing of PKA) leading to the inhibition of mitochondrial PARP1 phosphorylation and decreased PARP activity. In the Fig. 12, mitochondrial PKA is shown to be stimulated by cytosolic cAMP. It must be pointed out that multiple lines of data indicate that the cytosolic and mitochondrial cAMP pools do not communicate in most cases. At the same time, several sets of data also indicate that cell membrane and cytosolic signals can elevate intramitochondrial PKA activity, followed by phosphorylation of intramitochondrial proteins (reviewed in Lefkimmiatis and Zaccolo, 2014). Yet another possibility may be the extramitochondrial phosphorylation of PARP1, followed by its transport into the mitochondria, even though the early time course of extranuclear PARP1 phosphorylation (see Fig. 9) tends to speak against this possibility.

Although many additional details of the underlying mechanisms remain to be explored, the current findings may have implications for a number of cellular processes that are known to be regulated by PARP1, including cellular metabolism and DNA repair, given the fact that both of these processes can be regulated by the β -adrenoceptor/cAMP system (Carlucci et al., 2008; Cho and Juhn, 2012; Valsecchi et al., 2013; Lefkimmiatis and Zaccolo, 2014). However, these aspects remain to be directly investigated in further studies. The current findings may also provide a mechanistic explanation for our previously observed clinical/translational findings (Olah et al., 2011) showing that treatment of patients with severe burn injury with propranolol suppresses PARP1 activation in endothelial cells and tissue-resident mononuclear cells. Further studies are needed to determine whether modulation of PARP1 phosphorylation and its consequent catalytic activity by various drugs targeting the β -adrenoceptor/cAMP/PKA system may be used for therapeutic modulation of PARP1 in various pathophysiologic conditions associated with oxidative stress.

Acknowledgments

The authors thank Drs. Haag and Luo of the Mass Spectrometry Core Facility of UTMB for the mass spectrometry analysis.

Authorship Contributions

Participated in research design: Szabo, Brunyanszki and Szczesny.
Conducted experiments: Brunyanszki, Coletta, Oláh, Szczesny.

Performed data analysis: Brunyanszki, Szczesny.

Wrote or contributed to the writing of the manuscript: Szabo, Brunyanszki, Szczesny.

References

- Alamdary SZ, Digaleh H, and Khodagholfi F (2013) Dual contradictory effect of H-89 on neuronal retraction, death and inflammation in differentiated PC12 cells subjected to oxidative stress. *J Mol Neurosci* **51**:1030–1037.
- Bürkle A and Virág L (2013) Poly(ADP-ribose): PARadigms and PARadoxes. *Mol Aspects Med* **34**:1046–1065.
- Carlucci A, Lignitto L, and Feliciello A (2008) Control of mitochondria dynamics and oxidative metabolism by cAMP, AKAPs and the proteasome. *Trends Cell Biol* **18**: 604–613.
- Chen G, Pekary AE, Sugawara M, and Hershman JM (1993) Effect of exogenous hydrogen peroxide on iodide transport and iodine organification in FRTL-5 rat thyroid cells. *Acta Endocrinol (Copenh)* **129**:89–96.
- Cho EA and Juhn YS (2012) The cAMP signaling system inhibits the repair of γ -ray-induced DNA damage by promoting Epac1-mediated proteasomal degradation of XRCC1 protein in human lung cancer cells. *Biochem Biophys Res Commun* **422**:256–262.
- Curtin NJ and Szabo C (2013) Therapeutic applications of PARP inhibitors: anticancer therapy and beyond. *Mol Aspects Med* **34**:1217–1256.
- De Vos M, Schreiber V, and Dantzer F (2012) The diverse roles and clinical relevance of PARPs in DNA damage repair: current state of the art. *Biochem Pharmacol* **84**: 137–146.
- Dorn GW, 2nd (2013) Molecular mechanisms that differentiate apoptosis from programmed necrosis. *Toxicol Pathol* **41**:227–234.
- Du L, Zhang X, Han YY, Burke NA, Kochanek PM, Watkins SC, Graham SH, Carcillo JA, Szabó C, and Clark RS (2003) Intra-mitochondrial poly(ADP-ribose) contributes to NAD⁺ depletion and cell death induced by oxidative stress. *J Biol Chem* **278**:18426–18433.
- Gagné JP, Morel X, Gagné P, Labelle Y, Droit A, Chevalier-Paré M, Bourassa S, McDonald D, Hendzel MJ, and Prigent C et al. (2009) Proteomic investigation of phosphorylation sites in poly(ADP-ribose) polymerase-1 and poly(ADP-ribose) glycohydrolase. *J Proteome Res* **8**:1014–1029.
- Gancedo JM (2013) Biological roles of cAMP: variations on a theme in the different kingdoms of life. *Biol Rev Camb Philos Soc* **88**:645–668.
- Geraets L, Moonen HJ, Wouters EF, Bast A, and Hageman GJ (2006) Caffeine metabolites are inhibitors of the nuclear enzyme poly(ADP-ribose)polymerase-1 at physiological concentrations. *Biochem Pharmacol* **72**:902–910.
- Gerő D, Szoleczky P, Módis K, Pribis JP, Al-Abed Y, Yang H, Chevan S, Billiar TR, Tracey KJ, and Szabo C (2013a) Identification of pharmacological modulators of HMGB1-induced inflammatory response by cell-based screening. *PLoS ONE* **8**: e65994.
- Gerő D, Szoleczky P, Suzuki K, Módis K, Oláh G, oletta C, and Szabo C (2013b) Cell-based screening identifies paroxetine as an inhibitor of diabetic endothelial dysfunction. *Diabetes* **62**:953–964.
- Guicciardi ME, Malhi H, Mott JL, and Gores GJ (2013) Apoptosis and necrosis in the liver. *Compr Physiol* **3**:977–1010.
- Ha HC and Snyder SH (1999) Poly(ADP-ribose) polymerase is a mediator of necrotic cell death by ATP depletion. *Proc Natl Acad Sci USA* **96**:13978–13982.
- Hegedus C, Lakatos P, Oláh G, Tóth BI, Gergely S, Szabó E, Biró T, Szabó C, and Virág L (2008) Protein kinase C protects from DNA damage-induced necrotic cell death by inhibiting poly(ADP-ribose) polymerase-1. *FEBS Lett* **582**:1672–1678.
- Hottiger MO, Boothby M, Koch-Nolte F, Lüscher B, Martin NM, Plummer R, Wang ZQ, and Ziegler M (2011) Progress in the function and regulation of ADP-Ribosylation. *Sci Signal* **4**:mr5.
- Jagtap P, Soriano FG, Virág L, Liaudet L, Mabley J, Szabó E, Haskó G, Marton A, Lorigados CB, and Gallyas F, Jr et al. (2002) Novel phenanthridinone inhibitors of poly (adenosine 5'-diphosphate-ribose) synthetase: potent cytoprotective and antishock agents. *Crit Care Med* **30**:1071–1082.
- Jagtap P and Szabó C (2005) Poly(ADP-ribose) polymerase and the therapeutic effects of its inhibitors. *Nat Rev Drug Discov* **4**:421–440.
- Kauppinen TM, Chan WY, Suh SW, Wiggins AK, Huang EJ, and Swanson RA (2006) Direct phosphorylation and regulation of poly(ADP-ribose) polymerase-1 by extracellular signal-regulated kinases 1/2. *Proc Natl Acad Sci USA* **103**:7136–7141.
- Leadsham JE and Gourlay CW (2010) cAMP/PKA signaling balances respiratory activity with mitochondria dependent apoptosis via transcriptional regulation. *BMC Cell Biol* **11**:92.
- Lefkimmiatis K, Leronni D, and Hofer AM (2013) The inner and outer compartments of mitochondria are sites of distinct cAMP/PKA signaling dynamics. *J Cell Biol* **202**:453–462.
- Lefkimmiatis K and Zaccolo M (2014) cAMP signaling in subcellular compartments. *Pharmacol Ther* DOI: 10.1016/j.pharmthera.2014.03.008 [published ahead of print].
- Lopez-Hellin J, Garcia-Arumi E, and Schwartz S (1998) Oxidative stress induces age-dependent changes in lymphocyte protein synthesis and second messenger levels. *Life Sci* **63**:13–21.
- Mabley JG, Horváth EM, Murthy KG, Zsengeller Z, Vaslin A, Benko R, Kollai M, and Szabó C (2005) Gender differences in the endotoxin-induced inflammatory and vascular responses: potential role of poly(ADP-ribose) polymerase activation. *J Pharmacol Exp Ther* **315**:812–820.
- Mabley JG, Wallace R, Pacher P, Murphy K, and Szabó C (2007) Inhibition of poly (adenosine diphosphate-ribose) polymerase by the active form of vitamin D. *Int J Mol Med* **19**:947–952.
- Masmoudi A, Islam F, and Mandel P (1988) ADP-ribosylation of highly purified rat brain mitochondria. *J Neurochem* **51**:188–193.
- Módis K, Gerő D, Erdélyi K, Szoleczky P, DeWitt D, and Szabo C (2012) Cellular bioenergetics is regulated by PARP1 under resting conditions and during oxidative stress. *Biochem Pharmacol* **83**:633–643.
- Módis K, Gerő D, Stangl R, Rosero O, Sziárdó A, Lotz G, Mohácsik P, Szoleczky P, Coletta C, and Szabó C (2013a) Adenosine and inosine exert cytoprotective effects in an in vitro model of liver ischemia-reperfusion injury. *Int J Mol Med* **31**:437–446.

- Módos K, Coletta C, Erdélyi K, Papapetropoulos A, and Szabo C (2013b) Intra-mitochondrial hydrogen sulfide production by 3-mercaptopyruvate sulfurtransferase maintains mitochondrial electron flow and supports cellular bioenergetics. *FASEB J* **27**:601–611.
- Oei SL, Keil C, and Ziegler M (2005) Poly(ADP-ribosylation) and genomic stability. *Biochem Cell Biol* **83**:263–269.
- Oláh G, Finnerty CC, Sbrana E, Elijah I, Gerö D, Herndon DN, and Szabó C (2011) Increased poly(ADP-ribosylation) in skeletal muscle tissue of pediatric patients with severe burn injury: prevention by propranolol treatment. *Shock* **36**: 18–23.
- Rossi MN, Carbone M, Mostocotto C, Mancone C, Tripodi M, Maione R, and Amati P (2009) Mitochondrial localization of PARP-1 requires interaction with mitofilin and is involved in the maintenance of mitochondrial DNA integrity. *J Biol Chem* **284**: 31616–31624.
- Srinivasan S, Spear J, Chandran K, Joseph J, Kalyanaraman B, and Avadhani NG (2013) Oxidative stress induced mitochondrial protein kinase A mediates cytochrome c oxidase dysfunction. *PLoS ONE* **8**:e77129.
- Suzuki K, Olah G, Modis K, Coletta C, Kulp G, Gerö D, Szoleczky P, Chang T, Zhou Z, and Wu L et al. (2011) Hydrogen sulfide replacement therapy protects the vascular endothelium in hyperglycemia by preserving mitochondrial function. *Proc Natl Acad Sci USA* **108**:13829–13834.
- Szabó C, Zingarelli B, O'Connor M, and Salzman AL (1996) DNA strand breakage, activation of poly (ADP-ribose) synthetase, and cellular energy depletion are involved in the cytotoxicity of macrophages and smooth muscle cells exposed to peroxynitrite. *Proc Natl Acad Sci USA* **93**:1753–1758.
- Szabo C, Pacher P, and Swanson RA (2006) Novel modulators of poly(ADP-ribose) polymerase. *Trends Pharmacol Sci* **27**:626–630.
- Szántó M, Brunyánszki A, Kiss B, Nagy L, Gergely P, Virág L, and Bai P (2012) Poly (ADP-ribose) polymerase-2: emerging transcriptional roles of a DNA-repair protein. *Cell Mol Life Sci* **69**:4079–4092.
- Szczesny B, Olah G, Walker DK, Volpi E, Rasmussen BB, Szabo C, and Mitra S (2013) Deficiency in repair of the mitochondrial genome sensitizes proliferating myoblasts to oxidative damage. *PLoS ONE* **8**:e75201.
- Tanaka Y, Koide SS, Yoshihara K, and Kamiya T (1987) Poly (ADP-ribose) synthetase is phosphorylated by protein kinase C in vitro. *Biochem Biophys Res Commun* **148**:709–717.
- Tann AW, Boldogh I, Meiss G, Qian W, Van Houten B, Mitra S, and Szczesny B (2011) Apoptosis induced by persistent single-strand breaks in mitochondrial genome: critical role of EXOG (5'-exo/endonuclease) in their repair. *J Biol Chem* **286**: 31975–31983.
- Taylor SS, Zhang P, Steichen JM, Keshwani MM, and Kornev AP (2013) PKA: lessons learned after twenty years. *Biochim Biophys Acta* **1834**:1271–1278.
- Tulin A, Chinenov Y, and Spradling A (2003) Regulation of chromatin structure and gene activity by poly(ADP-ribose) polymerases. *Curr Top Dev Biol* **56**:55–83.
- Valsecchi F, Ramos-Espiritu LS, Buck J, Levin LR, and Manfredi G (2013) cAMP and mitochondria. *Physiology (Bethesda)* **28**:199–209.
- Virág L, Salzman AL, and Szabó C (1998) Poly(ADP-ribose) synthetase activation mediates mitochondrial injury during oxidant-induced cell death. *J Immunol* **161**: 3753–3759.
- Virág L and Szabó C (2001) Purines inhibit poly(ADP-ribose) polymerase activation and modulate oxidant-induced cell death. *FASEB J* **15**:99–107.
- Virág L, Robaszkiewicz A, Rodríguez-Vargas JM, and Oliver FJ (2013) Poly(ADP-ribose) signaling in cell death. *Mol Aspects Med* **34**:1153–1167.
- Wang Y, Dawson VL, and Dawson TM (2009) Poly(ADP-ribose) signals to mitochondrial AIF: a key event in parthanatos. *Exp Neurol* **218**:193–202.
- Yu SW, Andrabi SA, Wang H, Kim NS, Poirier GG, Dawson TM, and Dawson VL (2006) Apoptosis-inducing factor mediates poly(ADP-ribose) (PAR) polymer-induced cell death. *Proc Natl Acad Sci USA* **103**:18314–18319.
- Zhang J, Dawson VL, Dawson TM, and Snyder SH (1994) Nitric oxide activation of poly(ADP-ribose) synthetase in neurotoxicity. *Science* **263**:687–689.

Address correspondence to: Dr. Csaba Szabo, Department of Anesthesiology, The University of Texas Medical Branch at Galveston, 601 Harborside Drive, Galveston, TX 77555. E-mail: szabocsaba@aol.com
