

1 Problem statement

1.1 Strong formulation

We will work on a two-dimensional Helmholtz problem. Let $\Omega \subset \mathbb{R}^2$ be an open bounded region whose border $\partial\Omega$ is sufficiently smooth. Our strong formulation is the following:

$$\text{Find } u \text{ such that: } \begin{cases} \Delta u + k^2 u = -f & \text{in } \Omega \\ u = 0 & \text{on } \partial\Omega \end{cases}$$

$\Delta = \nabla^2$ is the Laplace operator, u could be the unknown acoustic pressure field, k is the wavenumber and f is a source term. In general, u and k can be complex-valued. However, the paramter studies will be conducted on real-valued cases, with $k > 0$. The extension to complex-valued cases should be straightforward.

1.2 Galerkin formulation

The usual Galerkin formulation for our problem is the following:

$$\text{Find } u^h \in \mathcal{V}^h \text{ such that } \forall v^h \in \mathcal{V}^h, a_G(u^h, v^h) = \langle f, v^h \rangle_{L^2(\Omega)}$$

Where $\mathcal{V}^h \subset H_0^1(\Omega)$ is finite-dimensional, $\langle u, v \rangle_{L^2(\Omega)}$ is the $L^2(\Omega)$ scalar product, and a_G is the following sesquilinear form:

$$\begin{aligned} a_G : H_0^1(\Omega)^2 &\rightarrow \mathbb{C} \\ (u, v) &\mapsto \langle \nabla u, \nabla v \rangle_{L^2(\Omega)} - k^2 \langle u, v \rangle_{L^2(\Omega)} \end{aligned}$$

We will use a mesh method to build \mathcal{V}^h , partitioning Ω in mutually exclusive elements.

1.3 GLS

The Galerkin/least-squares formulation uses the previous formulation and adds other terms whose purpose is to minimize the square of the residue over the element interiors $\tilde{\Omega}$. Their contribution is weighted by a parameter τ :

$$\text{Find } u^h \in \mathcal{V}^h, \forall v^h \in \mathcal{V}^h, a_{GLS}(u^h, v^h) = \langle f, v^h \rangle_{L^2(\Omega)} + \langle \tau f, \mathcal{L}v^h \rangle_{L^2(\tilde{\Omega})}$$

Where $\mathcal{L} = \Delta + k^2$ is the Helmholtz operator, and a_{GLS} is the following sesquilinear form:

$$\begin{aligned} a_{GLS} : H_0^1(\Omega)^2 &\rightarrow \mathbb{C} \\ (u, v) &\mapsto a_G(u, v) + \langle \tau \mathcal{L}u, \mathcal{L}v \rangle_{L^2(\tilde{\Omega})} \end{aligned}$$

2 Dispersion analysis and optimal parameter

The Helmholtz equation suffers from the pollution effect, causing numerical wavenumbers to lose accuracy with the physical wavenumber increasing.

Exact solutions When considering the sourceless ($f = 0$) and free-space version of our problem, the exact solutions are $\forall x \in \mathbb{R}$, $u(x) = Ce^{ikx}$. Our studies will revolve around this case.

$$\sum_{i,j \in -1,0,1} A_{G,0,0}^{i,j} U_{i,j}^h = 0 \quad (1)$$

Setting $2h_\xi = h_\xi^- + h_\xi^+$ with ξ being x or y .

The values of the nonzero entries are the following:

Four corner nodes ($i, j \in \{-, +\}$):

$$A_{G,0,0}^{i,j} = \frac{k^2 h_x^i h_y^j}{36} + \frac{h_x^i}{6h_y^j} + \frac{h_y^j}{6h_x^i}$$

Four side nodes:

$$\begin{aligned} A_{G,0,0}^{\pm,0} &= \frac{k^2 h_x^\pm h_y}{9} + \frac{2h_y}{3h_x^\pm} - \frac{h_x^\pm}{3} \frac{h_y}{h_y^- h_y^+} \\ A_{G,0,0}^{0,\pm} &= \frac{k^2 h_x h_y^\pm}{9} + \frac{2h_x}{3h_y^\pm} - \frac{h_y^\pm}{3} \frac{h_x}{h_x^- h_x^+} \end{aligned}$$

Center node:

$$A_{G,0,0}^{0,0} = \frac{4k^2 h_x h_y}{9} - \frac{4h_x}{3} \frac{h_y}{h_y^- h_y^+} - \frac{4h_y}{3} \frac{h_x}{h_x^- h_x^+}$$

Setting $\alpha_\xi^\pm h_\xi = h_\xi^\pm$ with ξ being x or y .

Four side nodes:

$$\begin{aligned} A_{G,0,0}^{\pm,0} &= \frac{k^2 h_x^\pm h_y}{9} + \frac{2h_y}{3h_x^\pm} - \frac{h_x^\pm}{3\alpha_y^- \alpha_y^+ h_y} \\ A_{G,0,0}^{0,\pm} &= \frac{k^2 h_x h_y^\pm}{9} + \frac{2h_x}{3h_y^\pm} - \frac{h_y^\pm}{3\alpha_x^- \alpha_x^+ h_x} \end{aligned}$$

Center node:

$$A_{G,0,0}^{0,0} = \frac{4k^2 h_x h_y}{9} - \frac{4h_x}{3\alpha_y^- \alpha_y^+ h_y} - \frac{4h_y}{3\alpha_x^- \alpha_x^+ h_x}$$

$$k^2 = \frac{6 \left(2f_x^+ h_x^- \left((h_x^+)^2 h_y - (h_y^+)^2 h_y^- - h_y^+ (h_y^-)^2 \right) + 2f_x^- h_x^+ \left((h_x^-)^2 h_y - (h_y^-)^2 h_y^- - h_y^- (h_y^+)^2 \right) + 2f_y^+ h_y^- \left(h_x \right) \right)}{}$$

$$\gamma_0 + \sum_{\xi \in \{x,y\}} \sum_{i \in \{-,+\}} \gamma_\xi^i f_\xi^i + \quad (2)$$

Terme indépendant en haut

$$\begin{aligned}
& + 12h_x^- h_y^+ \left((h_x^+)^2 + (h_y^-)^2 \right) \\
& \quad + 12h_x^- h_y^- \left((h_x^+)^2 + (h_y^+)^2 \right) \\
& \quad + 12h_x^+ h_y^+ \left((h_x^-)^2 + (h_y^-)^2 \right) \\
& \quad + 12h_x^+ h_y^- \left((h_x^-)^2 + (h_y^+)^2 \right)
\end{aligned}$$

Termes en f en haut

$$\begin{aligned}
& 12f_x^+ h_x^- \left((h_x^+)^2 h_y - (h_y^+)^2 h_y^- - h_y^+ (h_y^-)^2 \right) + 12f_x^- h_x^+ \left((h_x^-)^2 h_y - (h_y^+)^2 h_y^- - h_y^+ (h_y^-)^2 \right) \\
& + 12f_y^+ h_y^- \left(h_x (h_y^+)^2 - (h_x^+)^2 h_x^- - h_x^+ (h_x^-)^2 \right) + 12f_y^- h_y^+ \left(h_x (h_y^-)^2 - h_x^+ (h_x^-)^2 - (h_x^+)^2 h_x^- \right)
\end{aligned}$$

Terme indépendant en bas (en facteur des 4h)

$$16h_x h_y$$

Termes en f en bas (en facteur des 4h)

$$4f_x^+ h_x^+ h_y \quad + \quad 4f_x^- h_x^- h_y \quad + \quad 4f_y^+ h_x h_y^+ \quad + \quad 4f_y^- h_x h_y^-$$

Termes en g en bas (en facteur des 4h)

$$+ h_x^+ h_y^+ g^{++} + h_x^+ h_y^- g^{+-} + h_x^- h_y^+ g^{-+} + h_x^- h_y^- g^{--}$$