

# **Tutorial: Similarity and Numerical Solutions for Third Order, Partial Differential Equations**

Ray Fernandes

Department of Mathematical Sciences, University of Bath

8th July, 2008



# Similarity Solutions for the Third Order Linear PDE

## Third Order linear PDE

$$u_t = u_{xxx}, \quad \text{in } \mathbb{R} \times \mathbb{R}_+$$

# Similarity Solutions for the Third Order Linear PDE

## Third Order linear PDE

$$u_t = u_{xxx}, \quad \text{in } \mathbb{R} \times \mathbb{R}_+$$

## Similarity Solutions

$$u(x, t) = t^{-a} F(y), \quad y = xt^{-b}.$$

## Reduces to an ODE

$$-at^{-a-1}F - bt^{-a-1}F'y = t^{-a-3b}F'''.$$

## Reduces to an ODE

$$-at^{-a-1}F - bt^{-a-1}F'y = t^{-a-3b}F'''.$$

## Equating powers of $t$

$$a + 1 = a + 3b.$$

### Reduces to an ODE

$$-at^{-a-1}F - bt^{-a-1}F'y = t^{-a-3b}F'''.$$

### Equating powers of $t$

$$b = \frac{1}{3}.$$

### Reduces to an ODE

$$-at^{-a-1}F - bt^{-a-1}F'y = t^{-a-3b}F'''.$$

### Equating powers of $t$

$$b = \frac{1}{3}.$$

### Equating coefficients

$$a = \frac{1}{3}.$$

**$F$  solves the ODE**

$$F''' = -\frac{1}{3} F' y - \frac{1}{3} F.$$



**$F$  solves the ODE**

$$F''' = -\frac{1}{3} (Fy)'.$$

**$F$  solves the ODE**

$$F''' = -\frac{1}{3} (Fy)'.$$

**Can integrate once**

$$F'' = -\frac{1}{3} Fy.$$

**$F$  solves the ODE**

$$F''' = -\frac{1}{3} (Fy)'.$$

**Can integrate once**

$$F'' = -\frac{1}{3} Fy.$$

Yields the Airy equation.

# Asymptotics

Asymptotic analysis can give good approximations, when exact solutions may not be found. Describes limiting behaviour.

# Asymptotics

Asymptotic analysis can give good approximations, when exact solutions may not be found. Describes limiting behaviour.

## Asymptotic To

$$f(x) \sim g(x) \quad \text{as } x \rightarrow x_0 \quad \Longleftrightarrow \quad \lim_{x \rightarrow x_0} \frac{f(x)}{g(x)} = 1.$$

# Asymptotics

Asymptotic analysis can give good approximations, when exact solutions may not be found. Describes limiting behaviour.

## Asymptotic To

$$f(x) \sim g(x) \quad \text{as } x \rightarrow x_0 \quad \Longleftrightarrow \quad f(x) - g(x) = o(f(x)).$$

# Asymptotics

Asymptotic analysis can give good approximations, when exact solutions may not be found. Describes limiting behaviour.

## Asymptotic To

$$f(x) \sim g(x) \quad \text{as } x \rightarrow x_0 \quad \Longleftrightarrow \quad f(x) - g(x) = o(f(x)).$$

## Order Less Than

$$f(x) = o(g(x)) \quad \text{as } x \rightarrow x_0 \quad \Longleftrightarrow \quad \lim_{x \rightarrow x_0} \frac{f(x)}{g(x)} = 0.$$

# Asymptotic Analysis of the Linear Equation

Assume solution,  $F(y)$ , is exponential as  $y \rightarrow +\infty$

$$F(y) = e^{s(y)}.$$



# Asymptotic Analysis of the Linear Equation

**Assume solution,  $F(y)$ , is exponential as  $y \rightarrow +\infty$**

$$F(y) = e^{s(y)}.$$

**Derivatives of  $F$**

$$F' = s' e^s \quad \text{and} \quad F'' = [s'' + (s')^2] e^s.$$

# Asymptotic Analysis of the Linear Equation

Assume solution,  $F(y)$ , is exponential as  $y \rightarrow +\infty$

$$F(y) = e^{s(y)}.$$

Derivatives of  $F$

$$F' = s' e^s \quad \text{and} \quad F'' = [s'' + (s')^2] e^s.$$

Assume that  $s(y)$  is some polynomial, such that  $s(y) \sim \alpha y^\beta$ .  
Then  $s' \sim \alpha \beta y^{\beta-1}$  and  $s'' \sim \alpha \beta (\beta - 1) y^{\beta-2}$ .

# Asymptotic Analysis of the Linear Equation

Assume solution,  $F(y)$ , is exponential as  $y \rightarrow +\infty$

$$F(y) = e^{s(y)}.$$

Derivatives of  $F$

$$F' = s' e^s \quad \text{and} \quad F'' = [s'' + (s')^2] e^s.$$

Assume that  $s(y)$  is some polynomial, such that  $s(y) \sim \alpha y^\beta$ .  
Then  $s' \sim \alpha \beta y^{\beta-1}$  and  $s'' \sim \alpha \beta (\beta - 1) y^{\beta-2}$ .

Second order linear ODE

$$F'' = -\frac{1}{3} F y.$$

# Asymptotic Analysis of the Linear Equation

Assume solution,  $F(y)$ , is exponential as  $y \rightarrow +\infty$

$$F(y) = e^{s(y)}.$$

Derivatives of  $F$

$$F' = s' e^s \quad \text{and} \quad F'' = [s'' + (s')^2] e^s.$$

Assume that  $s(y)$  is some polynomial, such that  $s(y) \sim \alpha y^\beta$ .  
Then  $s' \sim \alpha \beta y^{\beta-1}$  and  $s'' \sim \alpha \beta (\beta - 1) y^{\beta-2}$ .

Substituting into the ODE

$$\alpha \beta (\beta - 1) y^{\beta-2} + \alpha^2 \beta^2 y^{2\beta-2} \sim -\frac{1}{3} y$$

Two cases:  $\beta \leq 0$  and  $\beta > 0$ .

Two cases:  $\beta \leq 0$  and  $\beta > 0$ .

$$\alpha\beta(\beta - 1)y^{\beta-2} + \alpha^2\beta^2y^{2\beta-2} \sim -\frac{1}{3}y$$

**For  $\beta \leq 0$**

$$y^{\beta-2} = o(y) \quad \text{and} \quad y^{2\beta-2} = o(y)$$

Two cases:  $\beta \leq 0$  and  $\beta > 0$ .

$$\alpha\beta(\beta - 1)y^{\beta-2} + \alpha^2\beta^2y^{2\beta-2} \sim -\frac{1}{3}y$$

**For  $\beta \leq 0$**

$$y^{\beta-2} = o(y) \quad \text{and} \quad y^{2\beta-2} = o(y)$$

small term

Two cases:  $\beta \leq 0$  and  $\beta > 0$ .

$$\alpha\beta(\beta - 1)y^{\beta-2} + \alpha^2\beta^2y^{2\beta-2} \sim -\frac{1}{3}y$$

**For  $\beta \leq 0$**

$$y^{\beta-2} = o(y) \quad \text{and} \quad y^{2\beta-2} = o(y)$$

small term + small term



Two cases:  $\beta \leq 0$  and  $\beta > 0$ .

$$\alpha\beta(\beta - 1)y^{\beta-2} + \alpha^2\beta^2y^{2\beta-2} \sim -\frac{1}{3}y$$

**For  $\beta \leq 0$**

$$y^{\beta-2} = o(y) \quad \text{and} \quad y^{2\beta-2} = o(y) \quad \Rightarrow \text{no balance.}$$

small term + small term  $\approx$  larger term

$$\alpha\beta(\beta - 1)y^{\beta-2} + \alpha^2\beta^2y^{2\beta-2} \sim -\frac{1}{3}y$$

**For  $\beta > 0$**

$$y^{\beta-2} = o(y^{2\beta-2}) \quad \text{as } y \rightarrow \infty$$

$$\alpha\beta(\beta - 1)y^{\beta-2} + \alpha^2\beta^2y^{2\beta-2} \sim -\frac{1}{3}y$$

**For  $\beta > 0$**

$$y^{\beta-2} = o(y^{2\beta-2}) \quad \text{as } y \rightarrow \infty$$

**small term**

$$\alpha\beta(\beta-1)y^{\beta-2} + \alpha^2\beta^2y^{2\beta-2} \sim -\frac{1}{3}y$$

For  $\beta > 0$

$$y^{\beta-2} = o(y^{2\beta-2}) \quad \text{as } y \rightarrow \infty$$

small term

larger term

$$\alpha\beta(\beta - 1)y^{\beta-2} + \alpha^2\beta^2y^{2\beta-2} \sim -\frac{1}{3}y$$

**For  $\beta > 0$**

$$y^{\beta-2} = o(y^{2\beta-2}) \quad \text{as } y \rightarrow \infty \quad \Rightarrow \text{Terms can balance}$$

small term + 2nd larger term  $\sim$  larger term

## Balancing Leading Order Terms

$$\alpha^2 \beta^2 y^{2\beta-2} \sim -\frac{1}{3}y$$

## Balancing Leading Order Terms

$$\alpha^2 \beta^2 y^{2\beta-2} \sim -\frac{1}{3}y$$

## Equating Powers of $y$ we have

$$\beta = \frac{3}{2} \quad \text{and} \quad \alpha = \frac{2i}{3\sqrt{3}}.$$

## Balancing Leading Order Terms

$$\alpha^2 \beta^2 y^{2\beta-2} \sim -\frac{1}{3}y$$

## Equating Powers of $y$ we have

$$\beta = \frac{3}{2} \quad \text{and} \quad \alpha = \frac{2i}{3\sqrt{3}}.$$

$$s(y) \sim -\frac{2i}{3\sqrt{3}}y^{3/2}$$



## Balancing Leading Order Terms

$$\alpha^2 \beta^2 y^{2\beta-2} \sim -\frac{1}{3}y$$

## Equating Powers of $y$ we have

$$\beta = \frac{3}{2} \quad \text{and} \quad \alpha = \frac{2i}{3\sqrt{3}}.$$

$$s(y) = -\frac{2i}{3\sqrt{3}}y^{3/2} + c(y), \quad c(y) = o(y^{3/2})$$

Insert  $s(y)$  into the ODE again. Higher order terms cancel due to previous balance.

Insert  $s(y)$  into the ODE again. Higher order terms cancel due to previous balance.

**Second term in the expansion.**

$$c(y) \sim -\frac{1}{4} \ln y$$

Insert  $s(y)$  into the ODE again. Higher order terms cancel due to previous balance.

### Second term in the expansion.

$$c(y) \sim -\frac{1}{4} \ln y$$

$$s(y) \sim -\frac{2i}{3\sqrt{3}} y^{3/2} - \frac{1}{4} \ln y$$

Insert  $s(y)$  into the ODE again. Higher order terms cancel due to previous balance.

### Second term in the expansion.

$$c(y) \sim -\frac{1}{4} \ln y$$

$$s(y) \sim -\frac{2i}{3\sqrt{3}} y^{3/2} - \frac{1}{4} \ln y$$

Can keep expanding with as many terms as you like for greater accuracy.

### Asymptotic Behaviour as $y \rightarrow +\infty$

$$F(y) \sim y^{-1/4} \exp\left(-\frac{2i}{3\sqrt{3}} y^{3/2}\right) \quad \text{as } y \rightarrow +\infty.$$

### Asymptotic Behaviour as $y \rightarrow +\infty$

$$F(y) \sim y^{-1/4} \exp\left(-\frac{2i}{3\sqrt{3}} y^{3/2}\right) \quad \text{as } y \rightarrow +\infty.$$

To find the behaviour as  $y \rightarrow -\infty$ , let  $y \mapsto -y$  and we can do the same analysis as before.

### Asymptotic Behaviour as $y \rightarrow +\infty$

$$F(y) \sim y^{-1/4} \exp\left(-\frac{2i}{3\sqrt{3}} y^{3/2}\right) \quad \text{as } y \rightarrow +\infty.$$

To find the behaviour as  $y \rightarrow -\infty$ , let  $y \mapsto -y$  and we can do the same analysis as before.

### Asymptotic Behaviour as $y \rightarrow -\infty$

$$F(y) \sim y^{-1/4} \exp\left(\frac{2i}{3\sqrt{3}} y^{3/2}\right) \quad \text{as } y \rightarrow -\infty.$$



# Numerical Solutions using the Boundary Value Problem Solver

For all solutions  $F$ , of the ODE,  $cF$  is also a solution for any  $c \in \mathbb{R}$ .

# Numerical Solutions using the Boundary Value Problem Solver

For all solutions  $F$ , of the ODE,  $cF$  is also a solution for any  $c \in \mathbb{R}$ .

$F = 0$  is a solution and may be found when attempting to solve numerically.

# Numerical Solutions using the Boundary Value Problem Solver

For all solutions  $F$ , of the ODE,  $cF$  is also a solution for any  $c \in \mathbb{R}$ .

$F = 0$  is a solution and may be found when attempting to solve numerically.

Need to ensure non-zero solution, so set constraint  $\max |F| = 1$ .

# Numerical Solutions using the Boundary Value Problem Solver

For all solutions  $F$ , of the ODE,  $cF$  is also a solution for any  $c \in \mathbb{R}$ .

$F = 0$  is a solution and may be found when attempting to solve numerically.

Need to ensure non-zero solution, so set constraint  $\max |F| = 1$ .

Maximum attained at some point  $\hat{a}$ .

Can solve for solutions left and right of this point. Match the two to find the full solution.

Can solve for solutions left and right of this point. Match the two to find the full solution.


Need to move the point  $\hat{a}$ , until correct profile is found.

Can solve for solutions left and right of this point. Match the two to find the full solution.

Need to move the point  $\hat{a}$ , until correct profile is found.


Profile is correct if the second derivative, for both left and right solutions, are the same at the point  $\hat{a}$ . This ensures continuity at the point  $\hat{a}$ .

## Rescaled Fundamental Solution to $u_t = u_{xxx}$







## Rescaled Fundamental Solution to $u_t = u_{xxx}$



## Rescaled Fundamental Solution to $u_t = u_{xxx}$



## Rescaled Fundamental Solution to $u_t = u_{xxx}$



# Third Order Semilinear Model

## Semilinear PDE

$$u_t = u_{xxx} - |u|^{p-1}u \quad \text{in } \mathbb{R} \times \mathbb{R}_+.$$

# Third Order Semilinear Model

## Semilinear PDE

$$u_t = u_{xxx} - |u|^{p-1}u \quad \text{in} \quad \mathbb{R} \times \mathbb{R}_+.$$

## Very Singular Solutions (VSS)

$$u_*(x, t) = t^{-1/(p-1)}f(y), \quad y = xt^{-1/3}.$$

# Third Order Semilinear Model

## Semilinear PDE

$$u_t = u_{xxx} - |u|^{p-1}u \quad \text{in } \mathbb{R} \times \mathbb{R}_+.$$

## Very Singular Solutions (VSS)

$$u_*(x, t) = t^{-1/(p-1)}f(y), \quad y = xt^{-1/3}.$$

## $f$ solves the ODE

$$f''' + \frac{1}{3}f'y + \frac{1}{p-1}f - |f|^{p-1}f = 0 \quad \text{in } \mathbb{R}.$$

Asymptotic methods like that used for the linear case, cannot be applied here.

Asymptotic methods like that used for the linear case, cannot be applied here.

Nonlinear term  $|f|^{p-1}f$ , means that  $e^{s(y)}$  terms will not cancel.



Asymptotic methods like that used for the linear case, cannot be applied here.

Nonlinear term  $|f|^{p-1}f$ , means that  $e^{s(y)}$  terms will not cancel.

Can use knowledge of linear results, since semilinear case is the same, with a perturbation of  $|f|^{p-1}f$ .

# Numerical Solutions of the Semilinear problem

Scaling group of solutions no longer present, due to the term  $|f|^{p-1}f$

# Numerical Solutions of the Semilinear problem

Scaling group of solutions no longer present, due to the term  $|f|^{p-1}f \Rightarrow$  not necessary to use a maximising constraint.

# Numerical Solutions of the Semilinear problem

Scaling group of solutions no longer present, due to the term  $|f|^{p-1}f \Rightarrow$  not necessary to use a maximising constraint.

Cannot integrate to reduce the order, so must use 3 boundary conditions.

# Numerical Solutions of the Semilinear problem

Scaling group of solutions no longer present, due to the term  $|f|^{p-1}f \Rightarrow$  not necessary to use a maximising constraint.

Cannot integrate to reduce the order, so must use 3 boundary conditions.

Use 2 boundary conditions at the initial point:  $f = 0$  and  $f' = 0$ .  
Use 1 boundary condition at the end point:  $f = 0$ .

Problems arise, since we do not know exactly where the oscillations cross  $f = 0$ .

Problems arise, since we do not know exactly where the oscillations cross  $f = 0$ .

If wrong end point is found, then the oscillations are forced through that point and symmetry may be compromised.


Problems arise, since we do not know exactly where the oscillations cross  $f = 0$ .

If wrong end point is found, then the oscillations are forced through that point and symmetry may be compromised.


Move end point until best profile is found.




**Wrong profile to  $u_t = u_{xxx} - u^p$  with  $p = 2.9$**




**Wrong profile to  $u_t = u_{xxx} - u^p$  with  $p = 2.9$**




**Wrong profile to  $u_t = u_{xxx} - u^p$  with  $p = 2.9$**




Wrong profile to  $u_t = u_{xxx} - u^p$  with  $p = 2.9$




Wrong profile to  $u_t = u_{xxx} - u^p$  with  $p = 2.9$




**Wrong profile to  $u_t = u_{xxx} - u^p$  with  $p = 2.9$**



## Fundamental solution to $u_t = u_{xxx} - u^p$ with $p = 2.9$




## Fundamental solution to $u_t = u_{xxx} - u^p$ with $p = 2.9$






## Fundamental solution to $u_t = u_{xxx} - u^p$ with $p = 2$



## Fundamental solution to $u_t = u_{xxx} - u^p$ with $p = 2$



Conditions we look at to ensure the best possible profiles:

- Symmetry of tail for  $y \gg 1$ .
- Symmetry of tail occurs as close to 0 as possible.
- Minimisation of tail.

Conditions we look at to ensure the best possible profiles:

- Symmetry of tail for  $y \gg 1$ .
- Symmetry of tail occurs as close to 0 as possible.
- Minimisation of tail.

Conditions we look at to ensure the best possible profiles:

- Symmetry of tail for  $y \gg 1$ .
- Symmetry of tail occurs as close to 0 as possible.
- Minimisation of tail.

Conditions we look at to ensure the best possible profiles:

- Symmetry of tail for  $y \gg 1$ .
- Symmetry of tail occurs as close to 0 as possible.
- Minimisation of tail.