

590.8
F 63

Linnean Society of New South Wales.

THE
MACLEAY MEMORIAL VOLUME

EDITED BY
J. J. FLETCHER, M.A., B.Sc.

PUBLISHED BY THE SOCIETY
AND SOLD BY
ANGUS & ROBERTSON, AND TURNER & HENDERSON, SYDNEY,
DULAU & CO., 37 SOHO SQUARE, LONDON, AND FRIEDLÄNDER & SOHN, BERLIN.

September, 1893.

Price Three Guineas.



THE UNIVERSITY OF SYDNEY

MACLEAY MUSEUM

085164

SYDNEY:

F. CUNNINGHAME AND CO., PRINTERS,

PITT STREET.

1881

J. B. HARRISON, F.R.S.

PRINTED BY THE UNIVERSITY

THE UNIVERSITY OF SYDNEY
THE MACLEAY MUSEUM
THE UNIVERSITY OF SYDNEY



CONTENTS.

	PAGE
THE HON. SIR WILLIAM MACLEAY, KT., F.L.S., M.L.C.	VII.-LI.
 <i>Contributions to our Knowledge of Ceratodus. Part I.—The Blood Vessels.</i>	
By W. BALDWIN SPENCER, M.A., Professor of Biology in the University of Melbourne. (Plates I.-v.)	1
<i>The Pliocene Mollusca of New Zealand.</i> By F. W. HUTTON, F.R.S., Professor of Geology in Canterbury College, Christchurch. (Plates VI.-IX.)	35
<i>A Monograph of the Temnocephaleæ.</i> By WILLIAM A. HASWELL, M.A., D.Sc., F.L.S., Challis Professor of Biology, University of Sydney. (Plates x.-xv.)	93
<i>On an apparently New Type of the Platyhelminthes (Trematoda?).</i> By WILLIAM A. HASWELL, M.A., D.Sc., Challis Professor of Biology, University of Sydney. (Plate xvi.)	153
<i>Observations on the Myology of Palinurus Edwardsii, Hutton.</i> By Professor T. JEFFERY PARKER, D.Sc., F.R.S., and JOSEPHINE GORDON RICH. (Plates xvii.-xxi.)	159
<i>Observations upon the Anatomy of the Muzzle of the Ornithorhynchus.</i> By J. T. WILSON, M.B., Professor of Anatomy, University of Sydney, and C. J. MARTIN, M.B., B.Sc., Demonstrator of Physiology, University of Sydney. (Plates xxii.-xxiii.)	179
<i>On the Peculiar Rod-like Tactile Organs in the Integument and Mucous Membrane of the Muzzle of Ornithorhynchus.</i> By J. T. WILSON, M.B., Professor of Anatomy, University of Sydney, and C. J. MARTIN, M.B., B.Sc., Demonstrator of Physiology, University of Sydney. (Plates xxiv.-xxvi.)	190
<i>On Parmacochlea Fischeri, Smith.</i> By C. HEDLEY, F.L.S. (Plate xxvii.)	201
<i>On the Geographic Relations of the Floras of Norfolk and Lord Howe Islands.</i> By Professor RALPH TATE, F.L.S., F.G.S., &c. ...	205
* <i>Notes on an Undescribed Acacia from New South Wales.</i> By Baron von MUELLER, K.C.M.G., M. & Ph.D., LL.D., F.R.S. (Plate xxviii.)	222
* <i>Description of a New Hakea from Eastern New South Wales.</i> By Baron von MUELLER, K.C.M.G., M. & Ph.D., LL.D., F.R.S., and J. H. MAIDEN, F.L.S. (Plate xxix.)	226
<i>A Description of some of the Weapons and Implements of the Alligator Tribe, Port Essington, North Australia.</i> By R. ETHERIDGE, JUNR., Palæontologist to the Australian Museum, and Geological Survey of New South Wales. (Plates xxx.-xxxv.)	228
<i>Nematodes, mostly Australian and Fijian.</i> By N. A. COBB. (Plates xxxvi.-xlii.)	252

* The two plates illustrating these papers should have been numbered xxviii. and xxix. instead of xxix. and xxx. They were printed off, however, before it was known that a plate to which the number xxviii. had been assigned, was to be withdrawn.

28907



CORRIGENDA.

Appendix to Paper No. IV. (On an apparently new Type of the *Platyhelminthes*, by Prof. HASWELL), p. 153.

"Since this paper passed through the press my attention has been directed by Prof. R. Blanchard, of Paris, to the fact that a genus *Actinodactylus* was established in 1890 by Duchassaing for a Tunicate. I, therefore, propose to alter the termination, making it *Actinodactylella*. I have omitted to give a specific name in the paper, and now propose to call this remarkable worm *Actinodactylella Blanchardi*."—W. A. H.

Page 88, line 23—a full point has been omitted after Dieff.

Page 128, line 21—*for* decribed *read* described.

Page 220, line 24—for *Passiflora Hebertiana* read *Passiflora Herbertiana*.

For Plate XXIX. *read* Plate XXVIII.

For Plate XXX. *read* Plate XXIX.



NEMATODES, MOSTLY AUSTRALIAN AND FIJIAN.

BY N. A. COBB.

(Plates XXXVI.-XLII.)

INTRODUCTORY NOTE.

The following pages contain descriptions, accompanied by about one hundred and seventy figures, of eighty-two species of Nematodes, of which about half have not been hitherto described. In a number of cases the anatomical details have been worked out in a manner worthy the attention of the morphologist.




FIG. 1.—Diagram in explanation of the descriptive formula used for Nematode worms; 6, 7, 8, 10, 6 are the transverse measurements, while 7, 14, 28, 50, 88 are the corresponding longitudinal measurements. The formula in this case is:—

$$\frac{7 \cdot 14 \cdot 28 \cdot 50 \cdot 88}{6 \cdot 7 \cdot 8 \cdot 10 \cdot 6}$$

The unit of measurement is the hundredth part of the length of the worm, whatever that may be. The measurements become, therefore, percentages of the length.

The measurements are taken with the animal viewed in profile; the first is taken at the base of the pharynx, the second at the nerve-ring, the third at the cardiac constriction, the fourth at the vulva in females and at the middle (M) in males, the fifth at the anus.

GENUS MONONCHUS, Bastian.

The genus *Mononchus* is composed of herbivorous free-living Nematodes, not marine, having the average formulæ $\frac{2.5 \cdot 7.6 \cdot 25 \cdot 60 \cdot 90}{2.5 \cdot 2.8 \cdot 3.6 \cdot 3.8 \cdot 2} \cdot 2\text{mm.}$ and $\frac{2.5 \cdot 7.6 \cdot 25 \cdot \text{—M—} \cdot 90}{2.5 \cdot 2.8 \cdot 3.6 \cdot 3.8 \cdot 2} \cdot 2 \cdot (-)\text{mm.}$ They are readily recognised by their capacious pharynx, containing from one to three commonly conspicuous teeth, whose function is, in conjunction with certain minute file-like or rasp-like areas on the wall of the pharynx, to masticate the food, which consists of fresh and succulent vegetable matter, such as rootlets, or the tissues of aquatic or sub-aquatic plants, or the protected tissues found among the sheaths of the leaves of certain land-plants. The thick transparent cuticle of these worms is destitute both of hairs and striæ. The neck is sometimes almost cylindrical, but is usually conoid, and invariably ends in a truncate head, which in one species

Hab.—Found among the bases of carrot leaves, Sydney, New South Wales, Australia, in July.

7. *D. domus Glauci*, n.sp. $\frac{.4}{1} \frac{12}{4.1} \frac{28}{4.7} \frac{.53^{.23}}{5.3} \frac{98.6}{2}$ 1.93 mm. Neck conoid,—convex-conoid anteriorly; lips six, each with at least one distinct papilla; spear acute, well developed; œsophagus at first only one-fifth as wide as the corresponding part of the neck, but gradually and uniformly widening until it is at last more than half as wide as the base of the neck; rectum nearly twice as long as the anal body-diameter; pre-rectum three times the length of the rectum, tapering posteriorly; ovaries reaching two-thirds to three-fourths the way back to the vulva; eggs thick-shelled, ellipsoidal, as long as the body is wide and over half as wide as long.

Hab.—Found among moss on the walls of the Casa Poetæ, Pompeii, Italy. Possibly the widths given in the formula are too great.

8. *D. Vesuvianus*, n.sp. $\frac{.3}{1.1} \frac{10.5}{3.2} \frac{20}{3.5} \frac{.47^{.40}}{4.5} \frac{98.3}{2.3}$ 1.15 mm. In the posterior part the neck is nearly cylindrical, but the anterior third is convex-conoid to the truncate head. The lip-region is not conspicuous, and the lips are either absent or very indistinct. There are perhaps two rows of papillæ, six in each row. The spear is very slender and can be traced backward only for a distance equalling one-fiftieth of the length of the worm. The anterior part of the œsophagus is only about one-fourth as wide as the corresponding part of the neck, but it expands suddenly behind the middle, so that the posterior third (male) or two-fifths (female) is twice as wide, *i.e.*, about half as wide as the base of the neck. The intestine is about half as wide as the body and is marked off from the œsophagus by a distinct constriction. The cardia is broad but not deep. The rectum is only three-fourths as long as the anal body-diameter; the pre-rectal portion of the intestine, however, is quite as long as the enlarged portion of the œsophagus. The lateral fields are nearly one-third as wide as the body. The tail in either sex is hemispherical or conoid-hemispherical. The inconspicuous vulva leads to a vagina half as long as the body is wide. The reflexed ovaries reach half-way back to the vulva. $\frac{.3}{1.3} \frac{11}{3.7} \frac{25}{4.2} \frac{—M—}{4.5} \frac{98.2}{2.5}$ 1 mm. A ventral row of eleven juxtaposed median innervated papillæ extends forward from opposite the proximal ends of the spicula. These latter are of the usual form and about twice as long as the tail. The internal male sexual organs occupy the posterior two-thirds of the body.

Hab.—Found among moss on the sides of Mount Vesuvius, Italy.

9. *D. labyrinthostoma*, n.sp. $\frac{1}{1} \frac{7}{1.9} \frac{29}{2.2} \frac{.50^{.9}}{2.2} \frac{90}{1.2}$ 1.75 mm. This species belongs to the group first made known through the researches of Dr. de Man, and characterised by the possession of elaborate mouth parts accessory to the spear. The group represents the acme of development in *Dorylaimus* so far as yet discovered. The spear presents