substance: gallium sulfide (GaS)

property: band structure, direct energy gap

The best known polytype is β -GaS.


band structure

see Fig. 1; Brillouin zone, Fig. 2; XPS-spectrum, Fig. 3.

GaS is an indirect gap semiconductor. The electronic band structure is very similar to that of GaSe. The valence band maximum is at the Γ -point The conduction band is characterized by 6 equivalent minima at the M point.

direct gap

$E_{\rm g,dir}(1_{\rm 4v}^{-}-1_{\rm 3c}^{-})$	3.050(2) eV	T = 77 K	optical absorption	69A
-	3.065 eV	T = 77 K	electroabsorption	77G
pressure coefficient of $E_{ m g,dir}$				





References:

- 69A Aulich, E., Brebner, J. L., Mooser, E.: Phys. Status Solidi 31 (1969) 129.
- Guseinov, G. D., Abdullayeva, S. G., Aksianov, I. G., Gajiyev, V. A.: Mater. Res. Bull. 12 (1977) 207.
- 78M Mejatty, M., Segura, A., LeToullec, R., Besson, J. M.: J. Phys. Chem. Solids 39 (1978) 25.
- Antonangeli, F., Piacentini, M., Baizarotti, A., Grasso, V., Girlanda, R., Doni, E.: Nuovo Cimento B 51 (1979) 181.
- 81D Depeursinge, Y.: Nuovo Cimento B 64 (1981) 111.



Fig. 1. GaS. Empirical pseudopotential band structure for the $\beta\text{--polytype}$ [81D].






Fig. 2. Brillouin zone for the hexagonal lattice. g_1 , g_2 , g_3 : reciprocal basis vectors; $g_1 \parallel k_x$, $g_3 \parallel k_z$.






Fig. 3.

GaS, GaSe, InSe. XPS-number of electrons vs. electron binding energy. The zero of energy corresponds to the upper valence band edge. Peaks α_3 and G are spurious and most probably due to the unmonochromatized AlK_{α} X-ray source used in the experiment. Peak A is the contribution of the uppermost valence bands which are essentially formed by the chalcogen s–p_z-orbitals. Peak B corresponds to the contribution of the chalcogen p_{xy} states, peaks E, C to that of the cation s-states and peak F to that of the chalcogen s-electrons (S3s, Se4s) [79A].

