

substance: gallium sulfide (GaS)

property: indirect energy gap and exciton energies

indirect energy gap

The values quoted in the literature correspond to the ground state of the indirect free exciton (see [77D1] and the transition energy of the indirect exciton, see below) . The binding energy of the indirect exciton is not known.

indirect exciton transition energy

$E_{\text{gx,ind}}$	2.593 eV	$T = 1.5 \text{ K}$	optical absorption	77D2
	2.600 eV	$T = 4.2 \text{ K}$	photoluminescence for $\text{GaS}_x\text{Se}_{1-x}$	80S
	2.5947(6) eV	$T = 4.2 \text{ K}$	photoluminescence	73R
	2.576(1) eV	$T = 77 \text{ K}$	wavelength modulated absorption	75K

For temperature dependence of indirect exciton transition energy, see Fig. 1.

pressure coefficient of $E_{\text{gx,ind}}$

$dE_{\text{gx,ind}}/dp$	$-13.5 \cdot 10^{-6} \text{ eV bar}^{-1}$	$T = 300 \text{ K}$	photoconductivity	69N
	$-11.0(15) \cdot 10^{-6} \text{ eV bar}^{-1}$	$T = 300 \text{ K},$ $p < 6 \text{ kbar}$	optical absorption, see Fig. 2	78M
	$-4 \cdot 10^{-6} \text{ eV bar}^{-1}$	$T = 300 \text{ K},$ $p \approx 40 \text{ kbar}$		

pressure coefficient of $E_{\text{gx,ind}}$

$dE_{\text{gx,ind}}/dp$	$-3.1 \cdot 10^{-11} \text{ eV Pa}^{-1}$	$T = 4.2 \text{ K}$ $0 < p \leq 80 \text{ MPa}$	optical absorption, uniaxial pressure	83S
-------------------------	--	--	---------------------------------------	-----

g -value of indirect free triplet exciton

g_{\parallel}	2.006(2)		optical detected magnetic resonance	79D
-----------------	----------	--	-------------------------------------	-----

References:

- 69N Niilisk, A. J., Kirs, J. J.: Phys. Status Solidi 31 (1969) K91.
- 73R Razbirin, B. S., Karaman, M. I., Mushinskii, V. P., Starukhin, A. N.: Fiz. Tekh. Poluprovodn. 7 (1973) 1112; Sov. Phys. Semicond. (English Transl.) 7 (1973) 753.
- 75K Karaman, M. I., Mushinskii, V. P.: Fiz. Tekh. Poluprovodn. 9 (1975) 1415; Sov. Phys. Semicond. (English Transl.) 9 (1975) 934.
- 77D1 Depeursinge, C., Le Chi Thanh: Proc. 13th Intern. Conf. Phys. Semicond., Rome 1976, ed. by F. Fumi, Veschi, Roma, 1977, p. 383.
- 77D2 Depeursinge, C.: Thesis, Lausanne 1977, unpublished.
- 78M Mejatty, M., Segura, A., LeToullec, R., Besson, J. M.: J. Phys. Chem. Solids 39 (1978) 25.
- 79D Dawson, P., Killoran, N., Cavenett, B. C.: Solid State Commun. 32 (1979) 1133.
- 80S Serizawa, H., Sasaki, Y., Nishina, Y.: J. Phys. Soc. Jpn. 48 (1980) 490.
- 83S Sasaki, Y., Hoshi, K., Saito, S., Yamaguchi, K., Nishina, Y.: J. Phys. Soc. Jpn. 52 (1983) 3706.



Fig. 1.

GaS. Energetic position of indirect exciton ground state transition vs. temperature [77D2].




Fig. 2.

GaS. Absorption coefficient for $E \perp c$ (K_{\perp}) at 300 K vs. photon energy for different hydrostatic pressures, 1, 1 bar; 2, 10 kbar; 3, 19.5 kbar; 4, 29.5 kbar; 5, 43.5 kbar [78M].

