


Refinement of the crystal structure of digallium trisulfide, Ga₂S₃

C. Y. Jones^{*I}, J. C. Bryan^{II}, K. Kirschbaum^{III} and J. G. Edwards^{III}^I Oak Ridge National Laboratory, Metals and Ceramics Division, Oak Ridge, TN 37831, USA^{II} Oak Ridge National Laboratory, Chemical and Analytical Sciences Division, Oak Ridge, TN 37831, USA^{III} University of Toledo, Department of Chemistry, Toledo, OH 43606, USA

Received February 21, 2001, CSD-No. 409550



● Ga

Abstract

Ga₂S₃, monoclinic, *C1c1* (No. 9), *a* = 11.107(2) Å, *b* = 6.395(1) Å, *c* = 7.021(1) Å, β = 121.17(3)°, *V* = 426.7 Å³, *Z* = 4, *R*_{gt}(*F*) = 0.035, *wR*_{ref}(*F*²) = 0.084, *T* = 294 K.

Source of material

A pale, yellow crystal of Ga₂S₃ was selected from crystals grown by vapor deposition from Ga₂S₃ powder in an effusion cell [1]. The selected crystal had smooth, uniformly-reflecting facet surfaces.

Experimental details

The unit cell parameters and atomic coordinates from this determination agree with the previously published structure [2], but are systematically smaller, possibly because of a difference in the temperature. An extinction coefficient, 0.142, is reported here for the first time.

Discussion

Gallium sulfide, Ga₂S₃, is a III–VI semiconductor whose monoclinic phase possesses at room-temperature a blue-green luminescence [3–5]. Often referred to as α-Ga₂S₃, the monoclinic Ga₂S₃ crystallizes in space group *Cc* and contains ordered vacancies [2]. In addition to its luminescence properties, Ga₂S₃ also exhibits partial vapor pressures of Ga₂S(g) that increase with decreasing temperature and decrease with increasing temperature in the range 1230 K < *T* < 1260 K, where a phase transformation was postulated [6]. Its anomalous vapor-pressure behavior has made Ga₂S₃ a model system for the study of a class of high-temperature condensed-phase transformations in it and other sulfides and chalcogenides exhibiting anomalous temperature dependence of vapor pressure. Recent data, from the first time-of-flight neutron powder diffraction experiments performed on samples undergoing effusion [1], revealed a disordering, monoclinic-to-hexagonal phase transformation in the same temperature range as the anomalous vapor pressure, 1240 K – 1260 K [7]. For the monoclinic phase of Ga₂S₃, four single-crystal structures have been published [2,8,9,10]. Work on the high-temperature structures of Ga₂S₃ determined by neutron scattering has made it desirable to reanalyze its single-crystal structure at a specified temperature. The figure shows the corner-sharing GaS₄ tetrahedra of the structure as viewed along the [001] direction and the channels created by the ordered vacancies.

Table 1. Data collection and handling.

Crystal:	yellow, irregular, size 0.30 × 0.30 × 0.40 mm
Wavelength:	Mo K _α radiation (0.71073 Å)
μ:	13.87 cm ^{−1}
Diffraction, scan mode:	Enraf-Nonius CAD4, ω/2θ
2θ _{max} :	52°
<i>N</i> (<i>hkl</i>) _{measured} , <i>N</i> (<i>hkl</i>) _{unique} :	436, 436
Criterion for <i>I</i> _{obs} , <i>N</i> (<i>hkl</i>) _{gt} :	<i>I</i> _{obs} > 2 σ(<i>I</i> _{obs}), 432
<i>N</i> (<i>param</i>) _{refined} :	47
Programs:	SHEXTL [11], SHELXL-97 [12], PLATON [13]

Table 2. Atomic coordinates and displacement parameters (in Å²).

Atom	Site	<i>x</i>	<i>y</i>	<i>z</i>	<i>U</i> ₁₁	<i>U</i> ₂₂	<i>U</i> ₃₃	<i>U</i> ₁₂	<i>U</i> ₁₃	<i>U</i> ₂₃
Ga(1)	4 <i>a</i>	0.04213(8)	0.4023(2)	0.1251(1)	0.0068(7)	0.0130(6)	0.0096(7)	0.0007(4)	0.0042(5)	0.0005(4)
Ga(2)	4 <i>a</i>	0.20026(9)	0.9321(2)	0.1156(1)	0.0065(7)	0.0133(6)	0.0102(7)	−0.0003(4)	0.0045(5)	−0.0012(4)
S(1)	4 <i>a</i>	−0.0050(3)	1.0861(4)	−0.0162(6)	0.007(1)	0.016(1)	0.018(1)	−0.0001(8)	0.004(1)	−0.0048(9)
S(2)	4 <i>a</i>	0.3378(3)	0.9075(3)	0.4999(4)	0.004(1)	0.015(1)	0.010(2)	−0.0009(8)	0.004(1)	−0.0019(8)
S(3)	4 <i>a</i>	0.1741(3)	0.5840(3)	0.0084(4)	0.008(1)	0.012(1)	0.010(1)	−0.0017(8)	0.006(1)	−0.0010(8)

* Correspondence author (e-mail: jonescy@ornl.gov)

Acknowledgment. We thank the College of Arts and Sciences of the University of Toledo for use of the X-ray diffractometer and computer facilities in the Arts and Sciences Instrumentation Center.

References

1. Jones, C. Y.; Edwards, J. G.: Observation of a Phase Transformation in Ga_2S_3 in an Quartz Effusion Cell above 1230 K by means of Neutron Scattering. *J. Phys. Chem. B* **105** (2001) 2718-2724.
2. Goodyear, J.; Steigmann, G. A.: The Crystal Structure of $\alpha\text{-Ga}_2\text{S}_3$. *Acta Crystallogr.* **16** (1963) 946-949.
3. Lee, J.-S.; Won, Y.-H.; Kim, H.-N.; Kim, C.-D.; Kim, W.-T.: Photoluminescence of Ga_2S_3 and $\text{Ga}_2\text{S}_3\text{:Mn}$ Single Crystals. *Sol. State. Commun.* **97** (1996) 1101-1104.
4. Márquez, F.; Fornés, V.: Synthesis and characterization of Ga_2S_3 semiconductor included in zeolite Y. *Sol. State Commun.* **112** (1999) 17-20.
5. Kadono, K.; Higuchi, H.; Takahashi, M.; Kawamoto, Y.; Tanaka, H.: Upconversion luminescence of Ga_2S_3 -based sulfide glasses containing Er^{3+} ions. *J. Noncryst. Solids* **184** (1995) 309-313.
6. Roberts, J. A.; Searcy, A.: Anomalous Temperature Dependence for a Partial Vapor Pressure. *J. Phys. Chem. B* **99** (1995) 525-527.
7. Edwards, J. G.; Uram, R. S.: Multiple Equilibrium States in Effusion Cells: Catastrophic Phase Transitions. *J. Phys. Chem. B* **96** (1992) 8561-8568.
8. Goodyear, J.; Duffin, W. A.; Steigmann, G. A.: [The Unit Cell of \$\alpha\text{-Ga}_2\text{S}_3\$](#) . *Acta Crystallogr.* **14** (1961) 1168-1170.
9. Hahn, H.; Frank, G.: Zur Struktur des Ga_2S_3 . *Z. Anorg. Allgem. Chem.* **278** (1955) 333-339.
10. Tomas, A.; Pardo, M. P.; Guittard, M.: Structural Determination of α and $\beta\text{-Ga}_2\text{S}_3$. *Mat. Res. Bull.* **22** (1987) 1549-1554.
11. SHELXTL 5.10/IRIX. Bruker Analytical X-ray Instruments Inc. Madison, Wisconsin, USA 1997.
12. Sheldrick, G. M.: SHELXL-97. Program for Crystal Structure Refinement. Universität Göttingen, Germany 1997.
13. Spek, A. L.: PLATON. A Multi-Purpose Crystallographic Tool, Universiteit Utrecht, The Netherlands 2000.