



# Engineering Physics

## (PHY1701)


**Dr. B. Ajitha**

Assistant Professor  
Division of Physics  
VIT University  
Chennai, India  
[ajitha.b@vit.ac.in](mailto:ajitha.b@vit.ac.in)

## Contents

- Introduction to Nano-materials,
- Moore's law,
- Properties of Nano-materials\* ,
- Quantum confinement, Quantum well, wire & dot (CF 226-236),
- Carbon Nano-tubes (CNT) (CF 115-125), &
- Applications of nanotechnology in industry

❖ Introduction to Nano Technology, Charles P Poole Jr., Frank J Owens, Willey Publication. (CF)



- The surface to volume ratio follows an inverse power law
- The fraction of atoms on the surface increases with decreasing the size
- Large surface is favorable for applications such as catalysis.
- As a particle decreases in size, a greater proportion of atoms are found at the surface compared to those inside.
- For example, a particle of
  - Size-30 nm- $\rightarrow$  5% of its atoms on its surface
  - Size-10 nm- $\rightarrow$  20% of its atoms on its surface
  - Size-3 nm- $\rightarrow$  50% of its atoms on its surface

**Nanoparticles are more reactive than large particles (Catalyst)**

Cubic:

Surface area  $S = 6a^2$

Volume  $V = a^3$

Surface area to volume ratio

$$\frac{S}{V} = \frac{6a^2}{a^3} = \frac{6}{a}$$

Sphere

Surface area  $S = 4\pi R^2$

Volume of sphere  $V = \frac{4}{3}\pi R^3$

Surface to volume ratio  $\frac{S}{V} = \frac{4\pi R^2}{\frac{4}{3}\pi R^3} = \frac{3}{R}$

# Increase in surface area

Imagine a sphere of radius  $R=1\text{cm}$ . How many no. of tiny spheres of radius  $r=1\text{nm}$  is possible from the material?

$$\frac{4}{3}\pi R^3 = \frac{4}{3}\pi r^3$$

$$\text{i.e. } R^3 = N r^3$$

$$N = \frac{R^3}{r^3} = \frac{1\text{cm}^3}{1\text{nm}^3} = \frac{10^{-6}\text{m}^3}{10^{-27}\text{m}^3} = 10^{21}$$

$10^{21}$  tiny 1 nm spheres are formed.

# Increase in surface area

Total surface area available on these particles:

Surface area of the big sphere

$$S = 4\pi R^2 = 4\pi (10^{-2} \text{ m})^2 = 4\pi (10^{-4}) \text{ m}^2$$

Surface area of the tiny sphere

$$S = 4\pi r^2 = 4\pi (10^{-9} \text{ m})^2 = 4\pi (10^{-18}) \text{ m}^2$$

There are  $10^{21}$  numbers of tiny spheres,  
so their total surface area

$$S_t = 10^{21} \times 4\pi (10^{-18}) \text{ m}^2 = 4\pi 10^3 \text{ m}^2$$

$$\text{So, } \frac{S_t}{S} = \frac{4\pi \times 10^3 \text{ m}^2}{4\pi \times 10^{-4} \text{ m}^2} = 1 \times 10^7$$

The surface area has increased by a factor of 10 million times.

# Increase in surface area

- Nanoparticles exhibit unique properties due to their high surface area to volume ratio.
- A spherical particle has a diameter (D) of 100nm.
  - Calculate the volume (V) and surface area (SA)

$$V = \frac{4}{3} \pi r^3 = \frac{\pi D^3}{6}$$

$$V = \frac{\pi(100 \times 10^{-9})^3}{6}$$

$$V = 5.24 \times 10^{-22} \text{ m}^3$$


$$SA = 4\pi r^2 = \pi D^2$$

$$SA = \pi(100 \times 10^{-9})^2$$

$$SA = 3.141 \times 10^{-14} \text{ m}^2$$



# Increase in surface area



8 Cubes Side  $L$   
Each has Surface area  $6L^2$   
Total Surface Area  $48L^2$


1 Cube  
Length of sides  $2L$   
Surface area  $24L^2$



For example, 5 cubic centimeters - about 1.7 cm per side - of material divided 24 times will produce 1 nanometer cubes and spread in a single layer could cover a football field


- Quantum effects can begin to dominate the behaviour of matter at the nanoscale – particularly at the lower end - affecting the optical, electrical and magnetic behaviour of materials.
- When the particle size is comparable to the debroglie wavelength of electrons or mean free path of electrons, the energy levels of electron changes leading to an effect – quantum confinement characteristics.
- The quantum confinement effect can be observed once the diameter of the particle is of the same magnitude as then wavelength of the electron Wave function.
- Quantum confinement is responsible for the increase of energy difference between energy states and band gap.

- A phenomenon tightly related with the optical and electronic properties of the materials.
- When materials are this small, their electronic and optical properties deviate substantially from those of bulk materials.(GOLD)




## Melting point as a function of particle size

- Nanoparticles have a lower melting point than their bulk counterparts




Melting point of gold nanoparticles as a function of size.



## Definition:

- Quantum Confinement is the spatial confinement of electron-hole pairs (excitons) in one or more dimensions within a material.
  - 1D confinement: Quantum Wells
  - 2D confinement: Quantum Wire
  - 3D confinement: Quantum Dot
- Quantum confinement is more prominent in semiconductors because they have an energy gap in their electronic band structure.
- Metals do not have a bandgap, so quantum size effects are less prevalent. Quantum confinement is only observed at dimensions below 2 nm.

- Energy level spacing and quantum confinement
  - The reduction in the number of atoms in a material results in the confinement of normally delocalized energy states.
  - Electron-hole pairs become spatially confined when the dimensions of a nanoparticle approach the de Broglie wavelength of electrons in the conduction band.
  - As a result the spacing between energy bands of semiconductor or insulator is **increased** (Similar to the *particle in a box* scenario, of introductory quantum mechanics.)





- Recall that when atoms are brought together in a bulk material the number of energy states increases substantially to form nearly continuous bands of states.







- This is very similar to the famous particle-in-a-box scenario and can be understood by examining the Heisenberg Uncertainty Principle.
- The Uncertainty Principle states that the more precisely one knows the position of a particle, the more uncertainty in its momentum (and vice versa).
- Therefore, the more spatially confined and localized a particle becomes, the broader the range of its momentum/energy.
- This is manifested as an increase in the average energy of electrons in the conduction band = increased energy level spacing = larger bandgap
- The bandgap of a spherical quantum dot is increased from its bulk value by a factor of  $1/R^2$ , where  $R$  is the particle radius.\*

no confinement  
bulk  
(3D materials)





bulk

1D confinement  
Q-well/ultrathin film  
(2D materials)





quantum well

2D confinement  
Q-wire  
(1D materials)



quantum wire

3D confinement  
Q-dot  
(0D material)




quantum dot

- Quantum effects arise in systems which confine electrons to regions comparable to their de Broglie wavelength.
- When such confinement occurs in one dimension only (say, by a restriction on the motion of the electron in the  $z$ -direction), with free motion in the  $x$ - and  $y$ -directions, a two-dimensional system is created.




(a) Q-Film



## Quantum Well





# Quantum Well



# Quantum wire

- Quantum effects in systems which confine electrons to regions comparable to their de Broglie wavelength.
- When such confinement occurs in two dimensions only (say, by two restrictions on the motion of the electron in the z- and y-directions), with free motion in the x-direction, a one dimensional electron is created.

(b) Q-Wire



Bulk



Well



Wire






Dot

# Quantum Dots


## What does this mean?

- Quantum dots are bandgap tunable by size. We can engineer their optical and electrical properties.
- Smaller QDs have a large bandgap.
- Absorbance and luminescence spectrums are blue shifted with decreasing particle size.

(c) Q-Box or Q-Dot




- Absorption and emission occur at specific wavelengths, which are related to QD size.





# Quantum Dots


- Quantum dots are usually regarded as semiconductors by definition. Similar behavior is observed in some metals.
- Therefore, in some cases it may be acceptable to speak about metal quantum dots.
- Typically, quantum dots are composed of groups II-VI, III-V, and IV-VI materials.
- QDs are bandgap tunable by size which means their optical and electrical properties can be engineered to meet specific applications.
- Nanocrystals (2-10 nm) of semiconductor compounds
- Small size leads to confinement of excitons (electron-hole pairs)
- Quantized energy levels and altered relaxation dynamics
- Examples: CdSe, PbSe, PbTe, InP







# Quantum Dots



# Moore's Law

- The number of transistors on a chip will approximately double every 18 to 24 months (Moore's Law).
- This law has given chip designers greater incentives to incorporate new features on silicon.
- Moore's Law works largely through shrinking transistors, the circuits that carry electrical signals.
- By shrinking transistors, designers can squeeze more transistors into a chip



- Moore's second law predicts that the cost of building a chip manufacturing plant doubles with every other chip generation or roughly every 36 months.

