




Fundamentos em Redes Neurais

Regressão Linear Univariada

Profº - Dr. Thales Levi Azevedo Valente

thales.l.a.valente@gmail.com.br

Sejam Bem-vindos !




**Os celulares devem
ficar no silencioso
ou desligados**

Pode ser utilizado
apenas em caso
de emergência



**Boa tarde/noite, por
favor e com licença
DEVEM ser usados**

Educação é
essencial




Machine Learning

Linear regression with one variable

Model representation

Housing Prices (Portland, OR)

Price
(in 1000s
of dollars)



Supervised Learning

Given the “right answer” for each example in the data.

Regression Problem

Predict real-valued output

Classification: Discrete-valued output

Training set of housing prices (Portland, OR)

Size in feet ² (x)	Price (\$) in 1000's (y)
→ 2104	460
1416	232
→ 1534	315
852	178
...	...

$m = 47$

Notation:

- m = Number of training examples
- x 's = "input" variable / features
- y 's = "output" variable / "target" variable

(x, y) - one training example

$(x^{(i)}, y^{(i)})$ - i^{th} training example

$$\begin{cases} x^{(1)} = 2104 \\ x^{(2)} = 1416 \\ y^{(1)} = 460 \end{cases}$$

Training Set

Learning Algorithm

Size of house

h

Estimated price

hypothesis


(estimated value of y)

h maps from x 's to y 's.

How do we represent h ?


$$h_{\theta}(x) = \theta_0 + \theta_1 x$$

Shorthand: $h(x)$



Linear regression with one variable. (x)
Univariate linear regression.

↳ one variable




Machine Learning

Linear regression with one variable

Cost function

Training Set

Size in feet ² (x)	Price (\$) in 1000's (y)
2104	460
1416	232
1534	315
852	178
...	...


 $m = 47$

Hypothesis: $h_{\theta}(x) = \theta_0 + \theta_1 x$

θ_i 's: Parameters


How to choose θ_i 's ?

$h_{\theta}(x) = \theta_0 + \theta_1 x$




$\rightarrow \theta_0 = 1.5$

$\rightarrow \theta_1 = 0$




$\rightarrow \theta_0 = 0$

$\rightarrow \theta_1 = 0.5$



$\rightarrow \theta_0 = 1$

$\rightarrow \theta_1 = 0.5$



$(x^{(i)}, y^{(i)})$

Idea: Choose θ_0, θ_1 so that $\underline{h_\theta(x)}$ is close to $\underline{y}$ for our training examples (x, y)

x, y

minimize θ_0, θ_1

$\frac{1}{2m} \sum_{i=1}^m \left(\underline{h_\theta(x^{(i)})} - \underline{y^{(i)}} \right)^2$

$\# \text{ training examples}$


$h_\theta(x^{(i)}) = \theta_0 + \theta_1 x^{(i)}$

$$J(\theta_0, \theta_1) = \frac{1}{2m} \sum_{i=1}^m (h_\theta(x^{(i)}) - y^{(i)})^2$$

minimize θ_0, θ_1 $J(\theta_0, \theta_1)$

Cost function

Squared error function



Machine Learning

Linear regression
with one variable


Cost function
intuition I

Hypothesis:

$$\underline{h_{\theta}(x) = \theta_0 + \theta_1 x}$$

Parameters:

$$\underline{\theta_0, \theta_1}$$



Cost Function:

$$\rightarrow J(\theta_0, \theta_1) = \frac{1}{2m} \sum_{i=1}^m (h_{\theta}(x^{(i)}) - y^{(i)})^2$$


Goal: minimize $J(\theta_0, \theta_1)$
 $\nearrow \theta_0, \theta_1$

Simplified

$$h_{\theta}(x) = \underline{\theta_1 x}$$

$$\theta_0 = 0$$

$$\underline{\theta_1}$$




$$J(\theta_1) = \frac{1}{2m} \sum_{i=1}^m (\underline{h_{\theta}(x^{(i)})} - y^{(i)})^2$$

minimize $\underline{J(\theta_1)}$ $\theta_1, x^{(i)}$

→ $h_{\theta}(x)$

(for fixed θ_1 , this is a function of x)




$J(\theta_1)$ = $\frac{1}{2m} \sum_{i=1}^m (h_{\theta}(x^{(i)}) - y^{(i)})^2$

= $\frac{1}{2m} \sum_{i=1}^m (\theta_1 x^{(i)} - y^{(i)})^2 = \frac{1}{2m} (0^2 + 0^2 + 0^2) = 0^2$

→ $J(\theta_1)$

(function of the parameter θ_1)




$\theta_1 = 0.5?$

$J(1) = 0$

$$h_{\theta}(x)$$

(for fixed θ_1 , this is a function of x)




$$J(0.5) = \frac{1}{2m} [(0.5-1)^2 + (1-2)^2 + (1.5-3)^2]$$

$$= \frac{1}{2 \times 3} (3.5) = \frac{3.5}{6} \approx \underline{0.58}$$

$$J(\theta_1)$$

(function of the parameter θ_1)




$$\theta_1 = 0?$$

$$J(0) = ?$$

$$h_{\theta}(x)$$

(for fixed θ_1 , this is a function of x)




$$J(0) = \frac{1}{2m} (1^2 + 2^2 + 3^2) = \frac{1}{6} \cdot 14 \approx 2.3$$


$$h(x) = -0.5x$$

minimize $J(\theta_1)$

$$J(\theta_1)$$

(function of the parameter θ_1)





Machine Learning

Linear regression with one variable

Cost function intuition II

Hypothesis: $h_{\theta}(x) = \theta_0 + \theta_1 x$


Parameters: θ_0, θ_1

Cost Function: $J(\theta_0, \theta_1) = \frac{1}{2m} \sum_{i=1}^m \left(h_{\theta}(x^{(i)}) - y^{(i)} \right)^2$

Goal: minimize $J(\theta_0, \theta_1)$
 θ_0, θ_1

$$\underline{h_{\theta}(x)}$$


(for fixed θ_0, θ_1 this is a function of x)




$$h_{\theta}(x) = 50 + 0.06x$$

$$\underline{J(\theta_0, \theta_1)}$$

(function of the parameters θ_0, θ_1)




Contour plots
Contour figures -



$J(\theta_0, \theta_1)$


$$h_{\theta}(x)$$

(for fixed θ_0, θ_1 this is a function of x)



$$J(\theta_0, \theta_1)$$


(function of the parameters θ_0, θ_1)



θ_0, θ_1


$$h_{\theta}(x)$$

(for fixed θ_0, θ_1 this is a function of x)



$$J(\theta_0, \theta_1)$$


(function of the parameters θ_0, θ_1)



$$\begin{cases} \theta_0 = 360 \\ \theta_1 = 0 \end{cases}$$


$$h_{\theta}(x)$$

(for fixed θ_0, θ_1 this is a function of x)




$$J(\theta_0, \theta_1)$$

(function of the parameters θ_0, θ_1)




$$h_{\theta}(x)$$


(for fixed θ_0, θ_1 this is a function of x)



$$J(\theta_0, \theta_1)$$

(function of the parameters θ_0, θ_1)





Machine Learning

Linear regression
with one variable


Gradient
descent


Have some function $J(\theta_0, \theta_1)$ $J(\theta_0, \theta_1, \theta_2, \dots, \theta_n)$

Want $\min_{\theta_0, \theta_1} J(\theta_0, \theta_1)$ $\min_{\theta_0, \dots, \theta_n} J(\theta_0, \dots, \theta_n)$

Outline:

- Start with some θ_0, θ_1 (say $\theta_0 = 0, \theta_1 = 0$)
- Keep changing θ_0, θ_1 to reduce $J(\theta_0, \theta_1)$
until we hopefully end up at a minimum





Gradient descent algorithm

repeat until convergence {

$$\theta_j := \theta_j - \alpha \frac{\partial}{\partial \theta_j} J(\theta_0, \theta_1)$$

learning rate

(for $j = 0$ and $j = 1$)

Simultaneously update θ_0 and θ_1

Correct: Simultaneous update

$$\rightarrow \text{temp0} := \theta_0 - \alpha \frac{\partial}{\partial \theta_0} J(\theta_0, \theta_1)$$

$$\rightarrow \text{temp1} := \theta_1 - \alpha \frac{\partial}{\partial \theta_1} J(\theta_0, \theta_1)$$

$$\rightarrow \theta_0 := \text{temp0}$$

$$\rightarrow \theta_1 := \text{temp1}$$

Incorrect:

$$\rightarrow \text{temp0} := \theta_0 - \alpha \frac{\partial}{\partial \theta_0} J(\theta_0, \theta_1)$$

$$\rightarrow \theta_0 := \text{temp0}$$

$$\rightarrow \text{temp1} := \theta_1 - \alpha \frac{\partial}{\partial \theta_1} J(\theta_0, \theta_1)$$

$$\rightarrow \theta_1 := \text{temp1}$$

Assignment


$$a = b$$

$$a = a + 1$$

Truth assertion

$$a = b$$

$$a = a + 1$$



Machine Learning

Linear regression with one variable

Gradient descent intuition

Gradient descent algorithm

repeat until convergence {

→ $\theta_j := \theta_j - \alpha \frac{\partial}{\partial \theta_j} J(\theta_0, \theta_1)$


learning
rate

derivative

(simultaneously update
 $j = 0$ and $j = 1$)


$$\min_{\theta_1} J(\theta_1)$$

$$\theta_1 \in \mathbb{R}.$$



$$\theta_1 := \theta_1 - \underbrace{\alpha}_{\substack{\leftarrow \\ \frac{\partial}{\partial \theta_1} J(\theta_1)}} \underbrace{\frac{\partial}{\partial \theta_1} J(\theta_1)}_{\geq 0}$$

$$\theta_1 := \theta_1 - \underline{\alpha} \cdot (\text{positive number})$$





$$\underbrace{\frac{\partial}{\partial \theta_1} J(\theta_1)}_{\leq 0}$$


$$\theta_1 := \underbrace{\theta_1}_{\uparrow} - \alpha \underbrace{(\text{negative number})}_{\uparrow}$$

$$\theta_1 := \theta_1 - \alpha \frac{\partial}{\partial \theta_1} J(\theta_1)$$

If α is too small, gradient descent
can be slow.

If α is too large, gradient descent
can overshoot the minimum. It may
fail to converge, or even diverge.






Current value of θ_1


$$\theta_1 := \theta_1 - \alpha \frac{d}{d\theta_1} J(\theta_1)$$

Gradient descent can converge to a local minimum, even with the learning rate α fixed.

$$\theta_1 := \theta_1 - \alpha \frac{d}{d\theta_1} J(\theta_1)$$

As we approach a local minimum, gradient descent will automatically take smaller steps. So, no need to decrease α over time.





Machine Learning

Linear regression
with one variable

Gradient descent for
linear regression

Gradient descent algorithm

repeat until convergence {
 $\theta_j := \theta_j - \alpha \frac{\partial}{\partial \theta_j} J(\theta_0, \theta_1)$
 (for $j = 1$ and $j = 0$)
}

Linear Regression Model

$$h_{\theta}(x) = \theta_0 + \theta_1 x$$

$$J(\theta_0, \theta_1) = \frac{1}{2m} \sum_{i=1}^m (h_{\theta}(x^{(i)}) - y^{(i)})^2$$

$$\frac{\partial}{\partial \theta_j} \underline{J(\theta_0, \theta_1)} = \frac{2}{2\theta_j} \underline{\frac{1}{2m} \sum_{i=1}^m (\underline{h_{\theta}(x^{(i)}) - y^{(i)}})^2}$$

$$= \frac{2}{2\theta_j} \frac{1}{2m} \sum_{i=1}^m (\underline{\theta_0 + \theta_1 x^{(i)} - y^{(i)}})^2$$

$$j = 0 : \underline{\frac{\partial}{\partial \theta_0} J(\theta_0, \theta_1)} = \frac{1}{m} \sum_{i=1}^m (h_{\theta}(x^{(i)}) - y^{(i)})$$

$$j = 1 : \underline{\frac{\partial}{\partial \theta_1} J(\theta_0, \theta_1)} = \frac{1}{m} \sum_{i=1}^m (h_{\theta}(x^{(i)}) - y^{(i)}) \cdot x^{(i)}$$

Gradient descent algorithm

repeat until convergence {

$$\theta_0 := \theta_0 - \alpha \frac{1}{m} \sum_{i=1}^m (h_{\theta}(x^{(i)}) - y^{(i)})$$


$$\theta_1 := \theta_1 - \alpha \frac{1}{m} \sum_{i=1}^m (h_{\theta}(x^{(i)}) - y^{(i)}) \cdot x^{(i)}$$


}

$$\frac{\partial}{\partial \theta_0} J(\theta_0, \theta_1)$$

update
 θ_0 and θ_1
simultaneously

$$\frac{\partial}{\partial \theta_1} J(\theta_0, \theta_1)$$






"Convex function"

Bowl-shaped


$$\underline{h_{\theta}(x)}$$

(for fixed θ_0, θ_1 this is a function of x)




$$\underline{J(\theta_0, \theta_1)}$$

(function of the parameters θ_0, θ_1)




$$h_{\theta}(x)$$

(for fixed θ_0, θ_1 this is a function of x)




$$J(\theta_0, \theta_1)$$

(function of the parameters θ_0, θ_1)




$$h_{\theta}(x)$$

(for fixed θ_0, θ_1 this is a function of x)




$$J(\theta_0, \theta_1)$$

(function of the parameters θ_0, θ_1)




$$h_{\theta}(x)$$

(for fixed θ_0, θ_1 this is a function of x)




$$J(\theta_0, \theta_1)$$

(function of the parameters θ_0, θ_1)




$$h_{\theta}(x)$$

(for fixed θ_0, θ_1 this is a function of x)




$$J(\theta_0, \theta_1)$$

(function of the parameters θ_0, θ_1)




$$h_{\theta}(x)$$

(for fixed θ_0, θ_1 this is a function of x)




$$J(\theta_0, \theta_1)$$

(function of the parameters θ_0, θ_1)




$$h_{\theta}(x)$$

(for fixed θ_0, θ_1 this is a function of x)




$$J(\theta_0, \theta_1)$$

(function of the parameters θ_0, θ_1)




$$h_{\theta}(x)$$

(for fixed θ_0, θ_1 this is a function of x)




$$J(\theta_0, \theta_1)$$

(function of the parameters θ_0, θ_1)




$$h_{\theta}(x)$$

(for fixed θ_0, θ_1 this is a function of x)



$$J(\theta_0, \theta_1)$$

(function of the parameters θ_0, θ_1)



“Batch” Gradient Descent

“Batch”: Each step of gradient descent uses all the training examples.


$$\rightarrow \sum_{i=1}^m (h_{\theta}(x^{(i)}) - y^{(i)})$$

Referências

NG, Andrew. *Machine Learning*. Stanford University, 2011. Curso oferecido via Coursera. Disponível em: <https://www.coursera.org/learn/machine-learning>.



Dúvidas?



Até a próxima...



Apresentador

Thales Levi Azevedo Valente

E-mail:

thales.l.a.valente@gmail.com