




Fundamentos em Redes Neurais

Introdução à Conceitos Avançados de
Regressão

Profº - Dr. Thales Levi Azevedo Valente
thales.l.a.valente@gmail.com.br

Sejam Bem-vindos !




**Os celulares devem
ficar no silencioso
ou desligados**

Pode ser utilizado
apenas em caso
de emergência



**Boa tarde/noite, por
favor e com licença
DEVEM ser usados**

Educação é
essencial



Machine Learning

Linear Regression with multiple variables

Multiple features

Multiple features (variables).

Size (feet ²)	Price (\$1000)
x	y
2104	460
1416	232
1534	315
852	178
...	...

$$\underline{h_{\theta}(x) = \theta_0 + \theta_1 x}$$

Multiple features (variables).

Size (feet ²)	Number of bedrooms	Number of floors	Age of home (years)	Price (\$1000)
x_1	x_2	x_3	x_4	y
2104	5	1	45	460
1416	3	2	40	232
1534	3	2	30	315
852	2	1	36	178
...	...	...	...	...

$m = 47$

Notation:

n = number of features

$n = 4$


$x^{(i)}$ = input (features) of i^{th} training example.

$x_j^{(i)}$ = value of feature j in i^{th} training example.

$$\underline{x^{(2)}} = \begin{bmatrix} 1416 \\ 3 \\ 2 \\ 40 \end{bmatrix}$$

$$x_3^{(2)} = 2$$

Hypothesis:

Previously: $\underline{h_{\theta}(x) = \theta_0 + \theta_1 x}$ 

$$h_{\theta}(x) = \theta_0 + \theta_1 x_1 + \theta_2 x_2 + \theta_3 x_3 + \theta_4 x_4$$

e.g. $\underline{h_{\theta}(x)} = \underline{80} + \underline{0.1}x_1 + \underline{0.01}x_2 + 3x_3 - 2x_4$

$\uparrow \quad \uparrow$

$\uparrow$
age

$$\rightarrow h_{\theta}(x) = \theta_0 + \theta_1 x_1 + \theta_2 x_2 + \dots + \theta_n x_n$$

For convenience of notation, define $x_0 = 1$. $(x_0^{(i)} = 1)$

$$x = \begin{bmatrix} x_0 \\ x_1 \\ x_2 \\ \vdots \\ x_n \end{bmatrix} \in \mathbb{R}^{n+1}$$

$$\theta = \begin{bmatrix} \theta_0 \\ \theta_1 \\ \theta_2 \\ \vdots \\ \theta_n \end{bmatrix} \in \mathbb{R}^{n+1}$$

$$h_{\theta}(x) = \theta_0 x_0 + \theta_1 x_1 + \dots + \theta_n x_n$$

$\downarrow = 1$


$$= \boxed{\theta^T x}$$

$$\underbrace{[\theta_0 \ \theta_1 \ \dots \ \theta_n]}_{\theta^T} \begin{bmatrix} x_0 \\ x_1 \\ \vdots \\ x_n \end{bmatrix}$$

$(n+1) \times 1$
matrix

$\theta^T x$

Multivariate linear regression. $\leftarrow$



Machine Learning

Linear Regression with multiple variables

Gradient descent for multiple variables

Hypothesis: $h_{\theta}(x) = \theta^T x = \theta_0 x_0 + \theta_1 x_1 + \theta_2 x_2 + \dots + \theta_n x_n$

Parameters: $\theta_0, \theta_1, \dots, \theta_n$ θ $n+1$ -dimensional vector

Cost function:

$$\underbrace{J(\theta_0, \theta_1, \dots, \theta_n)}_{J(\theta)} = \frac{1}{2m} \sum_{i=1}^m (h_{\theta}(x^{(i)}) - y^{(i)})^2$$

Gradient descent:

Repeat {

$\rightarrow \theta_j := \theta_j - \alpha \frac{\partial}{\partial \theta_j} \underbrace{J(\theta_0, \dots, \theta_n)}_{J(\theta)}$

}

(simultaneously update for every $j = 0, \dots, n$)

Gradient Descent

Previously ($n=1$):

Repeat {

→ $\theta_0 := \theta_0 - \alpha \underbrace{\frac{1}{m} \sum_{i=1}^m (h_{\theta}(x^{(i)}) - y^{(i)})}_{\frac{\partial}{\partial \theta_0} J(\theta)}$

→ $\theta_1 := \theta_1 - \alpha \frac{1}{m} \sum_{i=1}^m (h_{\theta}(x^{(i)}) - y^{(i)}) \underbrace{x_1^{(i)}}_{x_1^{(i)}}$
(simultaneously update θ_0, θ_1)

}

New algorithm ($n \geq 1$):

Repeat {

→ $\theta_j := \theta_j - \alpha \frac{1}{m} \sum_{i=1}^m (h_{\theta}(x^{(i)}) - y^{(i)}) x_j^{(i)}$
(simultaneously update θ_j for $j = 0, \dots, n$)

}


→ $\theta_0 := \theta_0 - \alpha \frac{1}{m} \sum_{i=1}^m (h_{\theta}(x^{(i)}) - y^{(i)}) \underbrace{x_0^{(i)}}_{x_0^{(i)}}$

→ $\theta_1 := \theta_1 - \alpha \frac{1}{m} \sum_{i=1}^m (h_{\theta}(x^{(i)}) - y^{(i)}) \underline{x_1^{(i)}}$

→ $\theta_2 := \theta_2 - \alpha \frac{1}{m} \sum_{i=1}^m (h_{\theta}(x^{(i)}) - y^{(i)}) x_2^{(i)}$
...

$\sqrt{\frac{2}{\alpha \theta_j} J(\theta)}$

$x_0^{(i)} = 1$



Machine Learning

Linear Regression with multiple variables


Gradient descent in practice I: Feature Scaling

Feature Scaling

Idea: Make sure features are on a similar scale.

E.g. $x_1 = \text{size (0-2000 feet}^2\text{)}$ ←


$x_2 = \text{number of bedrooms (1-5)}$ ←



$$\rightarrow x_1 = \frac{\text{size (feet}^2\text{)}}{2000} \quad \checkmark$$

$$\rightarrow x_2 = \frac{\text{number of bedrooms}}{5} \quad \checkmark$$

$$0 \leq x_1 \leq 1 \quad 0 \leq x_2 \leq 1$$



Feature Scaling

Get every feature into approximately a $-1 \leq x_i \leq 1$ range.

$$x_0 = 1$$

$$0 \leq x_1 \leq 3 \quad \checkmark$$

$$-2 \leq x_2 \leq 0.5 \quad \checkmark$$

$$-100 \leq x_3 \leq 100 \quad \times$$

$$-0.0001 \leq x_4 \leq 0.0001 \quad \times$$

$$-3 \text{ to } 3 \quad \checkmark$$

$$-\frac{1}{5} \text{ to } \frac{1}{5} \quad \checkmark$$

Mean normalization

Replace x_i with $x_i - \mu_i$ to make features have approximately zero mean
(Do not apply to $x_0 = 1$).

E.g. $\rightarrow x_1 = \frac{\text{size} - 1000}{2000}$

Average size = 1000

$$x_2 = \frac{\#bedrooms - 2}{5}$$

1-5 bedrooms


$\rightarrow -0.5 \leq x_1 \leq 0.5, -0.5 \leq x_2 \leq 0.5$

$$x_1 \leftarrow \frac{x_1 - \mu_1}{\sigma_1}$$

← avg value of x_1 in training set

$$x_2 \leftarrow \frac{x_2 - \mu_2}{\sigma_2}$$

range (max-min)
(or standard deviation)



Machine Learning

Linear Regression with multiple variables


Gradient descent in practice II: Learning rate

Gradient descent

$$\rightarrow \theta_j := \theta_j - \alpha \frac{\partial}{\partial \theta_j} J(\theta)$$

- “Debugging”: How to make sure gradient descent is working correctly.
- How to choose learning rate α .


Making sure gradient descent is working correctly.



→ Example automatic convergence test:



→ Declare convergence if $J(\theta)$ decreases by less than 10^{-3} in one iteration.

Making sure gradient descent is working correctly.



Gradient descent not working.

Use smaller α .



- For sufficiently small α , $J(\theta)$ should decrease on every iteration.
- But if α is too small, gradient descent can be slow to converge.

Summary:


- If α is too small: slow convergence.
- If α is too large: $J(\theta)$ may not decrease on every iteration; may not converge. (Slow converge also possible.)



To choose α , try

..., 0.001, 0.003, 0.01, 0.03, 0.1, 0.3, 1, ...

Arrows indicate the sequence of values: $\uparrow$ (under 0.001), $\curvearrowright$ (from 0.001 to 0.003), $\curvearrowright$ (from 0.003 to 0.01), $\curvearrowright$ (from 0.01 to 0.03), $\curvearrowright$ (from 0.03 to 0.1), $\curvearrowright$ (from 0.1 to 0.3), $\uparrow$ (under 1). The labels $\approx 3\times$ are written below the arrows, indicating that each value is approximately 3 times the previous one.



Machine Learning

Linear Regression with multiple variables

Features and polynomial regression

Housing prices prediction

$$h_{\theta}(x) = \theta_0 + \theta_1 \times \underbrace{\text{frontage}}_{x_1} + \theta_2 \times \underbrace{\text{depth}}_{x_2}$$

Area


$$x = \underline{\text{frontage} \times \text{depth}}$$

$$h_{\theta}(x) = \theta_0 + \theta_1 x$$

↑ land area



Polynomial regression



$\theta_0 + \theta_1 x + \theta_2 x^2$

$\rightarrow \theta_0 + \theta_1 x + \theta_2 x^2 + \theta_3 x^3$

$$h_{\theta}(x) = \theta_0 + \theta_1 x_1 + \theta_2 x_2 + \theta_3 x_3$$
$$= \theta_0 + \theta_1(\text{size}) + \theta_2(\text{size})^2 + \theta_3(\text{size})^3$$

$\rightarrow x_1 = (\text{size})$

$\rightarrow x_2 = (\text{size})^2$


$\rightarrow x_3 = (\text{size})^3$

Size: 1-1000

Size²: 1-1,000,000

Size³: 1-10⁹


Choice of features



→ $h_{\theta}(x) = \theta_0 + \theta_1(\text{size}) + \theta_2(\text{size})^2$

→ $h_{\theta}(x) = \theta_0 + \theta_1(\text{size}) + \theta_2\sqrt{(\text{size})}$






Machine Learning

Linear Regression with multiple variables

Normal equation

Gradient Descent




Normal equation: Method to solve for θ analytically.

Intuition: If 1D ($\theta \in \mathbb{R}$)

$\rightarrow \underline{J(\theta) = a\theta^2 + b\theta + c}$

$\frac{d}{d\theta} J(\theta) = \dots \stackrel{\text{set}}{=} 0$

Solve for θ



$$\underline{\theta \in \mathbb{R}^{n+1}} \quad \underline{J(\theta_0, \theta_1, \dots, \theta_m) = \frac{1}{2m} \sum_{i=1}^m (h_{\theta}(x^{(i)}) - y^{(i)})^2}$$

$$\underline{\frac{\partial}{\partial \theta_j} J(\theta) = \dots \stackrel{\text{set}}{=} 0} \quad (\text{for every } j)$$

Solve for $\underline{\theta_0, \theta_1, \dots, \theta_n}$

Examples: $m = 4$.

	Size (feet ²)	Number of bedrooms	Number of floors	Age of home (years)	Price (\$1000)
x_0	x_1	x_2	x_3	x_4	y
1	2104	5	1	45	460
1	1416	3	2	40	232
1	1534	3	2	30	315
1	852	2	1	36	178

$\underline{X} = \begin{bmatrix} 1 & 2104 & 5 & 1 & 45 \\ 1 & 1416 & 3 & 2 & 40 \\ 1 & 1534 & 3 & 2 & 30 \\ 1 & 852 & 2 & 1 & 36 \end{bmatrix}$
 $m \times (n+1)$

$\underline{y} = \begin{bmatrix} 460 \\ 232 \\ 315 \\ 178 \end{bmatrix}$
 m -dimensional vector

$$\theta = (X^T X)^{-1} X^T y$$

m examples $(x^{(1)}, y^{(1)}), \dots, (x^{(m)}, y^{(m)})$; n features.

$$\underline{x^{(i)}} = \begin{bmatrix} x_0^{(i)} \\ x_1^{(i)} \\ x_2^{(i)} \\ \vdots \\ x_n^{(i)} \end{bmatrix} \in \mathbb{R}^{n+1} \quad \bigg| \quad \begin{matrix} X \\ \text{(design} \\ \text{matrix)} \end{matrix} = \begin{bmatrix} \text{---} (x^{(1)})^T \text{---} \\ \text{---} (x^{(2)})^T \text{---} \\ \vdots \\ \text{---} (x^{(m)})^T \text{---} \end{bmatrix}$$

E.g. If $\underline{x^{(i)}} = \begin{bmatrix} 1 \\ x_1^{(i)} \end{bmatrix}$ $\rightarrow$ $X = \begin{bmatrix} | & x^{(1)} \\ | & x^{(2)} \\ \vdots & \vdots \\ | & x^{(m)} \end{bmatrix}$ $m \times (n+1)$

$\Theta = (X^T X)^{-1} X^T y$

$y = \begin{bmatrix} y^{(1)} \\ y^{(2)} \\ \vdots \\ y^{(m)} \end{bmatrix}$ $m \times 2$

$$\theta = \underbrace{(X^T X)^{-1} X^T y}$$

$(X^T X)^{-1}$ is inverse of matrix $X^T X$.

Set $\underbrace{A = X^T X}$

$$\boxed{(X^T X)^{-1}} = A^{-1}$$

Octave: `pinv(X' * X) * X' * y`

$$\underbrace{\text{pinv}(X^T * X) * X^T * y}$$

$$\theta = (X^T X)^{-1} X^T y$$

$\min_{\theta} J(\theta)$

X'

X^T

~~Feature Scaling~~

$$0 \leq x_1 \leq 1$$

$$0 \leq x_2 \leq 1000$$

$$0 \leq x_3 \leq 10^{-5} \quad \checkmark$$

m training examples, n features.

Gradient Descent

- • Need to choose α .
- • Needs many iterations.
- Works well even when n is large.

↗
 $n \approx 10^6$


← -

Normal Equation

- • No need to choose α .
- • Don't need to iterate.
- Need to compute
- • $(X^T X)^{-1}$ $\frac{n \times n}{\quad}$ $O(n^3)$
- Slow if n is very large.

$n \approx 100$
 $n \approx 1000$

- - - $n \approx 10000$



Machine Learning

Linear Regression with multiple variables

Normal equation and non-invertibility (optional)


Normal equation

$$\theta = \underline{(X^T X)^{-1} X^T y}$$

$$\underline{X^T X}$$

- What if $\boxed{X^T X}$ is non-invertible? (singular/degenerate)

- Octave: `pinv(X' * X) * X' * y`



What if $X^T X$ is non-invertible?

- Redundant features (linearly dependent).

E.g. $\begin{cases} x_1 = \text{size in feet}^2 \\ \cancel{x_2 = \text{size in m}^2} \\ x_1 = (3.28)^2 x_2 \end{cases}$

$1\text{m} = 3.28\text{ feet}$

$\rightarrow m = 10 \leftarrow$

$\rightarrow n = 100 \leftarrow$

$\odot \in \mathbb{R}^{101}$

- Too many features (e.g. $m \leq n$).

- Delete some features, or use regularization.


$\downarrow$ later

Referências

NG, Andrew. *Machine Learning*. Stanford University, 2011. Curso oferecido via Coursera. Disponível em: <https://www.coursera.org/learn/machine-learning>.



Dúvidas?



Até a próxima...



Apresentador

Thales Levi Azevedo Valente

E-mail:

thales.l.a.valente@gmail.com