

Applied Linear Algebra

by
Dr.Dinesh Naik
(B.E, M.Tech, Ph.D)

Assistant Professor, Dept. of Information Technology
National Institute of Technology Karnataka, Surathkal

March 15, 2023

Acknowledgement

- I would like to express my sincere gratitude Stephen Boyd, Department of Electrical Engineering Stanford University
- I also thank Lieven Vandenberghe Department of Electrical and Computer Engineering University of California, Los Angeles

Linear Algebra - study of systems of linear equations

A **linear equation** is of the form

$$a_1x_1 + a_2x_2 + \cdots + a_nx_n = b \quad \text{where}$$

- ▶ x_1, x_2, \dots, x_n are variables
- ▶ a_1, a_2, \dots, a_n are called coefficients
- ▶ b is the right-hand-side value
- ▶ a_1, a_2, \dots, a_n and b are typically real numbers, i.e., numbers found on a number line

Linear Algebra - study of systems of linear equations

A **linear equation** is of the form

$$a_1x_1 + a_2x_2 + \cdots + a_nx_n = b \quad \text{where}$$

- ▶ x_1, x_2, \dots, x_n are variables
- ▶ a_1, a_2, \dots, a_n are called coefficients
- ▶ b is the right-hand-side value
- ▶ a_1, a_2, \dots, a_n and b are typically real numbers, i.e., numbers found on a number line

Terms like x_1^2 or x_1x_2 are nonlinear and therefore not allowed.

Some sample linear equations: $5x_1 + 3x_2 = 10$

$$6x_1 + 8x_2 - 9x_3 = 14$$

System of linear equations—multiple equations.

A system of linear equations has either:

1. No solution.
2. Exactly one solution (unique solution).
3. Infinitely many solutions.

A system of linear equations has either:

1. No solution.
2. Exactly one solution (unique solution).
3. Infinitely many solutions.

A system is said to be consistent if it has at least one solution; otherwise it is inconsistent.

Need a systematic method to solve a system of equations.

Elementary row operations:

1. Interchange (swap) two rows.
2. Multiply all entries in a row by a nonzero constant.
3. Replace one row by the sum of itself and another row.

Two matrices are row equivalent if there is a sequence of elementary row operations that transforms one matrix into the other.

Need a systematic method to solve a system of equations.

Elementary row operations:

1. Interchange (swap) two rows.
2. Multiply all entries in a row by a nonzero constant.
3. Replace one row by the sum of itself and another row.

Two matrices are row equivalent if there is a sequence of elementary row operations that transforms one matrix into the other.

If the augmented matrices of two systems are row equivalent, then the two systems have the same solution set.

Two Fundamental Questions:

1. Is the system consistent, i.e., does at least one solution exist?
2. If the system is consistent, then is there a unique solution or an infinite number of solutions?

Inner product

- ▶ *inner product* (or *dot product*) of n -vectors a and b is

$$a^T b = a_1 b_1 + a_2 b_2 + \cdots + a_n b_n$$

- ▶ other notation used: $\langle a, b \rangle$, $\langle a|b \rangle$, (a, b) , $a \cdot b$
- ▶ example:

$$\begin{bmatrix} -1 \\ 2 \\ 2 \end{bmatrix}^T \begin{bmatrix} 1 \\ 0 \\ -3 \end{bmatrix} = (-1)(1) + (2)(0) + (2)(-3) = -7$$

Inner product. When a and b are n -vectors, $a^T b$ is exactly the inner product of a and b , obtained from the rules for transposing matrices and forming a matrix-vector product. We start with the (column) n -vector a , consider it as an $n \times 1$ matrix, and transpose it to obtain the n -row-vector a^T . Now we multiply this $1 \times n$ matrix by the n -vector b , to obtain the 1-vector $a^T b$, which we also consider a scalar. So the notation $a^T b$ for the inner product is just a special case of matrix-vector multiplication.

Properties of inner product

- ▶ $a^T b = b^T a$
- ▶ $(\gamma a)^T b = \gamma(a^T b)$
- ▶ $(a + b)^T c = a^T c + b^T c$

can combine these to get, for example,

$$(a + b)^T (c + d) = a^T c + a^T d + b^T c + b^T d$$

General examples

- ▶ $e_i^T a = a_i$ (picks out i th entry)
- ▶ $\mathbf{1}^T a = a_1 + \cdots + a_n$ (sum of entries)
- ▶ $a^T a = a_1^2 + \cdots + a_n^2$ (sum of squares of entries)

Examples

- ▶ w is weight vector, f is feature vector; $w^T f$ is score
- ▶ p is vector of prices, q is vector of quantities; $p^T q$ is total cost
- ▶ c is cash flow, d is discount vector (with interest rate r):

$$d = (1, 1/(1+r), \dots, 1/(1+r)^{n-1})$$

$d^T c$ is net present value (NPV) of cash flow

- ▶ s gives portfolio holdings (in shares), p gives asset prices; $p^T s$ is total portfolio value

Block vectors. If the vectors a and b are block vectors, and the corresponding blocks have the same sizes (in which case we say they *conform*), then we have

$$a^T b = \begin{bmatrix} a_1 \\ \vdots \\ a_k \end{bmatrix}^T \begin{bmatrix} b_1 \\ \vdots \\ b_k \end{bmatrix} = a_1^T b_1 + \cdots + a_k^T b_k.$$

The inner product of block vectors is the sum of the inner products of the blocks.

Applications. The inner product is useful in many applications, a few of which we list here.

- *Co-occurrence.* If a and b are n -vectors that describe occurrence, *i.e.*, each of their elements is either 0 or 1, then $a^T b$ gives the total number of indices for which a_i and b_i are both one, that is, the total number of co-occurrences. If we interpret the vectors a and b as describing subsets of n objects, then $a^T b$ gives the number of objects in the intersection of the two subsets. This is illustrated in figure 1.13, for two subsets A and B of 7 objects, labeled $1, \dots, 7$, with corresponding occurrence vectors

$$a = (0, 1, 1, 1, 1, 1, 1), \quad b = (1, 0, 1, 0, 1, 0, 0).$$

Here we have $a^T b = 2$, which is the number of objects in both A and B (*i.e.*, objects 3 and 5).




Figure 1 Two sets A and B , containing seven objects.