


Divide-and-Conquer Strategy




Divide-and-Conquer Strategy

◆ Divide-and conquer is a general algorithm design paradigm:

- **Divide**: Divide the input data S in two or more disjoint subsets S_1, S_2, \dots
- **Recur**: Solve each of these subproblems recursively
- **Conquer**: Combine the solutions for S_1, S_2, \dots , into a solution for S

◆ Base-cases for recursion are subproblems of constant size

◆ Analysis can be done using **Recurrence Equations**



Merge-Sort Review

- ◆ Merge-sort on a given input sequence S with n elements consists of three steps:
 - **Divide**: Partition S into two sequences S_1 and S_2 of about $n/2$ elements each
 - **Recur**: Sort S_1 and S_2 Recursively
 - **Conquer**: Merge S_1 and S_2 into a unique sorted sequence

Algorithm *mergeSort*(S, C)

Input sequence S with n elements, comparator C

Output sequence S sorted according to C

if $S.size() > 1$

$(S_1, S_2) \leftarrow partition(S, n/2)$

mergeSort(S_1, C)

mergeSort(S_2, C)

$S \leftarrow merge(S_1, S_2)$


Recurrence Equations



- ◆ The conquer step of merge-sort consists of merging two sorted sequences, each with $n/2$ elements and implemented by means of a doubly linked list, takes at most bn steps, for some constant b .
- ◆ Likewise, the basis case ($n < 2$) will take at b most steps.
- ◆ Therefore, if we let $T(n)$ denote the running time of merge-sort:

$$T(n) = \begin{cases} b & \text{if } n < 2 \\ 2T(n/2) + bn & \text{if } n \geq 2 \end{cases}$$

- ◆ We can therefore analyze the running time of merge-sort by finding a **closed form solution** to the above equation.
 - That is, a solution that has $T(n)$ only on the left-hand side.



Iterative Substitution

- ◆ In the iterative substitution, or “plug-and-chug,” technique, we iteratively apply the recurrence equation to itself and see if we can find a pattern:


$$\begin{aligned}T(n) &= 2T(n/2) + bn \\&= 2(2T(n/2^2)) + b(n/2) + bn \\&= 2^2T(n/2^2) + 2bn \\&= 2^3T(n/2^3) + 3bn \\&= 2^4T(n/2^4) + 4bn \\&= \dots \\&= 2^iT(n/2^i) + ibn\end{aligned}$$

- ◆ Note that base, $T(n)=b$, case occurs when $2^i=n$. That is, $i = \log n$.
- ◆ So, $T(n) = bn + bn \log n$; Thus, $T(n)$ is $O(n \log n)$.



- $$T(n) = \begin{cases} b & \text{if } n < 2 \\ 2T(n/2) + bn & \text{if } n \geq 2 \end{cases}$$

Total Time = $bn + bn \log n$
(Last Level plus All Previous Levels)



Guess-and-Test Method


- ◆ In the guess-and-test method, we guess a closed form solution and then try to prove that it is true by induction:

$$T(n) = \begin{cases} b & \text{if } n < 2 \\ 2T(n/2) + bn \log n & \text{if } n \geq 2 \end{cases}$$

- ◆ **Guess #1: $T(n) < cn \log n$.**

$$\begin{aligned} T(n) &= 2T(n/2) + bn \log n \\ &= 2(c(n/2) \log(n/2)) + bn \log n \\ &= cn(\log n - \log 2) + bn \log n \\ &= cn \log n - cn + bn \log n \end{aligned}$$

- ◆ **Wrong: We cannot make this last line be less than $cn \log n$**



Guess-and-Test Method

- ◆ Recall the Recurrence Equation:


$$T(n) = \begin{cases} b & \text{if } n < 2 \\ 2T(n/2) + bn \log n & \text{if } n \geq 2 \end{cases}$$

- ◆ **Guess #2: $T(n) < cn \log^2 n$.**


$$\begin{aligned} T(n) &= 2T(n/2) + bn \log n \\ &= 2(c(n/2) \log^2(n/2)) + bn \log n \\ &= cn(\log n - \log 2)^2 + bn \log n \\ &= cn \log^2 n - 2cn \log n + cn + bn \log n \\ &\leq cn \log^2 n \end{aligned}$$

- ◆ **If $c > b$, then $T(n)$ is $O(n \log^2 n)$.**
- ◆ **In general, to use this method, we need to have a good guess and further we need to be good at induction proofs.**


Master Method (Appendix)



Master Method, Example 1



Master Method, Example 2



◆ The form:
$$T(n) = \begin{cases} c & \text{if } n < d \\ aT(n/b) + f(n) & \text{if } n \geq d \end{cases}$$

◆ The Master Theorem:

Case-1. if $f(n)$ is $O(n^{\log_b a - \epsilon})$, then $T(n)$ is $\Theta(n^{\log_b a})$

Case-2. if $f(n)$ is $\Theta(n^{\log_b a} \log^k n)$, then $T(n)$ is $\Theta(n^{\log_b a} \log^{k+1} n)$


Case-3. if $f(n)$ is $\Omega(n^{\log_b a + \epsilon})$, then $T(n)$ is $\Theta(f(n))$,
provided $af(n/b) \leq \delta f(n)$ for some $\delta < 1$.

◆ Example:

$$T(n) = 2T(n/2) + n \log n$$

Solution: $\log_b a = 1$, so Case-2 says $T(n)$ is $O(n \log^2 n)$.

Master Method, Example 3



◆ The form:
$$T(n) = \begin{cases} c & \text{if } n < d \\ aT(n/b) + f(n) & \text{if } n \geq d \end{cases}$$

◆ The Master Theorem:

Case-1. if $f(n)$ is $O(n^{\log_b a - \epsilon})$, then $T(n)$ is $\Theta(n^{\log_b a})$

Case-2. if $f(n)$ is $\Theta(n^{\log_b a} \log^k n)$, then $T(n)$ is $\Theta(n^{\log_b a} \log^{k+1} n)$


Case-3. if $f(n)$ is $\Omega(n^{\log_b a + \epsilon})$, then $T(n)$ is $\Theta(f(n))$,
provided $af(n/b) \leq \delta f(n)$ for some $\delta < 1$.

◆ Example:

$$T(n) = T(n/3) + n \log n$$

Solution: $\log_b a = 0$, so Case-3 says $T(n)$ is $O(n \log n)$.

Master Method, Example 4



◆ The form:
$$T(n) = \begin{cases} c & \text{if } n < d \\ aT(n/b) + f(n) & \text{if } n \geq d \end{cases}$$

◆ The Master Theorem:

Case-1. if $f(n)$ is $O(n^{\log_b a - \epsilon})$, then $T(n)$ is $\Theta(n^{\log_b a})$

Case-2. if $f(n)$ is $\Theta(n^{\log_b a} \log^k n)$, then $T(n)$ is $\Theta(n^{\log_b a} \log^{k+1} n)$


Case-3. if $f(n)$ is $\Omega(n^{\log_b a + \epsilon})$, then $T(n)$ is $\Theta(f(n))$,
provided $af(n/b) \leq \delta f(n)$ for some $\delta < 1$.

◆ Example:

$$T(n) = 8T(n/2) + n^2$$

Solution: $\log_b a = 3$, so Case-1 says $T(n)$ is $O(n^3)$.

Master Method, Example 5



◆ The form:
$$T(n) = \begin{cases} c & \text{if } n < d \\ aT(n/b) + f(n) & \text{if } n \geq d \end{cases}$$

◆ The Master Theorem:

Case-1. if $f(n)$ is $O(n^{\log_b a - \epsilon})$, then $T(n)$ is $\Theta(n^{\log_b a})$

Case-2. if $f(n)$ is $\Theta(n^{\log_b a} \log^k n)$, then $T(n)$ is $\Theta(n^{\log_b a} \log^{k+1} n)$


Case-3. if $f(n)$ is $\Omega(n^{\log_b a + \epsilon})$, then $T(n)$ is $\Theta(f(n))$,
provided $af(n/b) \leq \delta f(n)$ for some $\delta < 1$.

◆ Example:

$$T(n) = 9T(n/3) + n^3$$

Solution: $\log_b a = 2$, so Case-3 says $T(n)$ is $O(n^3)$.

Master Method, Example 6



◆ The form:
$$T(n) = \begin{cases} c & \text{if } n < d \\ aT(n/b) + f(n) & \text{if } n \geq d \end{cases}$$

◆ The Master Theorem:

Case-1. if $f(n)$ is $O(n^{\log_b a - \epsilon})$, then $T(n)$ is $\Theta(n^{\log_b a})$

Case-2. if $f(n)$ is $\Theta(n^{\log_b a} \log^k n)$, then $T(n)$ is $\Theta(n^{\log_b a} \log^{k+1} n)$


Case-3. if $f(n)$ is $\Omega(n^{\log_b a + \epsilon})$, then $T(n)$ is $\Theta(f(n))$,
provided $af(n/b) \leq \delta f(n)$ for some $\delta < 1$.

◆ Example:

$$T(n) = T(n/2) + 1 \quad (\text{Binary Search})$$

Solution: $\log_b a = 0$, so Case-2 says $T(n)$ is $O(\log n)$.

Master Method, Example 7



◆ The form:
$$T(n) = \begin{cases} c & \text{if } n < d \\ aT(n/b) + f(n) & \text{if } n \geq d \end{cases}$$

◆ The Master Theorem:

Case-1. if $f(n)$ is $O(n^{\log_b a - \epsilon})$, then $T(n)$ is $\Theta(n^{\log_b a})$


Case-2. if $f(n)$ is $\Theta(n^{\log_b a} \log^k n)$, then $T(n)$ is $\Theta(n^{\log_b a} \log^{k+1} n)$

Case-3. if $f(n)$ is $\Omega(n^{\log_b a + \epsilon})$, then $T(n)$ is $\Theta(f(n))$,
provided $af(n/b) \leq \delta f(n)$ for some $\delta < 1$.

◆ Example:

$$T(n) = 2T(n/2) + \log n \quad (\text{Heap Construction})$$

Solution: $\log_b a = 1$, so Case-1 says $T(n)$ is $O(n)$.




Master Theorem Proof

- ◆ Using iterative substitution, let us see if we can find a pattern:

$$\begin{aligned}T(n) &= aT(n/b) + f(n) \\&= a(aT(n/b^2)) + f(n/b) + bn \\&= a^2T(n/b^2) + af(n/b) + f(n) \\&= a^3T(n/b^3) + a^2f(n/b^2) + af(n/b) + f(n) \\&= \dots \\&= a^{\log_b n}T(1) + \sum_{i=0}^{(\log_b n)-1} a^i f(n/b^i) \\&= n^{\log_b a}T(1) + \sum_{i=0}^{(\log_b n)-1} a^i f(n/b^i)\end{aligned}$$

- ◆ We then distinguish the three cases of Master theorem as
 - The first term is dominant
 - Each part of the summation is equally dominant
 - The summation is a geometric series



Integer Multiplication

◆ Algorithm: Multiply two n-bit integers I and J.

- Divide step: Split I and J into high-order and low-order bits


$$I = I_h 2^{n/2} + I_l$$

$$J = J_h 2^{n/2} + J_l$$

- We can then define $I * J$ by multiplying the parts and adding:

$$\begin{aligned} I * J &= (I_h 2^{n/2} + I_l) * (J_h 2^{n/2} + J_l) \\ &= I_h J_h 2^n + I_h J_l 2^{n/2} + I_l J_h 2^{n/2} + I_l J_l \end{aligned}$$

- So, $T(n) = 4T(n/2) + n$, which implies $T(n)$ is $O(n^2)$.
- But that is no better than the algorithm we already learnt.



Integer Multiplication

◆ Algorithm: Multiply two n-bit integers I and J.

- Divide step: Split I and J into high-order and low-order bits

$$I = I_h 2^{n/2} + I_l$$

$$J = J_h 2^{n/2} + J_l$$

- Observe that there is a different way to multiply parts:

$$\begin{aligned} I * J &= I_h J_h 2^n + [(I_h - I_l)(J_l - J_h) + I_h J_h + I_l J_l] 2^{n/2} + I_l J_l \\ &= I_h J_h 2^n + [(I_h J_l - I_l J_l - I_h J_h + I_l J_h) + I_h J_h + I_l J_l] 2^{n/2} + I_l J_l \\ &= I_h J_h 2^n + (I_h J_l + I_l J_h) 2^{n/2} + I_l J_l \end{aligned}$$

- $T(n) = 3T(n/2) + n$, it implies $T(n)$ is $O(n^{\log_2 3})$ (By Master Theorem)
- Thus, $T(n)$ is $O(n^{1.585})$.