


Label Size:-72 x 18mm

Composition
Gatifloxacin Sesquihydrate
Eq.to Gatifloxacin 0.5%w/v
Benzalkonium Chloride
Solution IP 0.02%v/v
(As Preservative)
Sterile Aqueous base q.s
Dosage: As directed by
the Physician.
Keep the medicine out of
reach of children.
For Direction, warning,
storage & instructions for
use, Please see the Carton.

SCHEDULE H PRESCRIPTION DRUG -
CAUTION Not to be sold by retail
without the prescription of a Registered
Medical Practitioner.



Marketed by:



Lifesciences Pvt Ltd

14L Floor 6th Navjeevan Commercial

Premises Mumbai Central

Mumbai-400008, Maharashtra India.

Manufactured in India by:

ABLE PHARMA.

Village Gullarwala, Sai Road,

Baddi, Distt.-Solani-173 205 (H.P.)

(WHO-GMP Certified Unit)

Mfg. Lic. No.: MB/06490 & MNB/06499

Batch No.:

बैच नं.:

Mfg. Date:

उत्पादन तिथि

Expiry Date:

समाप्ति की तिथि

M.R.P. ₹.:

(Inclusive of all taxes)



Label Size:-72 x 18mm

Gativix
Eye Drops



Marketed by:



Lifesciences Pvt Ltd

14L Floor 6th Navjeevan Commercial

Mumbai-400008, Maharashtra India.

Manufactured in India by:

ABLE PHARMA.

Village Gullarwala, Sai Road,

Baddi, Distt.-Solani-173 205 (H.P.)

(WHO-GMP Certified Unit)

Mfg. Lic. No.: MB/06490 & MNB/06499

Batch No.:

बैच नं.:

Mfg. Date:


उत्पादन तिथि

Expiry Date:

समाप्ति की तिथि

M.R.P. ₹.:

(Inclusive of all taxes)



Carton size : 45 x 32 x 80 mm

Gatifloxacin
Ophthalmic
Solution
Gativix
Eye Drops



STERILE



Composition
Gatifloxacin Sesquihydrate
Eq.to Gatifloxacin 0.5%w/v
Benzalkonium Chloride
Solution IP 0.02%v/v
(As Preservative)
Sterile Aqueous base q.s
Dosage: As directed by the
Physician.
Storage: Store in a cool &
dry place at a temperature
not exceeding 30°C. Protect
from light. Do not allow to
freeze.
Warning: 1.If irritation
persists or increases,
discontinue the use and
consult the Physician.
2.Do not touch the nozzle
tip to any surface. This may
contaminate the solution.
3.Do not use the vial if it
contains visible particulate
matter.
**Keep the medicine out of
reach of children.**

SCHEDULE H PRESCRIPTION DRUG -
CAUTION Not to be sold by retail
without the prescription of a Registered
Medical Practitioner.

Gatifloxacin
Ophthalmic
Solution
Gativix
Eye Drops



STERILE



- INSTRUCTIONS FOR USE**
- Remove the last ring by pulling the notch.
 - PULL**
 - SPIKE**
 - Tighten the cap on the nozzle as shown.
 - The spike in the cap will pierce the tip of the vial.
 - Dispense drops with gentle pressure.
 - Replace the cap after each use.

FOR EXTERNAL USE ONLY

NOT FOR INJECTION

Direction:- Use the solution within one month after opening the vial.

Mfg. Lic. No.: MB/06490 & MNB/06499
Manufactured in India by:

ABLE PHARMA

Village Gullarwala, Sai Road,
Baddi, Distt.-Solani-173 205 (H.P.)
(WHO-GMP Certified Unit)

Marketed By:



Lifesciences Pvt Ltd

14L Floor 6th Navjeevan Commercial
Premises Mumbai Central
Mumbai-400008, Maharashtra India.

Batch No. :

बैच नं.:

Mfg. Date :

उत्पादन तिथि

Expiry Date:

समाप्ति तिथि

M.R.P. ₹.:

(Inclusive of all taxes)

अधिकतम खुदरा मूल्य रु.
(सभी करों सहित)



Insert size:- 148 x 125 mm

For the use only of a Registered Medical Practitioner or a Hospital or a Laboratory

Gatifloxacin Ophthalmic Solution

Eye Drops

Gativix
Eye Drops

5ml
STERILE

Composition:-

Gatifloxacin Sesquihydrate
Eq. to Gatifloxacin 0.5%w/v
Benzalkonium Chloride Solution IP 0.02%w/v
(As Preservative)
Sterile Aqueous base q.s

DESCRIPTION:

Gatifloxacin Ophthalmic Solution is a sterile, clear, aqueous ophthalmic preparation containing Gatifloxacin 0.5% w/v. Gatifloxacin belongs to the fluoroquinolone class of antibacterial medicines and is intended for topical administration to the eye. The formulation is designed to provide antibacterial activity at the site of ocular infection. Benzalkonium Chloride Solution IP is incorporated at 0.02% w/v as a preservative, while sterile aqueous base is used as the vehicle. Gatifloxacin acts against susceptible bacteria by inhibiting bacterial enzymes required for DNA replication and cell division. This ultimately interferes with bacterial growth and survival.

PHARMACOLOGICAL CLASSIFICATION:

Therapeutic Class: Ophthalmic Antibacterial
Pharmacological Class: Fluoroquinolone Antibacterial
Gatifloxacin is a fourth-generation fluoroquinolone antibacterial agent. Its antibacterial activity is primarily associated with inhibition of two essential bacterial enzymes, DNA gyrase and topoisomerase IV. These enzymes are involved in the replication, transcription, repair and recombination of bacterial DNA. By interfering with these enzymes, Gatifloxacin prevents normal bacterial DNA processing and replication, resulting in inhibition of bacterial growth and death of susceptible organisms. Following topical ocular administration, Gatifloxacin acts primarily at the site of application. The extent of systemic exposure is considerably lower than that generally associated with systemic administration.

INDICATIONS:

Gatifloxacin Ophthalmic Solution is indicated for the topical treatment of bacterial infections of the eye caused by susceptible microorganisms. It may be used in conditions such as bacterial conjunctivitis and other superficial ocular bacterial infections when Gatifloxacin-sensitive organisms are suspected or identified. The medicine should be used only when a bacterial infection is suspected or confirmed. It is not indicated for the treatment of viral, fungal or non-infectious ocular conditions. If there is no clinical improvement, or if symptoms become worse during treatment, the patient should consult a physician for further evaluation.

DOSAGE AND ADMINISTRATION:

The dosage should be determined by the physician according to the severity and nature of the infection. Adults and Children: Instill one drop into the affected eye(s) as directed by the physician. The recommended frequency and duration of treatment should be followed as prescribed. Treatment should not be discontinued prematurely unless advised by a healthcare professional.

Method of Administration:

1. Wash hands thoroughly with soap and water before administration.
 2. Remove the cap carefully.
 3. Avoid touching the dropper tip with fingers or allowing it to come into contact with the eye, eyelids or any other surface.
 4. Tilt the head backwards and gently pull down the lower eyelid to form a small pocket.
 5. Instill the prescribed number of drops into the affected eye.
 6. Close the eye gently after administration and avoid excessive blinking.
 7. Replace the cap immediately after use.
- If more than one ophthalmic medicine is being used, an interval of approximately 5–10 minutes should be maintained between different eye preparations. Ophthalmic ointments should generally be applied after administration of eye drops.

CONTRAINDICATIONS:

Gatifloxacin Ophthalmic Solution is contraindicated in patients with:
- Known hypersensitivity to Gatifloxacin, other quinolone antibiotics, or any component of the formulation.
- A previous history of serious allergic or hypersensitivity reaction associated with quinolone therapy.
Patients should inform their physician about any previous allergy to antibiotics or ophthalmic medicines before starting treatment.

WARNINGS AND PRECAUTIONS:

For topical ophthalmic use only:

Do not swallow or inject this medicine. Avoid contamination of the dropper tip. Do not touch the tip to the eye, eyelids, fingers or other surfaces. If an allergic reaction develops, including eye swelling, severe itching, rash, difficulty in breathing or other signs of hypersensitivity, discontinue treatment and seek immediate medical attention. Prolonged use of topical antibacterial medicines may result in overgrowth of non-susceptible microorganisms, including fungi. If superinfection occurs, appropriate treatment should be considered. If symptoms worsen, persist or fail to improve within the expected period, medical evaluation is recommended. Temporary blurred vision, mild irritation, burning or discomfort may occur following administration. Patients should avoid driving or operating machinery until vision has returned to normal. Benzalkonium Chloride may cause ocular irritation and may be absorbed by soft contact lenses. Contact lenses should generally be removed before administration and may be reinserted after an appropriate interval as advised by the physician. Use with caution in patients with a known history of hypersensitivity to quinolone antibacterial agents. Do not use the solution if the seal is broken, the solution appears contaminated or its appearance has changed. Keep the container tightly closed when not in use.

USE IN PREGNANCY AND LACTATION:

Pregnancy:

There are limited data regarding the use of topical ophthalmic Gatifloxacin during pregnancy. Gatifloxacin Ophthalmic Solution should be used during pregnancy only when clearly necessary and when, in the judgement of the physician, the expected therapeutic benefit outweighs any potential risk. Pregnant women should consult their physician before using this medicine.

Lactation:

It is not known to what extent Gatifloxacin administered topically to the eye is excreted into human milk. Therefore, caution should be exercised when Gatifloxacin Ophthalmic Solution is administered to breastfeeding women. A decision regarding treatment during lactation should be made by the physician after considering the importance of treatment to the mother and any potential risk to the infant.

OVERDOSAGE:

Overdosage following topical ophthalmic administration is unlikely to result in significant systemic toxicity. If excessive quantities of the solution are accidentally instilled into the eye, the eye may be gently rinsed with clean, lukewarm water. If the medicine is accidentally swallowed or if unusual or severe symptoms occur, medical advice should be sought promptly. Treatment of overdose is generally symptomatic and supportive. There is no specific antidote for Gatifloxacin overdose following ophthalmic administration.

DRUG INTERACTIONS:

Clinically significant drug interactions are unlikely following topical ophthalmic administration because systemic exposure to Gatifloxacin is expected to be low. Nevertheless, patients should inform their physician about all medicines they are currently using, including prescription medicines, non-prescription medicines and other ophthalmic preparations. When multiple topical ophthalmic medicines are prescribed, they should be administered separately with an interval of approximately 5–10 minutes between applications to reduce the possibility of washout or dilution. If an ophthalmic ointment and eye drops are being used together, the eye drops should generally be administered first, followed by the ointment. Patients should not use additional antibiotic or steroid-containing eye preparations without medical advice.

Storage: Store in a cool & dry place at a temperature not exceeding 30°C. Protect from light. Do not allow to freeze.

Keep the medicine out of reach of children.

FOR EXTERNAL USE ONLY. NOT FOR INJECTION.

Pack Presentation: Gativix Eye Drops is available in LDPE vial.

Manufactured in India by:
ABLE PHARMA
Village Gullarwala, Sai Road,
Baddi, Distt.-Solani-173 205 (H.P.)
(WHO-GMP Certified Unit)

Marketed by:-



Rivixa

Lifesciences Pvt Ltd

141 Floor 6th Navjeevan Commercial Premises
Mumbai Central Mumbai-400008, Maharashtra India