







H10 · Goals are Legendre pushes · exploratory round 1 (2026-10-03)


Claim: $P_{\text{assigned}}(\text{state}) \propto P_{\text{free}}(\text{state}) \cdot \exp(\lambda \cdot \Sigma y)$, with y = statement alignment with the assigned goal $\hat{g}$. One λ per pair is fitted to the mean push;

everything else is predicted: who moves (P1), the variance (P2), the loop gain (P3) and the 32-d response direction (P4).

Combined verdicts: P1 failed (Stouffer $p = 0.890$) · P2 n/a (no pair in the perturbative range) · P3 descriptive (2/3 CIs contain 0) · P4 descriptive (0/3 pass the original rule) · overall FAILED



P5 kickoffs: jump vs pre-period fluctuation along $\hat{g}_{\text{new}}$; $\Delta m > 0$ in 100% of 19 transitions into assigned goals; Spearman (ρ) = -0.56



Synthetic validation (60 reps/cell)
P2 testable for $\epsilon \leq 0.34$

H	$\lambda=10$	P1 0.52	P4 0.00	H10/ 0.17
H	$\lambda=20$	P1 0.63	P4 0.00	H10/ 0.15
H	$\lambda=60$	P1 0.95	P4 0.00	H10/ 0.00
H	$\lambda=150$	P1 0.92	P4 0.00	H10/ 0.00
Haniso	$\lambda=20$	P1 0.72	P4 0.00	H10/ 0.13
Hgibbs	$\lambda=2$	P1 0.27	P4 0.00	H10/ 0.08
Hgibbs	$\lambda=6$	P1 0.60	P4 0.00	H10/ 0.00
Hgibbs	$\lambda=15$	P1 0.82	P4 0.00	H10/ 0.00
R1	$\lambda=0.1$	P1 0.10	P4 0.00	H10/ 0.00
R3	$\lambda=20$	P1 0.65	P4 0.00	H10/ 0.10
R4	$\lambda=20$	P1 0.52	P4 0.00	H10/ 0.07
R5up	$\lambda=20$	P1 0.57	P4 0.00	H10/ 0.08
R6	$\lambda=20$	P1 0.30	P4 0.00	H10/ 0.05

Per-period descriptives (G folders)

G11 free: $y = -0.51$, unimodal False, $g = 0.55$ [0.21, 0.64], $g_{\perp} = 0.53$; slope CI [-0.03, -0.006]
G16 free: $y = -0.30$, unimodal False, $g = 0.51$ [0.03, 0.69], $g_{\perp} = 0.42$; slope CI [-0.016, 0.016]
G37 free: $y = -0.11$, unimodal True, $g = 0.62$ [0.34, 0.7], $g_{\perp} = 0.56$; slope CI [-0.008, 0.024]
G31:g12 free: $y = +0.31$, unimodal True, $g = 0.71$ [0.59, 0.79], $g_{\perp} = 0.74$; slope CI [-0.008, 0.023]
G31:g17 free: $y = +0.62$, unimodal False, $g = 0.55$ [0.05, 0.74], $g_{\perp} = 0.74$; slope CI [-0.029, 0.013]
G12 assigned: all A days above free mean True; day1 - rest = +0.159; $g_A = 0.79$; trend slope CI [-0.05, -0.012]
G17 assigned: all A days above free mean True; day1 - rest = +0.019; $g_A = 0.70$; trend slope CI [-0.018, 0.045]
G38 assigned: all A days above free mean True; day1 - rest = -0.036; $g_A = 0.58$; trend slope CI [0.01, 0.03]