STA130 Fall 2019 – T0107

Week 2: Data Wrangling and R help

(Materials used in this presentation are provided by the U of T Statistical Sciences Department.

This presentation was prepared by Vivian Ngo.)

Github.com/vivianngo97/STA130-Fall-2019

viv.ngo@mail.utoronto.ca

Reminders

- Tutorials start 10 minutes after the hour

Agenda

- Vocabulary
- Group Discussion: Homework Question 1
- Writing Activity
- Mentorship Program

Vocabulary for this week's material

- Mean, average
- Median
- Standard deviation
- Variance
- Boxplot
- Interquartile range
- Quartile
- Proportion
- Outlier
- R object
- Vector
- Types of variables: e.g. character, numeric, logical
- Data frame
- Summary table, summary statistics

Vocabulary





- Mean, average
- Median
- **Standard deviation**
- Variance
- Boxplot
- Interquartile range
- Quartile
- Proportion
- Outlier
- R object
- Vector
- Types of variables: e.g. character, numeric, logical
- Data frame
- Summary table, summary statistics

- How far, on average, does our data deviate from the mean?
- Tells us about the spread
- Smaller standard deviation -> less spread

Vocabulary





- Mean, average
- Median
- Standard deviation
- Variance
- **Boxplot**
- Interquartile range
- Quartile
- Proportion
- Outlier
- R object
- Vector
- Types of variables: e.g. character, numeric, logical
- Data frame
- Summary table, summary statistics

- When to use boxplots?
  - To summarize the distribution of a quantitative variable
  - To compare distributions or summarize based on a categorical variable of interest
- Examples?

Vocabulary



- Mean, average
- Median
- Standard deviation
- Variance
- **Boxplot**
- Interquartile range
- Quartile
- Proportion
- Outlier
- R object
- Vector
- Types of variables: e.g. character, numeric, logical
- Data frame
- Summary table, summary statistics

Vocabulary







- Mean, average
- Median
- Standard deviation
- Variance
- Boxplot
- Interquartile range
- Quartile
- Proportion
- **Outlier**
- R object
- Vector
- Types of variables: e.g. character, numeric, logical
- Data frame
- Summary table, summary statistics

Vocabulary

```
> x<-1
> x
[1] 1

> x=1
> x
[1] 1

> x/10
[1] 0.1
```

Data Type	Description
Double (dbl)	Numbers (with or without decimals)
Integer (int)	Integers only (no decimals)
Character (chr)	Strings of letters and/or numbers and/or special characters surrounded by quotation marks
Logical (lgl)	TRUE or FALSE
Factor (fct)	A character type with a prespecified number and order of values (levels)

```
> die <- c(1,2,3,4,5,6)
> die
[1] 1 2 3 4 5 6
```

- Mean, average
- Median
- Standard deviation
- Variance
- Boxplot
- Interquartile range
- Quartile
- Proportion
- Outlier
- **R object**
- Vector
- Types of variables: e.g. character, numeric, logical
- Data frame
- Summary table, summary statistics

Vocabulary

min	mean	median	max	sd	n
9.5	1853.035	1001.7	60000	2646.909	6773

	GENDER	min	mean	median	max	sd	n
	<fct>	<dbl>	<dbl>	<dbl>	<dbl>	<dbl>	<int>
1	F	10	1864.	963.	60000	2761.	2582
2	M	9.5	1847.	1032.	59114.	2575.	4191

- Mean, average
- Median
- Standard deviation
- Variance
- Boxplot
- Interquartile range
- Quartile
- Proportion
- Outlier
- R object
- Vector
- Types of variables: e.g. character, numeric, logical
- Data frame
- **Summary table, summary statistics**

Group Discussion: Homework Question 1

- *For Question 1b, you used both histograms and boxplots to visualize your data. Which features were easier/harder to observe from each of the visualizations? In what situations may you want to choose a boxplot over a histogram, or vice versa? Explain.*





Writing Activity

Self-Reflection:

- *What questions, if any, do you have so far regarding the course materials?*
- *What is one of your favorite things about tutorial? Least favorite?*

Mentorship Program