Practice questions

Problem 1. For each of the logical formulas, indicate whether or not it is true when the domain of discourse is \mathbb{N} (the natural numbers 0, 1, 2, ...), \mathbb{Z} (the integers), \mathbb{Q} (the rationals), \mathbb{R} (the real numbers), and \mathbb{C} (the complex numbers).

$$\exists x \qquad (x^2 = 2) \\ \forall x \ \exists y \qquad (x^2 = y) \\ \forall y \ \exists x \qquad (x^2 = y) \\ \forall x \neq 0 \ \exists y \qquad (xy = 1) \\ \exists x \ \exists y \ (x+2y = 2) \land (2x+4y=5)$$

Problem 2. The goal of this problem is to translate some assertions about binary strings into logic notation. The domain of discourse is the set of all finite-length binary strings: λ , 0, 1, 00, 01, 10, 11, 000, 001, (Here λ denotes the empty string.) In your translations, you may use all the ordinary logic symbols (including =), variables, and the binary symbols 0, 1 denoting 0, 1.

A string like 01x0y of binary symbols and variables denotes the *concatenation* of the symbols and the binary strings represented by the variables. For example, if the value of x is 011 and the value of y is 1111, then the value of 01x0y is the binary string 0101101111.

Here are some examples of formulas and their English translations. Names for these predicates are listed in the third column so that you can reuse them in your solutions (as we do in the definition of the predicate NO-15 below).

Meaning	Formula	Name
x is a prefix of y	$\exists z \ (xz=y)$	PREFIX(x, y)
x is a substring of y	$\exists u \exists v \ (uxv = y)$	SUBSTRING(x, y)
x is empty or a string of 0's	$\neg SUBSTRING(1, x)$	NO-1s(x)

- (a) x consists of three copies of some string.
- **(b)** *x* is an even-length string of 0's.
- (c) x does not contain both a 0 and a 1.
- (d) x is the binary representation of $2^k + 1$ for some integer $k \ge 0$.