

Name : **Saroj Agarwal** Age/Gender : 63 y / F Collected : 26/03/2026 07:03
Patient ID : **0010663186** Visit Type : IP Received : 26/03/2026 07:50
Referred By : Dr. Darshit Shah DOB : 27/08/1962 Reported : 26/03/2026 10:04
Accession : 2600066852 Location : 1T13S`1TR1322`1TB1322`1ONCOMED

DEPARTMENT OF LABORATORY MEDICINE - HAEMATOLOGY

	Test Name	Unit	Reference Range	Low	Normal	High
	Activated Partial Thromboplastin Time (APTT)					
	APTT (Citrate Plasma, Nephelometry)	Secs	26.5-34.1			34.7
	Mean Normal APTT (Citrate Plasma)				30.3	
	APTT Ratio (Citrate Plasma, Calculated)				1.15	

APTT:

The activated partial thromboplastin time (APTT) test measures the functionality of the intrinsic and common pathways of the coagulation cascade. (factors XII, XI, IX, VIII, X, V, II and Fibrinogen).

Prolonged APTT tests may be due to:

- Inherited or acquired factor deficiencies.
- A nonspecific inhibitor such as the lupus anticoagulant (LA)
- Specific inhibitor: Antibodies that attack a particular factor.
- DIC, Fibrinolysis

Interference in APTT:

Preanalytical variables such as Insufficient/ excess sample, Heparin contamination from IV lines, Clotted blood samples, High hematocrit levels.

∞ End of Report ∞

Avatifdar

DR. ARCHANA VAZIFDAR

M.D (Pathology)
Head Of Hematopathology

DR. PREETI SINGH

M.D Pathology
Consultant Pathologist



Disclaimer: Tests marked with NABL symbol are accredited by NABL vide certificate number MC-7539