

Patient Name : SAROJ AGARWAL	Age/Sex : 63 Years / FEMALE
Ref. By : DR. SANJAY AGRAWAL	Date : 01/05/2026

RIGHT LOWER EXTREMITY ARTERIAL & VENOUS DOPPLER STUDY

Real time scan with colour flow mapping and pulsed Doppler analysis of arterial & venous drainage of right lower limb done.

GRAY SCALE IMAGING:

- Common femoral artery, superficial femoral artery, popliteal artery, anterior tibial, posterior tibial and peroneal arteries are normal in course and caliber
- Evidence of mild atheromatous plaque & intimo-medial thickening seen.
- E/o minimal to mild degree superficial subcutaneous fat plane inhomogeneous echogenicity involving right thigh region all around.
P/o changes of superficially subtle haemorrhage - Petechial lesions sequelae likely
- E/o moderate to severe degree of subcutaneous fat plane oedema changes noted involving right leg ankle foot region

Common Femoral vein, Superficial Femoral vein, Deep Femoral vein, Popliteal vein, Anterior Tibial vein, Posterior Tibial vein are normal in course and calibre.

Probe Compressibility: present in above mentioned veins.

COLOUR FLOW IMAGING

- Above mentioned arteries show normal colour filling.
- Normal colour filling also noted in dorsalis pedis artery.

Above mentioned veins show **normal Colour filling**.

Sapheno-Femoral junction show normal flow direction.

SPECTRAL ANALYSIS:

- Above mentioned arteries show normal wave form with triphasic pattern.
- No abnormal arterial occlusion seen

Above mentioned veins show normal wave form with normal Cardiac and Respiratory phasic variations.

- Augmentation ++
- No e/o DEEP VEIN THROMBOSIS - DVT

The superficial venous system was also examined.

No abnormal dilated greater saphanous varicocities are seen.
No varicocities are seen in the region of short sephanous vein territory.
There is no superficial vein thrombosis.

No e/o abnormal focal lesion or collection noted in examination of soft tissue plane of lower limb.

COMMENT:

- No signs of acute or chronic thrombosis in arterial or venous system
- No e/o hemodynamically significant stenosis or occlusion
- No abnormal varicocities or superficial - Deep venous thrombosis or abnormal perforators in lower limb
- No abnormal collection in soft tissue
- E/o minimal to mild degree superficial subcutaneous fat plane inhomogeneous echogenicity involving right thigh region all around.
P/o changes of superficially subtle haemorrhage - Petechial lesions sequelae likely
- E/o moderate to severe degree of subcutaneous fat plane oedema changes noted involving right leg ankle foot region

Adv : clinical - lab data correlation, Local & systemic examination correlation,
: SOS follow up with treatment / further evaluate if symptoms not relieved

Thanks for reference,

 **Dr. Jenish Shah**
— MBBS, MD Radiologist —