

Name : **Saroj Agarwal** Age/Gender : 63 y / F Collected : 20/02/2026 05:10
Patient ID : **0010663186** Visit Type : IP Received : 20/02/2026 06:51
Referred By : Dr. Darshit Shah DOB : 27/08/1962 Reported : 20/02/2026 09:47
Accession : 2600040488 Location : 1T06N`1TR0601`1TB0605`1ONCOMED

DEPARTMENT OF LABORATORY MEDICINE - BIOCHEMISTRY

	Test Name	Unit	Reference Range	Low	Normal	High
	Liver Profile					
	Bilirubin Total + Direct					
	Total Bilirubin (Serum, Diazo)	mg/dL	<= 1.2		0.33	
	Direct Bilirubin (Serum, Diazo)	mg/dL	<= 0.3		0.18	
	Indirect Bilirubin (Serum, Calculated)	mg/dL	<= 1.2		0.15	
	Alanine Aminotransferase [ALT/PT] (Serum, IFCC-Enzymatic with P5P)	U/L	<= 35		14.50	
	Aspartate Aminotransferase [AST/OT] (Serum, IFCC-Enzymatic with P5P)	U/L	<= 35		21.30	
	Gamma Glutamyltransferase [GGT] (Serum, Enzymatic IFCC)	U/L	< 40		26.30	
	Alkaline Phosphatase [ALP] (Serum, Enzymatic IFCC)	U/L	35-104		65.90	
	Proteins					
	Total Protein (Serum, Colorimetric - biuret)	g/dL	6.4-8.3	5.10		
	Albumin (Serum, Colorimetric - BCG)	g/dL	3.5-5.2	2.91		
	Globulin (Serum, Calculated)	g/dL	2-3.5		2.19	

Purva

DR. KSHAMA PIMPALGAONKAR
M D Biochemistry
Consultant Biochemist
Laboratory Medicine

DR. PURVA JAGETIA
M.D (Pathology)
Clinical Associate
Laboratory Medicine

DR. SRIKANT GHARPURE
M.D, DPB, DCP (London)
Consultant Pathologist
Laboratory Medicine



Name : **Saroj Agarwal** Age/Gender : 63 y / F Collected : 20/02/2026 05:10
Patient ID : **0010663186** Visit Type : IP Received : 20/02/2026 06:51
Referred By : Dr. Darshit Shah DOB : 27/08/1962 Reported : 20/02/2026 09:47
Accession : 2600040488 Location : 1T06N`1TR0601`1TB0605`1ONCOMED

DEPARTMENT OF LABORATORY MEDICINE - BIOCHEMISTRY

	Test Name	Unit	Reference Range	Low	Normal	High
	A / G ratio (Serum, Calculated)		1.0-2.5		1.33	

Total Bilirubin:

- Diseases or conditions which, through hemolytic processes, produce bilirubin faster than the liver can metabolize it, cause the levels of unconjugated (indirect) bilirubin to increase in the circulation.
- Liver immaturity and several other diseases in which the bilirubin conjugation mechanism is impaired cause similar elevations of circulating unconjugated bilirubin.
- Bile duct obstruction or damage to hepatocellular structure causes increases in the levels of both conjugated (direct) and unconjugated (indirect) bilirubin in the circulation.

∞ End of Report ∞

Purva

DR. KSHAMA PIMPALGAONKAR
M D Biochemistry
Consultant Biochemist
Laboratory Medicine

DR. PURVA JAGETIA
M.D (Pathology)
Clinical Associate
Laboratory Medicine

DR. SRIKANT GHARPURE
M.D, DPB, DCP (London)
Consultant Pathologist
Laboratory Medicine

