

Name : **Saroj Agarwal** Age/Gender : 63 y / F Collected : 08/03/2026 15:57
Patient ID : **0010663186** Visit Type : IP Received : 08/03/2026 17:20
Referred By : Dr. Darshit Shah DOB : 27/08/1962 Reported : 08/03/2026 19:01
Accession : 2600053233 Location : 1T05S`1TR0509`1TB0509`1ONCOMED

DEPARTMENT OF LABORATORY MEDICINE - BIOCHEMISTRY

	Test Name	Unit	Reference Range	Low	Normal	High
	Troponin I hs (Serum, CMIA)	pg/mL	<= 15.6		Less than 3.2	

Troponin I hs:

Troponin is a complex that regulates the contraction of striated muscle. It consists of 3 subunits (C, T, and I). Troponin I is an inhibitory protein and exhibits in 3 isoforms: cardiac muscle, slow-twitch skeletal muscle, and fast-twitch skeletal muscle. The cardiac specificity of this isoform improves the accuracy of diagnosis in patients with acute or chronic skeletal muscle injury and possible concomitant myocardial injury. Troponin I is released into the bloodstream within hours of the onset of symptoms of myocardial infarction or ischemic damage. It can be detected at 3 to 6 hours following onset of chest pain with peak concentrations at 12 to 16 hours, and remains elevated for 5 to 9 days. A positive troponin result is not always indicative of myocardial infarction. Other conditions resulting in myocardial cell damage can contribute to elevated cardiac troponin I levels. These conditions include, but are not limited to, myocarditis, cardiac surgery, angina, unstable angina, congestive heart failure and noncardiac-related causes, such as, renal failure and pulmonary embolism.

∞ End of Report ∞

DR. KSHAMA PIMPALGAONKAR

M D Biochemistry
Consultant Biochemist

DR. SRIKANT GHARPURE

M.D, DPB, DCP (London)
Consultant Pathologist

