

Name : **Saroj Agarwal** Age/Gender : 63 y / F Collected : 30/03/2026 23:06
Patient ID : **0010663186** Visit Type : IP Received : 31/03/2026 00:15
Referred By : Dr. Darshit Shah DOB : 27/08/1962 Reported : 31/03/2026 09:45
Accession : 2600070501 Location : 1T13S`1TR1322`1TB1322`1ONCOMED

DEPARTMENT OF LABORATORY MEDICINE - IMMUNOLOGY


	Test Name	Unit	Reference Range	Low	Normal	High
	Procalcitonin Quantitative (Serum,CMIA/Electrochemiluminescence)	ng/mL	<= 0.15			0.65

Procalcitonin Quantitative:

1. Procalcitonin (PCT) is an amino acid precursor of calcitonin peptide. PCT becomes detectable within 2 to 4 hours after a triggering event and peaks by 12 to 24 hours. PCT secretion parallels closely the severity of the inflammatory insult, with higher levels associated with more severe disease and declining levels with resolution of illness.
2. During severe systemic inflammation, especially related to bacterial infection, PCT and CCP-1 are secreted in large quantities. Noninfectious inflammatory stimuli need to be extremely severe to result in PCT elevations, making it a more specific marker for severe infections. In the absence of an ongoing stimulus. It is suitable for serial monitoring.
3. Reported results are for the information of the referring Clinician. Results should be interpreted in correlation with medical history, clinical presentation and other findings.
4. Reference ranges and clinical information/interpretations are traceable to kit inserts/textbooks/defined guidelines.

∞ End of Report ∞

DR. KSHAMA PIMPALGAONKAR
M D Biochemistry
Consultant Biochemist


DR. RINA SHAH
M.D Pathology
Consultant Pathologist

DR. SRIKANT GHARPURE
M.D, DPB, DCP (London)
Consultant Pathologist



Disclaimer: Tests marked with NABL symbol are accredited by NABL vide certificate number MC-7539