HIMGIRI INSTITUTE OF NUCLEAR MEDICINE

Radioiodine Therapy | PET-CT | Gamma Camera

Sahastradhara Road, Dehradun, Uttarakhand 248 013

Tel: 0135-2608 270 | Email: nm@himgiri-nuclear.example

AERB Reg. No. UK/NM/2017/0009

DISCHARGE SUMMARY

Radioiodine therapy ward (isolation)

Name:	**Ms. NANDITA RAWAT**	IP No.:	**HINM/IP/26/0071**
Age / Sex:	34 / F	Consultant:	Dr. Pallavi Dobhal (Nuclear Medicine)
Admitted:	20/04/2026	Discharged:	23/04/2026

Diagnosis

Papillary carcinoma thyroid, pT2 pN1a (AJCC 8th), ATA intermediate risk; status post total thyroidectomy with bilateral central compartment dissection (03/03/2026).

Pre-therapy work-up

Levothyroxine withdrawn for 4 weeks; low-iodine diet for 2 weeks. Blood tests on 17/04/2026: TSH 68.4 mIU/L, stimulated thyroglobulin 4.6 ng/mL, anti-thyroglobulin antibodies negative. Urine pregnancy test before therapy: negative.

Treatment

Oral radioiodine (I-131) 100 mCi administered on 20/04/2026 for adjuvant ablation. Kept in the shielded isolation room for 72 hours; dose rate at 1 metre at discharge 18 microSv/h.

Post-therapy scan

Whole-body I-131 scan on 22/04/2026: uptake in the thyroid bed (remnant) and a small focus in the right paratracheal region (nodal); no uptake in the lungs or skeleton.

Course

Mild neck discomfort and nausea on day 1, settled with ondansetron. No sialadenitis. Calcium normal on oral supplements.

SYNTHETIC DOCUMENT — FOR DEMONSTRATION ONLY

Medications on discharge

Medicine	Dose	When	Duration
Tab. Levothyroxine 100 mcg	1 tab	Empty stomach, 6 am	Long term (start 23/04/2026)
Tab. Calcium carbonate 1250 mg	1 tab	Twice daily after meals	1 month
Cap. Calcitriol 0.25 mcg	1 cap	Once daily	1 month
Tab. Ondansetron 4 mg	1 tab	If nausea	3 days
Tab. Pantoprazole 40 mg	1 tab	Before breakfast	7 days

Advice

- Drink plenty of water; suck lemon drops or chew sugar-free gum for 3 days to protect the salivary glands.
- For 7 days: sleep alone, keep 1 metre from adults and 2 metres from children and pregnant women; use a separate toilet and flush twice.
- Avoid pregnancy for 6 months after radioiodine.
- TSH and free T4 after 6 weeks; diagnostic whole-body scan with stimulated thyroglobulin after 6 months. Review in Nuclear Medicine OPD on 04/06/2026.

Dr. Pallavi Dobhal
M.D. (Nuclear Medicine)
Reg. No. UKMC-3412
Consultant Nuclear Medicine Physician

SYNTHETIC DOCUMENT — FOR DEMONSTRATION ONLY