Package 'IndexNumber'

October 12, 2022

Type Package

Title Index Numbers in Social Sciences
Version 1.3.2
Date 2021-03-15
Author Alejandro Saavedra-Nieves, Paula Saavedra-Nieves
Maintainer Alejandro Saavedra-Nieves <alejandro.saavedra.nieves@gmail.com></alejandro.saavedra.nieves@gmail.com>
Description We provide an R tool for teaching in Social Sciences. It allows the computation of index numbers. It is a measure of the evolution of a fixed magnitude for only a product of for several products. It is very useful in Social Sciences. Among others, we obtain simple index numbers (in chain or in serie), index numbers for not only a product or weighted index numbers as the Laspeyres index (Laspeyres, 1864), the Paasche index (Paasche, 1874) or the Fisher index (Lapedes, 1978).
License GPL-2
LazyData true
RoxygenNote 7.0.2
NeedsCompilation no
Repository CRAN
Date/Publication 2021-03-15 12:30:03 UTC
R topics documented:
IndexNumber-package
ActivePeople
aggregated.index.number
ECResources
edgeworth.index.number
fisher.index.number
Food
index.number.chain
index.number.serie
laspeyres.index.number

Index 15

IndexNumber-package Index Numbers in Social Sciences

Description

We provide an R tool for teaching in Social Sciences. It allows the computation of index numbers. It is a measure of the evolution of a fixed magnitude for only a product of for several products. It is very useful in Social Sciences. Among others, we obtain simple index numbers (in chain or in serie), index numbers for not only a product or weighted index numbers as the Laspeyres index (Laspeyres, 1864), the Paasche index (Paasche, 1874) or the Fisher index (Lapedes, 1978).

Details

The DESCRIPTION file:

Package: IndexNumber Type: Package

Title: Index Numbers in Social Sciences

Version: 1.3.2 Date: 2021-03-15

Author: Alejandro Saavedra-Nieves, Paula Saavedra-Nieves

Maintainer: Alejandro Saavedra-Nieves <alejandro.saavedra.nieves@gmail.com>

Description: We provide an R tool for teaching in Social Sciences. It allows the computation of index numbers. It is a meaning the computation of index numbers.

License: GPL-2 LazyData: true RoxygenNote: 7.0.2

Index of help topics:

ActivePeople Economically active people in Spain from 2002

to 2019.

ECResources Combustibles and energy resources for the main

home in Spain from 2006 to 2015.

Food Food in Spain from 2006 to 2015. IndexNumber-package Index Numbers in Social Sciences

Mortgages Mortgages constituted on urban properties in

Spain from 2003 to 2018.

aggregated.index.number

Calculate an aggregate index number

edgeworth.index.number

Calculate the Edgeworth index number

fisher.index.number
Calculate the Fisher index number
index.number.chain
Calculate an index number in chain
Calculate an index number in serie

laspeyres.index.number

ActivePeople 3

```
Calculate the Laspeyres index number paasche.index.number Calculate the Paasche index number
```

Once we have defined a magnitude for a product (of several products), we can analyse how it (they) evolves along the time. Index Numbers model this effect in Social Science. In this sense, several approaches may be considered. We include in this package several options of analysing this problem.

Author(s)

Alejandro Saavedra-Nieves, Paula Saavedra-Nieves

Maintainer: Alejandro Saavedra-Nieves <alejandro.saavedra.nieves@gmail.com>

References

- (2004) Consumer Price Index Manual: Theory and Practice. ILO, IMF. CPI Manual OECD, UN, Eurostat, and The World Bank by ILO, Geneva.

Index Number (2008) In: The Concise Encyclopedia of Statistics. Springer, New York, NY. <doi:https://doi.org/10.1007/978-0-387-32833-1>.

Laspeyres, E. (1871) Die Berechnung einer mittleren Waarenpreissteigerung. Jahrb. Natl. Stat. 16, 296–314.

Paasche, H. (1874) Uber die Preisentwicklung der letzten Jahre nach den Hamburger Borsennotirungen. Jahrb. Natl. Stat. 23, 168–178.

Examples

```
prices<-c(70,75,77,77,85,90,85,75,77,87,85,90,70,75,77,77,85,90)
index.number.serie(prices,name="Prices",opt.plot=TRUE,opt.summary=TRUE)</pre>
```

ActivePeople

Economically active people in Spain from 2002 to 2019.

Description

Number (thousands) of economically active women and men in Spain between 2002 and 2019

Usage

```
data(ActivePeople)
```

Format

A data frame with columns:

Time A trimester (T1, T2, T3 and T4) between 2002 and 2019.

TotalWomen Number (thousands) of economically active women.

TotalMen Number (thousands) of economically active men.

Source

Spanish Statistical Office (INE), http://www.ine.es

Examples

```
## Not run:
ActivePeople
## End(Not run)
```

aggregated.index.number

Calculate an aggregate index number

Description

This function determines index numbers without weights for those cases in which there exist more than an only product (in chain or in serie)

Usage

```
aggregated.index.number(x, base, type, name,opt.plot=FALSE, opt.summary=FALSE)
```

Arguments

It is a matrix containing that magnitude to be studied. In each column, it con-Х

tains the magnitud of a different product. Thus, we have nrow(x) values of a

magnitud for ncol(x) products.

base Chain of characters that indicates the nature of the index number. If we intro-

> duce base="serie", we compare each value with respect to the initial one. In this case, it is said to be an index number in serie. Otherwise, if we introduce base="chain", we obtain the index number in chain, by comparing each value

with the immediately previous value.

Chain of characters to indicate the type of non-weighted index number to evaltype uate the evolution of a set of magnitudes (even for different products).

> By considering base="serie", if we introduce type="arithmetic", we obtain the Sauerbeck index number. If we introduce type="geometric", we obtain the Geometric index. If we choose type="harmonic", we obtain the Harmonic mean index. If we write type="BDutot", we will obtain the Bradstreet-Dutot

index.

Otherwise, if we take base="chain" and type="Carli", we obtain the Carli index number. If we introduce type="Jevons", we obtain the Jevons index and

if we choose type="Dutot", we obtain the Dutot index.

Chain of characters to indicate the name of the variable under study. name

opt.plot Logical option to indicate if a graphical description of the index number along

the different stages is required. It takes the value TRUE or FALSE.

ECResources 5

opt.summary Logical option to indicate if a statistical summary of the index number is required. It takes the value TRUE or FALSE.

Value

Summary Statistical summary (optional) of the index number along the considered period.

Agg. index number

Table containing the values of the index number for the considered stages and

the aggregate value.

Graphical Graphical summary (optional) of the index number along the considered period.

Author(s)

A. Saavedra-Nieves, P. Saavedra-Nieves

References

CPI Manual (2004). Consumer Price Index Manual: Theory and Practice. OECD, UN, Eurostat, and TheWorld Bank by ILO, Geneva.

Index Number (2008). In: The Concise Encyclopedia of Statistics. Springer, New York, NY. <doi:https://doi.org/10.1007/978-0-387-32833-1>

Examples

```
prices<-matrix(c(70,75,77,77,85,90,85,75,77,87,85,90,70,75,77,77,85,90),ncol=3)
aggregated.index.number(prices,"chain","geometric","Price",opt.plot=TRUE,opt.summary=TRUE)</pre>
```

ECResources Combustibles and energy resources for the main home in Spain from 2006 to 2015.

Description

Unitary value (euros) and consumed amount (thousands of units) of combustibles and other energy resources for the main home in Spain from 2006 to 2015.

Usage

data(ECResources)

Format

A data frame with columns:

Time Year between 2006 and 2015.

ElectricityPrice Unitary value of electricity (KWh).

NaturalGasPrice Unitary value of natural gas (m3).

LiquifiedGasPrice Unitary value of liquified gas (kilo).

LiquifiedCombustiblesPrice Unitary value of liquified combustibles (1).

SolidCombustiblesPrice Unitary value of Solid combustibles(1).

ElectricityConsumed Consumed (thousands of units) of electricity (KWh).

NaturalGasConsumed Consumed (thousands of units) of natural gas (m3).

LiquifiedGasConsumed (thousands of units) of liquified gas (kilo).

LiquifiedCombustiblesConsumed (thousands of units) of liquified combustibles (1).

SolidCombustiblesConsumed (thousands of units) of solid combustibles (1).

Source

```
Spanish Statistical Office (INE), http://www.ine.es
```

Examples

```
## Not run:
   ECResources
## End(Not run)
```

edgeworth.index.number

Calculate the Edgeworth index number

Description

This function determines the Marshall-Edgeworth index number for those cases in which there exist more than an only product.

Usage

```
edgeworth.index.number(x, y, name, opt.plot = FALSE, opt.summary = FALSE)
```

Arguments

X	Matrix that contains the magnitude(s) under study. In each column, it contains the magnitude of a different product considered. Thus, we have nrow(x) values of a magnitude for ncol(x) products.
У	Matrix that contains that magnitude used as weight. In each column, it contains another magnitude associated to each different product along the time. Thus, we have $nrow(x)$ values of magnitudes for the set of $ncol(x)$ products.
name	Chain of characters to indicate the name of the variable under study.
opt.plot	Logical option to indicate if a graphical descriptiony of the index number along the different stages is required. It takes the value TRUE or FALSE.
opt.summary	Logical option to indicate if a statistical summary of the index number is required. It takes the value TRUE or FALSE.

fisher.index.number 7

Value

Summary Statistical summary (optional) of the index number along the considered period. Agg. index number

> Table containing the values of the index number for the considered stages and the aggregate value.

Graphical Graphical summary (optional) of the index number along the considered period.

Author(s)

A. Saavedra-Nieves, P. Saavedra-Nieves

References

Edgeworth, F. (1887) Measurement of change in value of money i. First Memorandum presented to the British Association for the Advancement of Science. Reprinted in his Papers Relating to Political Economy, 1, 198–259.

Marshall, A. (1887) Remedies for fluctuations of general prices. The Contemporary review, 1866-1900, 51, 355-375.

Examples

```
prices<-matrix(c(70,75,77,77,85,90,85,75,77,87,85,90,70,75,77,77,85,90),ncol=3)
weights < -matrix(c(1,1.5,0.8,1.1,1,0.9,0.7,0.8,0.6,1,1.1,0.9,1,1,0.9,1.1,0.6,0.8),ncol=3)
edgeworth.index.number(prices,weights,"Price",opt.plot=TRUE,opt.summary=TRUE)
```

fisher.index.number

Calculate the Fisher index number

Description

This function determines the Fisher index number for those cases in which there exist more than an only product.

Usage

```
fisher.index.number(x, y, name, opt.plot = FALSE, opt.summary = FALSE)
```

Arguments

X	It is a matrix containing that magnitude to be studied. In each column, it con-
	tains the magnitud of a different product. Thus, we have nrow(x) values of a magnitud for ncol(x) products.
у	It is a matrix containing that magnitude used as weight. In each column, it contains another magnitud of the different products along the time. Thus, we

have nrow(x) values of a magnitud for ncol(x) products. Chain of characters to indicate the name of the variable under study.

name

8 Food

opt.plot	Logical option to indicate if a graphical descriptiony of the index number along the different stages is required. It takes the value TRUE or FALSE.
opt.summary	Logical option to indicate if a statistical summary of the index number is required. It takes the value TRUE or FALSE.

Value

Summary Statistical summary (optional) of the index number along the considered period.

Agg. index number

Table containing the values of the index number for the considered stages and

the aggregate value.

Graphical Graphical summary (optional) of the index number along the considered period.

Author(s)

A. Saavedra-Nieves, P. Saavedra-Nieves

References

Fisher, I. (1922) The making of index numbers: a study of their varieties, tests, and reliability, volume 1. Houghton Mifflin.

Lapedes, Daniel N. (1978) Dictionary of Physics and Mathematics. McGrow–Hill. p. 367. ISBN 0-07-045480-9.

Examples

```
\label{eq:prices-matrix} $$ prices - matrix(c(70,75,77,77,85,90,85,75,77,87,85,90,70,75,77,77,85,90), ncol=3) $$ weights - matrix(c(1,1.5,0.8,1.1,1,0.9,0.7,0.8,0.6,1,1.1,0.9,1,1,0.9,1.1,0.6,0.8), ncol=3) $$ fisher.index.number(prices,weights,name="Price",opt.plot=TRUE,opt.summary=TRUE) $$
```

Food

Food in Spain from 2006 to 2015.

Description

Unitary value (euros) and consumed amount (thousands of units) of food in Spain from 2006 to 2015.

Usage

data(Food)

Food 9

Format

A data frame with columns:

Year Year from 2006 and 2015.

RicePrice Unitary value of rice (kilo).

BreadPrice Unitary value of bread (kilo).

PorkPrice Unitary value of pork meat (kilo).

FishPrice Unitary value of fish (kilo).

WholeMilkPrice Unitary value of whole milk (1).

EggsPrice Unitary value of eggs (unit).

OliveOilPrice Unitary value of olive oil (1).

ApplesPrice Unitary value of apples (kilo).

DriedFruitAndNutsPrice Unitary value of dried fruit and nuts (kilo).

GreenVegetablePrice Unitary value value of green vegetables (kilo).

PotatoesPrice Unitary value value of potatoes (kilo)

SugarPrice Unitary value value of sugar (kilo)

ChocolatePrice Unitary value value of chocolate (kilo)

CoffeePrice Unitary value value of coffee (1).

MineralWaterPrice Unitary value value of mineral water (1).

WinePrice Unitary value value of wine (1).

BeerPrice Unitary value value of beer (1).

RiceConsumed Total amount (thousands of units) of consumed rice (kilo).

BreadConsumed Total amount (thousands of units) of consumed bread (kilo).

PorkConsumed Total amount (thousands of units) of consumed pork meat (kilo).

FishConsumed Total amount (thousands of units) of consumed fish (kilo).

WholeMilkConsumed Total amount (thousands of units) of consumed whole milk (l).

EggsConsumed Total amount (thousands of units) of consumed eggs (unit).

OliveOilConsumed Total amount (thousands of units) of consumed olive oil (l).

ApplesConsumed Total amount (thousands of units) of consumed apples (kilo).

DriedFruitAndNutsConsumed Total amount (thousands of units) of consumed dried fruit and nuts (kilo).

GreenVegetableConsumed Total amount (thousands of units) of consumed green vegetables (kilo).

PotatoesConsumed Total amount (thousands of units) of consumed potatoes (kilo)

SugarConsumed Total amount (thousands of units) of consumed sugar (kilo)

ChocolateConsumed Total amount (thousands of units) of consumed chocolate (kilo)

CoffeeConsumed Total amount (thousands of units) of consumed coffee (1).

MineralWaterConsumed Total amount (thousands of units) of consumed mineral water (1).

WineConsumed Total amount (thousands of units) of consumed wine (1).

BeerConsumed Total amount (thousands of units) of consumed beer (1).

10 index.number.chain

Source

```
Spanish Statistical Office (INE), http://www.ine.es
```

Examples

```
## Not run:
Food
## End(Not run)
```

index.number.chain

Calculate an index number in chain

Description

This function determines index numbers "in chain" for those cases with an only product.

Usage

```
index.number.chain(x, name, opt.plot = FALSE, opt.summary = FALSE)
```

Arguments

X	It is a vector containing that magnitude to be studied for a product. Thus, we
	have length(x) values of it.

name Chain of characters to indicate the name of the variable under study.

opt.plot Logical option to indicate if a graphical descriptiony of the index number along

the different stages is required. It takes the value TRUE or FALSE.

opt.summary Logical option to indicate if a statistical summary of the index number is re-

quired. It takes the value TRUE or FALSE.

Value

Summary	Statistical summary (optional) of the index number along the considered period.
Index number	Table containing the values of the index number for the considered stages.
Graphical	Graphical summary (optional) of the index number along the considered period.

Author(s)

A. Saavedra-Nieves, P. Saavedra-Nieves

References

Index Number (2008) In: The Concise Encyclopedia of Statistics. Springer, New York, NY. <doi:https://doi.org/10.1007/978-0-387-32833-1>.

index.number.serie 11

Examples

```
prices<-c(70,75,77,77,85,90,85,75,77,87,85,90,70,75,77,77,85,90)
index.number.chain(prices,"Prices",opt.plot=TRUE,opt.summary=TRUE)</pre>
```

index.number.serie

Calculate an index number in serie

Description

This function determines index numbers "in serie" for those cases with an only product.

Usage

```
index.number.serie(x, name, opt.plot = FALSE, opt.summary = FALSE)
```

Arguments

X	It is a vector containing that magnitude to be studied for a product. Thus, we have length(x) values of it.
name	Chain of characters to indicate the name of the variable under study.
opt.plot	Logical option to indicate if a graphical descriptiony of the index number along the different stages is required. It takes the value TRUE or FALSE.
opt.summary	Logical option to indicate if a statistical summary of the index number is required. It takes the value TRUE or FALSE

Value

Summary	Statistical summary (optional) of the index number along the considered period.
Index number	Table containing the values of the index number for the considered stages.
Graphical	Graphical summary (optional) of the index number along the considered period.

Author(s)

A. Saavedra-Nieves, P. Saavedra-Nieves

References

Index Number (2008). In: The Concise Encyclopedia of Statistics. Springer, New York, NY. <doi:https://doi.org/10.1007/978-0-387-32833-1>

Examples

```
prices<-c(70,75,77,77,85,90,85,75,77,87,85,90,70,75,77,77,85,90)
index.number.serie(prices,"Prices",opt.plot=TRUE,opt.summary=TRUE)</pre>
```

laspeyres.index.number

Calculate the Laspeyres index number

Description

This function determines the Laspeyres index number for those cases in which there exist more than an only product.

Usage

laspeyres.index.number(x, y, name, opt.plot = FALSE, opt.summary = FALSE)

Arguments

x	Matrix that contains the magnitude(s) under study. In each column, it contains the magnitude of a different product considered. Thus, we have nrow(x) values of a magnitude for ncol(x) products.
у	Matrix that contains that magnitude used as weight. In each column, it contains another magnitude associated to each different product along the time. Thus, we have nrow(x) values of magnitudes for the set of ncol(x) products.
name	Chain of characters to indicate the name of the variable under study.
opt.plot	Logical option to indicate if a graphical descriptiony of the index number along the different stages is required. It takes the value TRUE or FALSE.
opt.summary	Logical option to indicate if a statistical summary of the index number is required. It takes the value TRUE or FALSE.

Value

Summary	Statistical summary (optional) of the index number along the considered period.
Agg. index number	er
	Table containing the values of the index number for the considered stages and the aggregate value.
Graphical	Graphical summary (optional) of the index number along the considered period.

Author(s)

A. Saavedra-Nieves, P. Saavedra-Nieves

References

Laspeyres, E. (1864) Hamburger Warenpreise 1850–1863 und die kalifornisch-australischen Geldentdeckung seit. Jahrb. Natl. Stat. 3, 81–118, 209–236.

Laspeyres, E. (1871) Die Berechnung einer mittleren Waarenpreissteigerung. Jahrb. Natl. Stat. 16, 296–314.

Mortgages 13

Examples

```
\label{local_prices} $$ prices < -matrix(c(70,75,77,77,85,90,85,75,77,87,85,90,70,75,77,77,85,90),ncol=3) $$ weights < -matrix(c(1,1.5,0.8,1.1,1,0.9,0.7,0.8,0.6,1,1.1,0.9,1,1,0.9,1.1,0.6,0.8),ncol=3) $$ laspeyres.index.number(prices,weights,"Price",opt.plot=TRUE,opt.summary=TRUE) $$
```

Mortgages Constituted on urban properties in Spain from 2003 to 2018.

Description

Number of mortgages constituted on urban properties and mortgages amounts (thousands of euros) from 2003 to 2018.

Usage

```
data(Mortgages)
```

Format

A data frame with columns:

Year Year from 2003 to 2018

Number.of.bank.mortgages Numbers of bank mortages.

Amount.of.bank.mortgages Amount (thousands of euros) of bank mortages.

Number.of.savings.bank.mortgages Numbers of savings bank mortages.

Amount.of.savings.bank.mortgages Amount (thousands of euros) of savings bank mortages.

Number.of.other.entities.mortgages Numbers of other entities mortages.

Amount.of.other.entities.mortgages Amount (thousands of euros) of other entities mortages.

Source

```
Spanish Statistical Office (INE), http://www.ine.es
```

Examples

```
## Not run:
  Mortgages
## End(Not run)
```

14 paasche.index.number

paasche.index.number Calculate the Paasche index number

Description

This function determines the Paasche index number for those cases in which there exist more than an only product.

Usage

```
paasche.index.number(x, y, name, opt.plot = FALSE, opt.summary = FALSE)
```

Arguments

х	Matrix that contains the magnitude(s) under study. In each column, it contains the magnitude of a different product considered. Thus, we have $nrow(x)$ values of a magnitude for $ncol(x)$ products.
У	Matrix that contains that magnitude used as weight. In each column, it contains another magnitude associated to each different product along the time. Thus, we have nrow(x) values of magnitudes for the set of ncol(x) products.
name	Chain of characters to indicate the name of the variable under study.
opt.plot	Logical option to indicate if a graphical descriptiony of the index number along the different stages is required. It takes the value TRUE or FALSE.
opt.summary	Logical option to indicate if a statistical summary of the index number is required. It takes the value TRUE or FALSE.

Value

Summary Statistical summary (optional) of the index number along the considered period.

Agg. index number

Table containing the values of the index number for the considered stages and the aggregate value.

Graphical Graphical summary (optional) of the index number along the considered period.

Author(s)

A. Saavedra-Nieves, P. Saavedra-Nieves

References

Paasche, H. (1874) Uber die Preisentwicklung der letzten Jahre nach den Hamburger Borsennotirungen. Jahrb. Natl. Stat. 23, 168–178.

Examples

```
prices<-matrix(c(70,75,77,77,85,90,85,75,77,87,85,90,70,75,77,77,85,90),ncol=3)
weights<-matrix(c(1,1.5,0.8,1.1,1,0.9,0.7,0.8,0.6,1,1.1,0.9,1,1,0.9,1.1,0.6,0.8),ncol=3)
paasche.index.number(prices,weights,"Price",opt.plot=TRUE,opt.summary=TRUE)</pre>
```

Index

```
* Fisher
    fisher.index.number, 7
* IndexNUMBER
    IndexNumber-package, 2
* aggregated
    {\it aggregated.index.number, 4}
* datasets
    ActivePeople, 3
    ECResources, 5
    Food, 8
    Mortgages, 13
* edgeworth
    edgeworth.index.number, 6
* indexnumber chain
    index.number.chain, 10
* indexnumberserie
    index.number.serie, 11
* laspeyres
    laspeyres.index.number, 12
* paasche
    paasche.index.number, 14
ActivePeople, 3
aggregated.index.number, 4
ECResources, 5
edgeworth.index.number, 6
fisher.index.number, 7
Food, 8
index.number.chain, 10
index.number.serie, 11
IndexNumber (IndexNumber-package), 2
IndexNumber-package, 2
laspeyres.index.number, 12
Mortgages, 13
paasche.index.number, 14
```