Conditioned States as Toric Kähler Data

A companion note to "Probability, Context and Differentiation: The Layers"
(working draft)

Vinay K Awasthi
Johns Hopkins University
vawasth1@jh.edu

November 2026

Abstract

The central point of this note is that *leading-order balance identifies candidates, not the selected state*. On a compact Delzant polytope with potential $W_T = \sum_k T^{-\lambda_k} y^{v_k}$, the unique positive critical point (Galkin's lemma) has a valuation u^* at which the leading terms balance with positive weights: Birch at target 0. The converse fails; on F_1 a whole segment balances and higher order selects. A selection theorem settles which point is chosen: u^* is the lexicographic max–min point, with error $O(1/\log(1/T))$. The positive orthant is a Lefschetz thimble, and its law $e^{-W/\hbar}dy/y$ has a tropical limit: uniform on shrunk polytopes, then concentrated on the candidates, level by level. Around this sits the toric Kähler dictionary in probabilistic terms: Fisher metric, relative entropy, Duistermaat–Heckman. Soft evidence of size n shifts $\lambda \mapsto \lambda - (n/N)\rho$, and acts only through circuits; F_1 has one, with a wall inside the Kähler cone where the leading set jumps.

Part II reads this on contingency tables. The margins are a point of the base and the Kähler class of the fibre, the coupling polytope. On the fibre, Tier A is Fisher's noncentral hypergeometric family, the best smallest cell is $\min(f_i/m, g_j/n)$, and the selected coupling is the facet-leximin one. One count of data can cross a leximin wall and change the Floer fibres without changing the surface.

Part III turns to morphisms. By the cited theorems, distinct objects are orthogonal, each has a Clifford endomorphism algebra (for the probabilistic object, of its Fisher information), and no higher products survive. Across an outcome wall the category over Λ is unchanged up to A_∞ equivalence. But the probabilistic object's Clifford form, reduced mod T, drops rank, and the Stokes walls, where thimble bases mutate, crowd into an $O(1/\log(1/T))$ window. Floer statements are cited, not proved. A ledger records every operation.

Part IV goes to 2×4, where the fibre is three-dimensional. Its 148 Kähler chambers are classified exactly, with ampleness certified by integer intersection numbers on invariant curves (Nakai–Moishezon, in its toric form): 14 are Fano and 134 weak Fano, and coupled three-step selection occurs only in the weak Fano ones. In three examples, one per role, the Clifford form of the positive object has one scale per dimension that selection removes, including a double scale where one level removes two; we state this as a conjecture. A path through the column-binding example (Stage IV) crosses two leximin walls in one Kähler chamber: the example itself sits on a level-two wall, which forces a collision of critical points, while the level-one wall is crossed without one. 74 numerical, 4 certificate and 3 ledger checks corroborate the statements.

Part I
The geometry

1 What this note does and does not do

The layers paper ends its toric section with an outlook remark: the precise dictionary is between conditioned states and moment fibres, not between sources and Lagrangians [7, Remark "outlook: a symplectic reading"]. This note makes that remark exact in two places and leaves the rest open. Its components have different status.

component	status	source
toric Kähler structure, moment map, Fisher metric, symplectic potential, Duistermaat–Heckman	Used	[29, 1, 18, 2]
Birch's theorem; relative interiors of convex hulls	Used	[8, 36]
one positive critical point (theorem 3(1)); the positive locus as a Lefschetz thimble	Used	Galkin's lemma [25]; [26, 31]
the valuation u^* maximises $\min_k \ell_k$; a recursive construction of u^*	Used	[32, Thm. 1.2]; [23]
the bridge (theorem 3(2)–(4)): leading-order balance, candidates versus selection, the sign split	Derived	this note
the orthant is the ascending thimble J_+; Laplace concentration (proposition 7)	Adapted	[25, 26]
the tropical limit of the thimble law (theorem 8) and its F_1 levels (example 9)	New	this note
linear-programming duality, strict complementarity; regular subdivisions and secondary fans	Used	[37, 28, 35]
soft evidence as $\lambda - s\rho$; the balance programme; candidates = relative interior of argmax (definition 10 and proposition 11)	Derived	this note
where data can act: circuits, least evidence, the F_1 wall (corollary 12 and example 14)	New	this note
coupling polytopes are Delzant; Kähler class = product coupling; chambers and blow-ups	Used	[5]
Tier A on the fibre is Fisher's noncentral hypergeometric family (proposition 22)	Adapted	[22, 9]
the max–min coupling $\min(f_i/m, g_j/n)$; leximin walls; the one-count example (theorem 23 and example 25)	New	this note
selection: the positive valuation is the lexicographic max–min point (theorem 27)	Adapted	[23, 32, 15]; proof here
morphisms: orthogonality, Clifford endomorphisms, summands (proposition 31)	Cited	[11, 12, 40]
no higher products; A_∞ equivalence across the F_1 wall (proposition 32 and theorem 34)	Derived (on cited m_2)	this note
the Clifford shadow of selection; Stokes walls crowd the outcome wall (theorem 33 and proposition 35)	New (35: computed)	this note
the column-binding example classified; the reduction of the Clifford form; its Hom table (propositions 36 and 39 and theorem 38)	New (Hom: on cited theorems)	this note
Stage IV: a path across two leximin walls; the collision at the level-two wall (propositions 40 and 41)	New (exact tropical; collision numerical)	this note
Floer cohomology of toric fibres and the potential W	Cited, not proved	[11, 23]
ampleness by invariant curves: Nakai–Moishezon, toric Kleiman, wall relations (proposition 42)	Used	[33, 14]
the 148 chambers of 2×4; selection depth in the Fano chambers; examples A, B, C (theorem 43 and propositions 44 and 46)	New (exact computation)	this note
weak Fano corrections $W^{\mathrm{LF}} = \sum(1+\delta_k)T^{-\lambda_k}y^{v_k}$ (remark 48)	Cited	[13]
one scale per removed dimension (Conjecture 50); proved at leading order in A, B, C (proposition 51)	Conjecture; New	this note
Novikov weights against the environment orders of the layers paper; decision, contextuality and VOI walls	Open	section 14

Two probability structures must be kept apart. The measures in this note live on *conditioned states*, points of the polytope. They are not the source weights π_l of the layers paper. The uniform measure of proposition 1(5) is a prior on states, not on sources.

2 Tier A: the toric Kähler structure of conditioned states

Setting. Let $S = \{S_i\} \subset \mathbb{Z}^d$ be a finite support whose affine span is $\mathbb{R}^d$, with coefficients $c_i > 0$. In log coordinates $u = \log x$ put

$$\varphi(u) = \log \sum_i c_i e^{\langle S_i, u\rangle}, \qquad q_i(u) = \frac{c_i e^{\langle S_i, u\rangle}}{\sum_j c_j e^{\langle S_j, u\rangle}},$$

so $q(u)$ is the monomial law of the layers paper and $\mu(u) = \sum_i q_i S_i$ is its moment map. On the torus $(\mathbb{C}^*)^d$, write $z_j = e^{(u_j + i\theta_j)/2}$.

Proposition 1 (the toric Kähler data)**.** *(1) $F(z) = \varphi(\log|z|^2)$ is a Kähler potential for a torus-invariant Kähler form ω on $(\mathbb{C}^*)^d$, the pullback of the Fubini–Study form by the monomial map $z \mapsto [\sqrt{c_i}\, z^{S_i}]$. In the coordinates (u, θ), up to the normalising constant of one's convention, $\omega = \sum_{j,k} \varphi_{jk}(u)\, du_j \wedge d\theta_k$, and the moment map of the rotation action is $\nabla\varphi = \mu$.*

(2) Fisher information is the metric. *The Hessian $\varphi_{jk} = \mathrm{Cov}_{q(u)}(S)_{jk}$ is the Fisher information of the exponential family $q(u)$ in its natural parameter, and the Kähler metric is $\sum \varphi_{jk}(du_j du_k + d\theta_j d\theta_k)$.*

(3) The positive real locus is a Lagrangian section. *$\{\theta = 0\}$ is Lagrangian, and it meets each moment fibre $\mu^{-1}(p)$, p in the interior of $Q = \mathrm{conv}(S)$, in exactly one point (Birch).*

(4) The symplectic potential is a relative entropy. *The Legendre dual $\psi(p) = \sup_u(\langle p, u\rangle - \varphi(u))$ equals $\sum_i q_i \log(q_i/c_i)$ evaluated at the exponential-family law $q = q(u(p))$. That law is the unique minimiser of $D(q\|c) = \sum_i q_i \log(q_i/c_i)$ among the laws on S with mean p; many other laws have mean p. So $\psi(p) = \min\{D(q\|c) : \sum_i q_i S_i = p\}$, the entropy of the minimiser relative to the weights c, with the sign reversed. Moreover $\nabla\psi(p) = u(p)$ and $\mathrm{Hess}\,\psi = (\mathrm{Hess}\,\varphi)^{-1}$, the Fisher information in the expectation parameter.*

(5) Duistermaat–Heckman. *The pushforward of the Liouville measure $\omega^d/d!$ by μ is $(2\pi)^d$ times Lebesgue measure on the interior of Q. In natural parameters it has density* $\det \mathrm{Hess}\,\varphi$.

Proof. These are classical. For (1), see [29, 1] and [24, Section 4.2]. (2) holds because φ is the log-partition function of the exponential family [2]. (3): on $\{\theta = 0\}$ every $d\theta_k$ vanishes, and a moment fibre is $\{u(p)\} \times T^d$, which meets $\theta = 0$ once because $u(p)$ is unique [8]. (4) is Legendre duality for a strictly convex function, and $\langle p, u\rangle - \varphi(u) = \sum_i q_i \log(q_i/c_i)$ at $\mu(u) = p$. That $q(u(p))$ minimises $D(\cdot\|c)$ under the mean constraint is the Lagrange (Gibbs) characterisation of exponential families [2]. (5) is the change of variables $p = \nabla\varphi(u)$, whose Jacobian is $\mathrm{Hess}\,\varphi$ [18]; the θ-integral gives $(2\pi)^d$. Checks: KB1 checks (1)–(2) on 30 random supports, against finite differences. KB2 checks (4) on 20 supports. KB3 checks (5) by comparing $\int \det \mathrm{Hess}\,\varphi\, du$ over boxes with the areas of their moment images. □

In general this structure lives on the open orbit $(\mathbb{C}^*)^d$ only. A Delzant polytope Q determines a compact symplectic toric manifold X_Q [16], but a metric built from an arbitrary support need not extend smoothly to it. It does extend in one case. Suppose S is the set of lattice points of a Delzant lattice polytope Q. On a smooth projective toric variety an ample line bundle is very ample [24, Section 3.4]. So the monomial map extends to an embedding of X_Q in $\mathbb{P}^{|S|-1}$, and ω is the restriction of the Fubini–Study form; the weights c_i act by a diagonal automorphism. The polytope of the layers paper's blueprint is not Delzant: its vertex at $(1, 1)$ has determinant 3. So the blueprint gives an orbifold, and the examples below are $\mathbb{CP}^2$ and F_1.

3 The bridge

Setting. Let $Q = \{p : \ell_k(p) = \langle v_k, p\rangle - \lambda_k \geq 0,\ k = 1, \dots, m\}$ be a compact Delzant polytope, with primitive inward facet normals v_k. Its potential, in the convention of Hori–Vafa and Cho–Oh [30, 11], is

$$W_T(y) = \sum_k T^{-\lambda_k} y^{v_k}, \qquad y \in (\mathbb{C}^*)^n,\ 0 < T < 1.$$

A family of critical points y_T has *valuation* $u = \lim_{T\to 0} \log|y_T| / \log T$ when the limit exists. At u, the term k has order $\ell_k(u)$, and the *leading terms* are $K(u) = \{k : \ell_k(u) \text{ minimal}\}$. Say the leading terms *balance* at u if $0 = \sum_{k\in K(u)} a_k v_k$ for some $a_k > 0$.

Lemma 2 (balance is Birch at target zero)**.** *For vectors v_k, $k \in K$, the following are equivalent:*

- *there are weights $a_k > 0$ with $\sum_k a_k v_k = 0$;*
- *there is a law q with $q_k > 0$ for every $k \in K$ and mean $\sum_k q_k v_k = 0$;*
- *0 lies in the relative interior of $\operatorname{conv}\{v_k : k \in K\}$.*

When they hold, the positive monomial law with support $\{v_k\}$ and mean 0 is unique, up to the torus directions orthogonal to the span of the differences (Birch).

Proof. Normalising the weights gives the first equivalence. The relative interior of the convex hull of finitely many points is the set of their convex combinations with all coefficients positive [36, Theorem 6.9]. Uniqueness is Birch's theorem [8], which is [7, Theorem "the toric realization"(1)] with target 0. □

Theorem 3 (the Birch bridge)**.** *Let the λ_k be rational.*

(1) One positive critical point *(Galkin's lemma [25]; Used). For each $T \in (0,1)$, W_T has exactly one critical point y_T in the positive torus $(0,\infty)^n$. A positive y is critical iff the law $w_k \propto T^{-\lambda_k} y^{v_k}$ of the terms of W_T has mean exponent $\sum_k w_k v_k = 0$: Birch at target 0.*

(2) Its valuation balances. *The valuation u^* of y_T exists. As $T \to 0$ the law w tends to a law supported exactly on $K(u^*)$, with positive weights and mean 0. So the leading terms balance at u^*.*

(3) Candidates versus selection. *The converse of (2) fails: the set of points where the leading terms balance can be larger than $\{u^*\}$, and higher-order terms select u^* among them.*

(4) The sign split. *At the valuation of a critical point that is not positive, the leading coefficients ρ_k of the surviving terms give a linear dependence $0 = \sum_K \rho_k v_k$ with every $\rho_k \neq 0$, real of either sign or complex. Unlike the positive case, this dependence does not define a probability law, and the leading terms there need not balance.*

Proof. (1) In log coordinates $b = \log y$, $\log W_T(e^b) = \log \sum_k T^{-\lambda_k} e^{\langle v_k, b\rangle}$ is strictly convex. Its Hessian is the covariance of the v_k under w, which is positive definite because the v_k *affinely* span $\mathbb{R}^n$. Indeed the v_k of a compact polytope positively span $\mathbb{R}^n$, so 0 lies in the interior of $\operatorname{conv}\{v_k\}$, which is therefore full-dimensional. Positive spanning also makes the function coercive: every direction increases some term. It therefore has exactly one critical point. Since $W_T > 0$ there, a critical point of $\log W_T$ is a critical point of W_T, and its gradient is the mean of w. (2) Write $T = \tau^N$, with N a common denominator of the λ_k. The positive critical point is a semialgebraic

function of τ, so it has a Puiseux expansion with positive leading coefficients [10], and its valuation exists. The weights w_k then have orders $\ell_k(u^*)$ in T, up to a common shift, so they concentrate on $K(u^*)$. The limits are strictly positive on every $k \in K(u^*)$, namely $w_k \to \rho_k / \sum_{j \in K(u^*)} \rho_j$, where the $\rho_k > 0$ are the leading coefficients of the positive Puiseux series $T^{-\lambda_k} y_T^{v_k}$. That strict positivity is what places 0 in the relative interior. This is the arc-limit theorem of the layers paper. The mean stays 0 in the limit; apply lemma 2. Judd and Rietsch prove that the valuation satisfies tropical critical conditions of this kind for complete positive Laurent polynomials over generalised Puiseux series [32]. (3) is example 6(b). (4) The same expansion, with complex leading coefficients ρ_k, gives $0 = \sum_K \rho_k v_k$ with every $\rho_k \neq 0$. Example 6(a) shows a real critical point with one negative coordinate, whose leading terms admit no positive balance. The checks below corroborate these statements; they are not proofs. KB4 checks (1)–(2) on $\mathbb{CP}^2$ and on F_1 at five blow-up sizes, from four starts each. KB5 checks (3) at exact rational points. KB6 finds all four critical points of F_1 in 80-digit arithmetic and checks (4). □

4 The Floer reading

For a compact toric Fano manifold X_Q, Cho and Oh show that the potential of the Lagrangian torus fibres is exactly W, with no corrections, and that a fibre $L(u)$ has non-vanishing Floer cohomology for some local system iff a critical point of W has valuation u [11]. Fukaya, Oh, Ohta and Ono extend the picture to all compact toric manifolds, where W acquires corrections [23]. We use only the Fano statement, and we prove nothing about Floer cohomology. Both examples below are Fano: $\mathbb{CP}^2$, and F_1, the blow-up of $\mathbb{CP}^2$ at a point, which is a del Pezzo surface. The Hirzebruch surfaces F_n with $n \geq 2$ are not Fano.

Corollary 4 (the probabilistic fibre)**.** *On a compact toric Fano manifold, exactly one critical point of W is positive. Its valuation determines* one *torus fibre $L(u^*)$ with non-vanishing Floer cohomology, not the only one. At u^*, the terms of W of least order form a conditioned law with mean exponent* 0*. Other critical points, including non-positive ones, may determine further fibres with non-vanishing Floer cohomology, but those are not represented by the positive probabilistic law. The fibre $L(u^*)$, a Lagrangian torus, is a different object from the positive real Lagrangian section of proposition 1(3), which meets it once.*

Proof. Combine theorem 3(1)–(2) with the cited theorem. □

Example 5 ($\mathbb{CP}^2$)**.** Here $v = (1,0), (0,1), (-1,-1)$ and $\lambda = (0,0,-1)$, so $W_T = y_1 + y_2 + T/(y_1 y_2)$. The critical points are $y_1 = y_2$ with $y_1^3 = T$: three of them, one positive, all with valuation $(\frac{1}{3}, \frac{1}{3})$, the barycentre. There all three terms tie, and $\frac{1}{3}\big((1,0) + (0,1) + (-1,-1)\big) = 0$. The barycentre is also the only point where the leading terms balance. Its fibre is the Clifford torus.

Example 6 (the Hirzebruch surface F_1)**.** Take $Q = \{p_1 \geq 0,\ p_2 \geq 0,\ p_1 + p_2 \leq 1,\ p_2 \leq 1 - \alpha\}$, the triangle of $\mathbb{CP}^2$ with the corner $(0,1)$ cut by a facet of length $\alpha \in (0,1)$. So $v = (1,0), (0,1), (-1,-1), (0,-1)$ and $W_T = y_1 + y_2 + T/(y_1 y_2) + T^{1-\alpha}/y_2$. The critical equations are $y_1^2 y_2 = T$ and $y_2^2 = T/y_1 + T^{1-\alpha}$. With $y_2 = w^2$ and $y_1 = T^{1/2}/w$, they become $w^4 - T^{1/2} w - T^{1-\alpha} = 0$, so there are four critical points. The Newton polygon of this quartic gives their valuations.

(a) $\alpha < \frac{1}{3}$. Three critical points lie over $(\frac{1}{3}, \frac{1}{3})$, one of them positive, and there the first three terms balance. The fourth lies over $(\alpha, 1 - 2\alpha)$. Its leading coefficient is $w \approx -T^{1/2-\alpha}$, so $y_1 < 0$. Its leading terms are $(1,0), (-1,-1), (0,-1)$. They admit the signed balance

$(1,0)+(-1,-1)-(0,-1)=0$ (in general $a_1=a_3=-a_4$), but no positive one. By corollary 4, the probabilistic fibre is over the barycentre, and the second fibre that the cited theorem provides is outside the positive theory.

(b) $\alpha>\frac{1}{3}$. All four critical points lie over $u^*=(\frac{1+\alpha}{4},\frac{1-\alpha}{2})$, one of them positive. But the leading terms $(0,1)$ and $(0,-1)$ balance at every point of the open segment $u_2=\frac{1-\alpha}{2}$, $\frac{1-\alpha}{2}<u_1<\alpha$. The endpoints are excluded. There a third term joins the leading set, which is $\{v_1,v_2,v_4\}$ at $u_1=\frac{1-\alpha}{2}$ and $\{v_2,v_3,v_4\}$ at $u_1=\alpha$. Neither set balances positively, because the first coordinates of its vectors are $1,0,0$, or $0,-1,0$. Leading order gives a segment of candidates, and the next order selects u^*: the same mechanism by which the layers paper's ties at the limit are broken at higher order.

At $\alpha=\frac{1}{3}$ all four facets tie at the barycentre. KB5 and KB6 corroborate every statement here at $\alpha=\frac{1}{5}$ and $\frac{3}{5}$; KB4 also uses $\frac{1}{4},\frac{2}{5},\frac{4}{5}$ (fig. 1).

5 The positive thimble and its tropical limit

The positive critical point has a second, analytic role: it is the critical point of an oscillatory integral. In mirror symmetry the integral $\int e^{-W/\hbar}\,dy/y$ over the positive real cycle is the one tied to Galkin's conifold point and to Gamma conjecture I; under homological mirror symmetry that cycle is expected to correspond to the structure sheaf [25, 26, 31]. Here we only need its elementary properties, and what the tropical limit does to the law it carries. Give $(\mathbb{C}^*)^n$ the flat metric $\sum_j|d\log y_j|^2$, and let $\hbar>0$.

Proposition 7 (the positive thimble)**.** *(1) The positive orthant $(0,\infty)^n$ is the ascending Lefschetz thimble J_+ of y_T for $\operatorname{Re}(W_T/\hbar)$, the set of points whose downward gradient flow tends to y_T. In $b=\log y$, the Hessian of $\operatorname{Re}W_T(e^b)$ at $b_T=\log y_T$, in the coordinates $(\operatorname{Re}b,\operatorname{Im}b)$, is $\operatorname{diag}(H,-H)$ with $H=\operatorname{Hess}(W_T\circ\exp)(b_T)$ positive definite. So the real directions ascend, and $\operatorname{Im}W_T=0$ on J_+.*

(2) $\mu_\hbar=Z^{-1}e^{-W_T(y)/\hbar}\,dy/y$ on J_+ is a probability law, and $Z<\infty$.

(3) Laplace. *As $\hbar/W_T(y_T)\to 0$, the law of $b=\log y$ under $\mu_\hbar$ is asymptotically normal, with mean b_T and covariance $\hbar H^{-1}$.*

Proof. (1) $W_T\circ\exp(b)=\sum_k T^{-\lambda_k}e^{\langle v_k,b\rangle}$ is holomorphic, and its restriction to real b is strictly convex and coercive (theorem 3(1)). For a holomorphic f, the Hessian of $\operatorname{Re}f$ in $(\operatorname{Re}b,\operatorname{Im}b)$ is $\operatorname{diag}(H,-H)$ with H the real Hessian, by the Cauchy–Riemann equations. The metric is real, so the gradient flow of $\operatorname{Re}W_T$ preserves the real locus. There the flow is the gradient flow of a strictly convex, coercive function, and every orbit tends to b_T. So the orthant lies in the stable manifold J_+, which is an n-ball. The orthant is an n-manifold, so it is open in J_+; it is closed in $(\mathbb{C}^*)^n$, so it is closed in J_+. Since J_+ is connected, the two are equal. (2) $W_T>0$ on J_+, and $W_T(e^b)\geq c|b|-C$ by positive spanning, so $Z<\infty$. (3) This is Laplace's method at the non-degenerate minimum b_T of $W_T\circ\exp$; the relative corrections are $O(\hbar/W_T(y_T))$. KB7 corroborates (1) and (3), at $\hbar/W_T(y_T)=10^{-2},10^{-3},10^{-4}$. □

Laplace describes the limit $\hbar\to 0$ at fixed T. The tropical limit couples the two, $\hbar=T^h$, and reads y on the scale of the valuation, $u=\log y/\log T$. Since $T^{-\lambda_k}y^{v_k}=T^{\ell_k(u)}$, the pushforward




Figure 1: F_1 at two blow-up sizes. (a) $\alpha = 0.2$: the only point where the leading terms balance is the barycentre, which carries the positive critical point; a real critical point with $y_1 < 0$ sits over $(\alpha, 1 - 2\alpha)$, outside the positive theory. (b) $\alpha = 0.6$: the leading terms balance along an open segment, whose endpoints (open circles) have no positive balance, and the critical points all sit over one point of it. (c) $\alpha = 0.6$: the u_1-marginal of the thimble law with $\hbar = T^h$ (theorem 8 and example 9). At $h = 0$ it is the marginal of the uniform law on Q; at $h = h^* = 0.2$ it is uniform on the segment; at $h = 0.3$ it is uniform on $[0.3, 0.5]$; above 0.4 it is concentrated at $u_1^* = 0.4$. Here $\log T = -200$, and -100 for $h = 0.45$.

$\nu_{T,h}$ of μ_{T^h} to u has density proportional to

$$f_{T,h}(u) = \exp\Big(-\sum_k T^{\ell_k(u)-h}\Big).$$

Let $g(u) = \min_k \ell_k(u)$, which is the tropicalisation of W in the min convention. Let $h^* = \max_u g$, and let $Q_h = \{u : g(u) \geq h\} = \{\langle v_k, u\rangle \geq \lambda_k + h\}$, so that $Q_0 = Q$.

Theorem 8 (the tropical limit of the thimble law)**.** *(1) If $h < h^*$, then $\nu_{T,h}$ tends, in total variation, to the uniform law on the polytope Q_h. At $h = 0$ this is the normalised Duistermaat–Heckman measure of X_Q.*

(2) If $h \geq h^*$, then $\nu_{T,h}$ concentrates on $Q_{h^*} = \operatorname{argmax} g$: $\nu_{T,h}(U) \to 1$ for every open $U \supseteq Q_{h^*}$.

(3) $\operatorname{argmax} g = \{u : 0 \in \operatorname{conv}\{v_k : k \in K(u)\}\}$. *It is a polytope. It contains every candidate, every u with $0 \in \operatorname{relint}\operatorname{conv}\{v_k : k \in K(u)\}$, and in particular u^*; it can also contain non-candidates. Every u outside it has $g(u) < h^*$. This applies to the valuation of a signed critical point of example 6(a) as well.*

Proof. Write $L = \log(1/T)$. Since the v_k positively span, $g(u) \leq C - c|u|$, so h^* is finite and attained, and the level sets of g are compact. (1) For u with $g(u) > h$, every exponent is positive and $f_{T,h}(u) \to 1$. Off Q_h, some exponent is negative and $f_{T,h}(u) \to 0$, and ∂Q_h is null. For $T \leq T_0 < 1$, $f_{T,h} \leq \min(1, \exp(-T_0^{g-h}))$, which is integrable. By dominated convergence, $\int f_{T,h} \to \operatorname{vol} Q_h > 0$, since Q_h has interior when $h < h^*$. Scheffé's lemma then gives convergence in total variation. (2) Choose $\delta > 0$ with $\{g \geq h^* - 2\delta\} \subseteq U$. Put $s = h - h^* \geq 0$. Off U, $T^{g-h} \geq T^{-(s+2\delta)}$, so $\int_{U^c} f_{T,h} \leq C\exp(-\frac{1}{2}T^{-(s+2\delta)})$. Near a maximiser u_0, on the ball of radius $1/(ML)$, where $M = \max|v_k|$, every $\ell_k \geq h^* - 1/L$. So $f_{T,h} \geq \exp(-me\,T^{-s})$ there, and the mass of the ball is at least $cL^{-n}\exp(-me\,T^{-s})$. The ratio of the two bounds tends to 0, because $T^{-2\delta} \to \infty$. (3) g is concave and piecewise linear, and its superdifferential at u is $\operatorname{conv}\{v_k : k \in K(u)\}$ [36, Theorem 23.8]. A point maximises a concave function iff 0 lies in its superdifferential. Candidates satisfy this, and so does u^* by theorem 3(2). This recovers, in our setting, the theorem of Judd and Rietsch that the tropical critical point maximises the tropicalisation [32, Thm. 1.2]. KB8 corroborates (1)–(3) on $\mathbb{CP}^2$ and on F_1 with $\alpha = \frac{1}{5}$ and $\frac{3}{5}$. □

So the first level of the tropical limit sees exactly the max–min set. That set is the first step of the iterated max–min by which Fukaya, Oh, Ohta and Ono locate a fibre [23], and of the recursive construction of Judd and Rietsch [32]. The selection of u^* inside it is a second-level question.

Example 9 (the levels on F_1)**.** For $\mathbb{CP}^2$, and for F_1 with $\alpha < \frac{1}{3}$, $h^* = \frac{1}{3}$ and $\operatorname{argmax} g$ is the barycentre. The signed fibre of example 6(a) has $g = \alpha < h^*$. For F_1 with $\alpha > \frac{1}{3}$, $h^* = \frac{1-\alpha}{2}$, and $\operatorname{argmax} g$ is the closed segment $S = \{u_2 = \frac{1-\alpha}{2},\ \frac{1-\alpha}{2} \leq u_1 \leq \alpha\}$. Its relative interior consists of the candidates, and its endpoints are maximisers that are not candidates.

(a) At $h = h^*$ the u_1-marginal of $\nu_{T,h}$ tends to the *uniform* law on $[\frac{1-\alpha}{2}, \alpha]$. Leading order gives the selected point no extra mass.

(b) On S, $\ell_1 = u_1$ and $\ell_3 = \frac{1+\alpha}{2} - u_1$. The next max–min level is $h_2 = \max_S \min(\ell_1, \ell_3) = \frac{1+\alpha}{4}$, attained at $u_1 = \frac{1+\alpha}{4} = u_1^*$. For $h^* < h < h_2$, the u_1-marginal tends to the uniform law on $[h, \frac{1+\alpha}{2} - h]$. For $h > h_2$, it concentrates at u_1^*.

Sketch. Near S, the transverse factor $\exp(-T^{u_2-h} - T^{1-\alpha-u_2-h})$ does not depend on u_1. The coupling of u_2 to ℓ_3 matters only in boundary layers of width $O(1/L)$. So the u_1-marginal behaves as in theorem 8, now for $\min(\ell_1, \ell_3)$ on S. KB8 corroborates (a) and (b) for $\alpha = \frac{3}{5}$, at $h = \frac{1}{5}, \frac{3}{10}, \frac{9}{20}$ (fig. 1(c)). That the mean at level h^* equals u_1^* here is a symmetry of this example: u_1^* is the midpoint of S. The mechanism of selection is the second level, not the mean.

6 Soft evidence moves λ

This section starts the programme of conditioning existing structures on data and recording what changes. It treats one kind of data, soft evidence whose strength grows with the amount of data. Hard evidence (zero cells) and forgetting are not treated here.

Definition 10 (the data action)**.** Fix a reference scale N and put $T = e^{-N}$. Evidence that multiplies term k of W_T by $e^{-n\rho_k}$ after n observations, with rates $\rho_k \geq 0$, multiplies it by $T^{(n/N)\rho_k}$. So it replaces λ by

$$\lambda^{\text{eff}} = \lambda - s\rho, \qquad s = n/N.$$

Which term each observation weighs, and the scale N, are modelling choices. A likelihood of fixed size has $s \to 0$ as $N \to \infty$. It changes the coefficients smoothly, as in proposition 1, and changes no valuation.

Let Δ_m be the laws on the m facets, and let $B = \{q \in \Delta_m : \sum_k q_k v_k = 0\}$ be the *balance polytope*, the laws with mean exponent 0. It has dimension $m - n - 1$, because 0 lies in the interior of $\operatorname{conv}\{v_k\}$. Write $g_\lambda = \min_k \ell_k$, $h^*(\lambda) = \max g_\lambda$, and $F(\lambda)$ for the face of B on which $\langle q, \lambda\rangle$ is maximal.

Proposition 11 (the max–min level is a balance programme)**.** *(1) $h^*(\lambda) = \min_{q\in B}\big(-\langle q, \lambda\rangle\big)$. So h^* is concave and piecewise linear in λ.*

(2) Let K_0 be the union of the supports of the laws in $F(\lambda)$, and $\sigma_0 = \operatorname{conv}\{v_k : k \in K_0\}$. Then $K(u) = K_0$ on the relative interior of $\operatorname{argmax} g_\lambda$, and $K(u) \supsetneq K_0$ on its relative boundary. The candidates (theorem 8(3)) are exactly the relative interior, and $\dim \operatorname{argmax} g_\lambda = n - \dim \sigma_0$.

(3) The limit law of the positive critical point (theorem 3(2)) lies in $F(\lambda)$.

Proof. (1) $\max_u \min_k \ell_k = \max_u \min_{q\in\Delta_m} \sum_k q_k \ell_k(u)$, and the minimax theorem exchanges the two. For fixed q, $\sup_u \sum_k q_k(\langle v_k, u\rangle - \lambda_k)$ is $-\langle q, \lambda\rangle$ if $\sum_k q_k v_k = 0$ and $+\infty$ otherwise. (2) By complementary slackness, $q_k > 0$ for an optimal q forces $\ell_k(u) = h^*$ at every maximiser u, so $K(u) \supseteq K_0$ on $\operatorname{argmax} g_\lambda$. By strict complementarity [37], some maximiser has $\ell_j > h^*$ for every $j \notin K_0$. The same then holds on the relative interior of $\operatorname{argmax} g_\lambda$, which is $\{u : \ell_k(u) = h^*\ (k \in K_0),\ \ell_j(u) > h^*\ (j \notin K_0)\}$. A law in the relative interior of $F(\lambda)$ is positive on K_0 and has mean 0, so those points are candidates (lemma 2). At a boundary point, suppose 0 were in the relative interior of $\operatorname{conv}\{v_k : k \in K(u)\}$. Then there would be a balanced law positive on all of $K(u)$. That law would be optimal, since $\sum_k q_k \ell_k(u) = h^*$, so its support would lie in K_0, a contradiction. The linear span of $\{v_k : k \in K_0\}$ is its affine span, since it contains 0, so the equations $\ell_k = h^*$ on K_0 cut out dimension $n - \dim \sigma_0$. (3) The limit law q is balanced and supported on $K(u^*)$, where every $\ell_k(u^*) = h^*$. So $-\langle q, \lambda\rangle = \sum_k q_k \ell_k(u^*) = h^*$, and q is optimal. DW2 and DW3 corroborate (1)–(3) on 40 and 23 configurations. □

So candidates against selection appears twice, in dual spaces. In u-space, u^* is selected inside the relative interior of $\operatorname{argmax} g_\lambda$, which has dimension $n - \dim \sigma_0$. In law space, the limit law is selected inside $F(\lambda)$. The two can be non-trivial at different parameters (example 14).

Corollary 12 (where data can act)**.** *(1) The leading set K_0, the candidate dimension and the face F change only on the walls of the normal fan of B, the* outcome walls*. Within a chamber, h^* is affine in λ, and so is u^* when σ_0 is an n-simplex.*

(2) Evidence along a translation, $\rho_k = \langle v_k, t\rangle$, changes nothing but $u^* \mapsto u^* - st$. Evidence along the constant, $\rho = \mathbf{1}$, changes nothing but $h^* \mapsto h^* + s$. Only the quotient of $\mathbb{R}^m$ by these $n+1$ directions can cross a wall. It has dimension $m - n - 1$, the number of independent affine dependences (circuits) of $\{v_k\}$.

(3) Suppose there is one circuit a, normalised to be primitive and integral. The wall is $c(\lambda) = -\langle a, \lambda\rangle = 0$, and along $\lambda_0 - s\rho$ it is reached at $s^* = -c(\lambda_0)/\langle a, \rho\rangle$ when this is positive, never otherwise. Over rates with $\max_k \rho_k - \min_k \rho_k \le 1$, the least such s is $|c(\lambda_0)|/(\frac{1}{2}\|a\|_1)$.

Proof. (1) $F(\lambda)$ is constant on the relative interior of each cone of the normal fan of B; apply proposition 11. On a chamber, $h^* = -\langle q_0, \lambda\rangle$ for a fixed vertex q_0. When σ_0 is an n-simplex, u^* is the unique solution of $\ell_k(u) = h^*$, $k \in K_0$, which is linear in λ. (2) Under a translation, ℓ_k becomes $\ell_k(u + st)$. Under the constant, every ℓ_k shifts by s. Both are orthogonal to $B - B$, which consists of affine dependences. (3) B is a segment, and its normal fan has one wall, $\langle q_A - q_B, \lambda\rangle = 0$, where $q_A - q_B$ is a multiple of a. Since $\sum_k a_k = 0$, adding a constant to ρ leaves $\langle a, \rho\rangle$ unchanged. So the best value over range at most 1 is $\sum_k a_k^+ = \frac{1}{2}\|a\|_1$. □

Example 13 ($\mathbb{CP}^2$: no wall)**.** Here $m = n + 1$, so there is no circuit, and B is the single law $(\frac{1}{3}, \frac{1}{3}, \frac{1}{3})$. No soft evidence changes the leading set of $\mathbb{CP}^2$. Data only move the selected point and the level (DW4, DW7).

Example 14 (F_1 across its wall)**.** For general λ, put $L = -(\lambda_1 + \lambda_2 + \lambda_3)$ and $\alpha = -\lambda_1 - \lambda_3 + \lambda_4$. These are the size of the triangle and the length of the blown-up facet, and both are invariant under translation. The Kähler cone is $0 < \alpha < L$. Its walls are viability walls, where the polytope loses facet 4 or collapses, and X_Q changes. The one circuit is $a = (2, -1, 2, -3)$, with $c(\lambda) = 3\alpha - L$. The balance polytope is the segment from $q_A = (\frac{1}{3}, \frac{1}{3}, \frac{1}{3}, 0)$ to $q_B = (0, \frac{1}{2}, 0, \frac{1}{2})$, and $h^* = \min\big(L/3, (L - \alpha)/2\big)$.

(a) The outcome wall $\alpha = L/3$ lies *inside* the Kähler cone: the manifold does not change there. Below it, $K_0 = \{1, 2, 3\}$ and the candidate is a point. Above it, $K_0 = \{2, 4\}$, the candidates form an open segment, and u^* is selected at the second level (example 9). The lower diagonal of the subdivision flips from v_1v_3 to v_2v_4.

(b) u^* is continuous and piecewise affine across the wall (fig. 2(a)). What jumps is the leading set, the candidate dimension ($0 \to 1$) and the mechanism of selection (leading order $\to$ second level). The signed fibre over $(\alpha, 1 - 2\alpha)$ reaches the barycentre at the wall. By the cited theorem [11], the number of torus fibres with non-vanishing Floer cohomology drops from two to one there. We claim nothing about the Fukaya category.

(c) *At the wall* ($\alpha = 1/3$, $L = 1$), the substitution $w = T^{1/6}x$ turns the quartic of example 6 into $x^4 - x - 1 = 0$ for every T. So the positive law is $(1/x, x^2, 1/x, 1/x^2)/\Sigma \approx (0.2156, 0.3922, 0.2156, 0.1766)$ for every T, where $x \approx 1.2207$. It is the interior point $(1 - t)q_A + tq_B$ of $F = B$, with $t \approx 0.353$. Here the candidate in u-space is a single point, but the law is selected inside a segment.

(d) The discriminant of $w^4 - T^{1/2}w - T^{1-\alpha}$ is $-27T^2 - 256T^{3-3\alpha}$, which never vanishes: no two critical points collide for $T > 0$. The outcome wall is where its two terms have equal order. It is a wall of the tropical discriminant, not of the discriminant itself.

(e) *Data.* Start at $\alpha = 1/5$, $L = 1$, so $c = -2/5$. Evidence against term 1, $\rho = e_1$, has $\langle a, \rho\rangle = 2$ and reaches the wall at $s^* = 1/5$, that is $n^* = N/5$ (fig. 2(b)). Along the way $\alpha/L = (1/5+s)/(1+s)$ stays in the Kähler cone. Evidence against term 4 ($\langle a, \rho\rangle = -3$) never reaches the wall. The most efficient rates with range at most 1 are $\rho = (1, 0, 1, 0)$, against terms 1 and 3, with $s = 1/10$.

DW4–DW7 corroborate (a)–(e).




Figure 2: Soft evidence on F_1. (a) The selected valuation u^* across the Kähler cone. The curves are the chamber formulas and the circles the positive critical point at $T = 10^{-30}$. The outcome wall at $\alpha = 1/3$ changes the leading set and the candidate dimension, not the position of u^*; here $h^* = u_2^*$. (b) A data path from $\alpha = 1/5$ with evidence against term 1: $u^*(s)$ is affine on each side of $s^* = 1/5$, with a kink at the wall.

On F_1, then, the least data that changes the perspective changes *which terms carry the limit law*, not where the selected state sits. A decision that reads the leading set jumps at n^*. A decision that reads u^* continuously does not. Decision, contextuality and value-of-information walls need their own definitions, a decision model, a named complex and obstruction class, and an information-acquisition model, before they can be compared with the outcome walls. We record them separately and have not computed them.

Part II

Back to contingency tables

Part I went from toric geometry to conditioned probability. Part II runs the dictionary backwards, from a table of counts to the potential, and asks what the Floer-selected fibre means for the table. It works the independence model exactly. Every operation it uses, and every operation it does not yet treat, is recorded in the ledger (appendix A).

7 Two configurations of one model

An $r \times c$ table of counts t_{ij} has row margins t_{i+} and column margins t_{+j}. The independence model is the log-linear model whose design matrix A has one column per cell, the row indicator stacked on the column indicator. It is toric [17, 27]: its sufficient statistic is the vector of margins, and by Birch the MLE is the unique law in the model with the observed margins, $\hat{p}_{ij} = t_{i+}t_{+j}/n^2$ [8, 9]. The moment polytope of this model is $\Delta_{r-1} \times \Delta_{c-1}$, so its manifold is $\mathbb{CP}^{r-1} \times \mathbb{CP}^{c-1}$, which is Delzant and Fano.

The model carries two different point configurations, and they must not be confused.

Proposition 15 (cells and facets)**.** *(1)* Cells. *The rc columns of A satisfy $(r-1)(c-1)$ independent linear relations. These are spanned by the basic moves $+1$ at $(i,j),(k,l)$ and -1 at $(i,l),(k,j)$, which change a table without changing its margins. This is the Tier A support of proposition 1.*

(2) Facets. *The $r+c$ inward facet normals of $\Delta_{r-1}(L_1) \times \Delta_{c-1}(L_2)$ have exactly one circuit, $a = (c\,\mathbf{1}_r, -r\,\mathbf{1}_c)/\gcd(r,c)$. This is the configuration W is built from.*

Proof. (1) A has rank $r+c-1$, and the moves lie in its kernel and have rank $(r-1)(c-1)$ [17]. (2) $m - n - 1 = (r+c) - (r+c-2) - 1 = 1$. The row normals sum to 0, and so do the column normals. So $\sum_k a_k v_k = 0$, and $\sum_k a_k = rc - cr = 0$. CT1 checks both for $r, c \in \{2,3,4\}$. □

So the 2×3 model has two moves but one circuit. The moves act on tables with fixed margins, within a fibre. The circuit acts on λ, across Kähler classes. The second statement is a special case of a general count.

Proposition 16 (products add one circuit)**.** *If $v^{(1)}$ and $v^{(2)}$ both have 0 in the interior of their convex hulls, the product configuration has $(m_1 - n_1 - 1) + (m_2 - n_2 - 1) + 1$ independent circuits. The extra one is $(q^{(1)}, -q^{(2)})$ for balanced laws $q^{(i)}$ of the factors. Its wall is $h_1^* = h_2^*$ when each factor has a single balanced law.*

Proof. The count is $(m_1+m_2)-(n_1+n_2)-1$. The vector $(q^{(1)}, -q^{(2)})$ sums to $1-1=0$, and both halves balance. It is independent of the circuits of the factors, which sum to zero on each factor separately. When $B_i = \{q^{(i)}\}$, the balance programme gives $h^* = \min(h_1^*, h_2^*)$ (proposition 11), with a wall where they are equal. □

8 The independence model: its wall and its selection

In the coordinates $b =$ the first $r-1$ row margins and the first $c-1$ column margins, the facets of $Q = \Delta_{r-1}(L_1) \times \Delta_{c-1}(L_2)$ have $\ell_k(b)$ equal to the k-th margin. Here L_1 and L_2 are the totals of the row and column margins. So $W = W_{\text{row}}(y) + W_{\text{col}}(z)$ with $W_{\text{row}} = y_1 + \cdots + y_{r-1} + T^{L_1}/(y_1 \cdots y_{r-1})$, and similarly for columns.

Theorem 17 (the independence model)**.** *(1) $h^* = \min(L_1/r,\, L_2/c)$. The leading set is the row facets if $cL_1 < rL_2$, the column facets if $cL_1 > rL_2$, and all $r+c$ facets on the wall $cL_1 = rL_2$. The wall is $c(\lambda) = -\langle a, \lambda\rangle = (cL_1 - rL_2)/\gcd(r,c) = 0$.*

(2) The critical points of W are $y_i = \zeta T^{L_1/r}$, $z_j = \eta T^{L_2/c}$ with $\zeta^r = \eta^c = 1$, so there are rc of them. All have the valuation $u^* = (L_1/r, \ldots; L_2/c, \ldots)$, the uniform margins, for every L_1, L_2. The positive law is exactly proportional to $T^{L_1/r}$ on each row facet and $T^{L_2/c}$ on each column facet.

(3) *Off the wall the candidates have dimension* $c-1$ *(if* $cL_1 < rL_2$*) or* $r-1$*, and the uniform margins are selected at the second level. On the wall the candidate is the single point* u^**.*

(4) *By the cited theorem [11], exactly one torus fibre has non-vanishing Floer cohomology, for every* L_1, L_2*. So the outcome wall of the independence model changes no Floer data. On* F_1 *it does (example 14(b)).*

Proof. (1) Each factor has the single balanced law, uniform on its facets, with values L_1/r and L_2/c. Apply proposition 16. (2) The variables separate. $\partial W_{\text{row}}/\partial y_i = 0$ gives $y_i \prod_j y_j = T^{L_1}$ for every i, so all y_i are equal and $y^r = T^{L_1}$. The term values are then $T^{L_1/r}$ times a root of unity. (3) Off the wall the leading set is one factor's facets, so σ_0 has dimension $r-1$ (or $c-1$); apply proposition 11(2). (4) All critical points share one valuation. CT2 and CT3 check (1)–(3), with exact rationals and gradients. □

9 Margins against evidence

Data enter in two ways, and the note keeps them apart.

Proposition 18 (two roles of data)**.** *(1)* Observed margins b *pick a point of* Q*, the fibre* $\mu^{-1}(b)$ *and the MLE (Birch). For the independence model,* $\ell_k(b)$ *is exactly the* k*-th margin, the facet* $\{\ell_k = 0\}$ *is a zero margin, and the MLE lies in the open model iff every margin is positive [9]. For general hierarchical models the facets correspond to facial sets of cells, and the MLE fails to exist exactly when the margins lie on a proper face [20, 21]. We use only the independence case.*

(2) Evidence on the terms of W *moves* λ *(definition 10), and with it* u^**,* h^* *and the walls. With* λ *fixed,* u^* *does not depend on* b*.*

(3) *Both* b *and* u^* *are points of* Q*, in moment coordinates. The valuation of a critical point is the moment coordinate of its torus fibre* $L(u^*) = \mu^{-1}(u^*)$ *[11]. So the MLE fibre and the Floer-selected fibre can be compared in one space. For the independence model we use the Fisher–Rao distance within the model, between* $\hat{p}$ *and the law at* u^* *(uniform margins). With surveys of sizes* n_1 *for rows and* n_2 *for columns,*

$$d(\hat{p}, p^*)^2 = n_1\, d_\Delta(\hat{p}_{\text{row}}, \mathbf{1}/r)^2 + n_2\, d_\Delta(\hat{p}_{\text{col}}, \mathbf{1}/c)^2, \qquad d_\Delta(p,q) = 2\arccos \sum_i \sqrt{p_i q_i}.$$

Joint sampling of n *cells is the case* $n_1 = n_2 = n$*. A coarser index is the* centrality $g(b)/h^*$*: the smallest margin divided by its largest possible value.*

Proof. (1) The first statements are direct, and the MLE formula shows positivity. (2) is corollary 12 together with the fact that W does not involve b. (3) The Fisher information of the independence family is block diagonal, $n_1 \operatorname{Cov}(\text{row}) \oplus n_2 \operatorname{Cov}(\text{col})$, because the log-likelihood separates into a row part and a column part. So the model is a Riemannian product of two scaled simplices, and on each simplex the Fisher–Rao distance is $2\arccos$ of the Bhattacharyya coefficient [2]. CT4 and CT5 check (1) and the block structure. □

The MLE fibre and the Floer-selected fibre coincide exactly when the margins are uniform. That the Floer fibre is the uniform-margin table, the one farthest from every zero-margin degeneracy in lattice distance, is a reading of the cited theorem, not a new statistical result. The polytope used here is the *base*, the space of margins. Section 11 turns to the *fibre* over a point of it, the coupling polytope, where the same margins become the Kähler class.

10 Data and walls in the independence model

Example 19 (2×2 and 2×3)**.** Under joint sampling, $L_1 = L_2 = n$. The 2×2 table therefore sits on its wall for every n, which is the monotone $\mathbb{CP}^1 \times \mathbb{CP}^1$. For 2×3, $cL_1 - rL_2 = n > 0$, so $h^* = n/3$, the column facets lead, and the candidates are the open segment of row margins $b_1 \in (n/3, 2n/3)$ with uniform column margins. Its endpoints are maximisers but not candidates. The selected point is $b_1 = n/2$. To reach the wall the column survey must grow to $n_2 = 3n/2$: $n/2$ more column observations. In the terms of corollary 12(3), this is ρ on a column facet with $\langle a, \rho \rangle = -r/\gcd$, so $s^* = n/2$. Crossing it changes the leading set and the candidate dimension. By theorem 17(4) it changes no Floer data. CT6 checks every number.

Example 20 (hard evidence: a structural zero)**.** Declare cell $(2, 2)$ of the 2×2 table structurally zero. This is the quasi-independence model on three cells. The three remaining cells span a unimodular triangle, which is *not* a face of the square: hard evidence need not restrict to a toric divisor. Its facet configuration is that of $\mathbb{CP}^2$, which has no circuit, so the one wall of the 2×2 model is gone. CT7 checks this. The general case, deleting cells of larger tables, is open.

11 One coupling polytope through every layer

What this section shows, in plain words. Fix two margins, a row law f and a column law g. The tables (couplings) with those margins form a polygon, the *fibre*. That polygon is the picture of a toric surface, and everything in Part I can be drawn on it.

- The usual conditional law of a table given its margins, the one behind Fisher's exact test, *is* the toric Kähler structure (fig. 4).
- The positive critical point of W picks the *leximin* coupling: the one whose smallest cell is as large as possible (fig. 3).
- The positive thimble is a family of laws on couplings. It runs from "every coupling equally likely" to "only the leximin coupling" (fig. 5).
- Two kinds of wall cut the space of margins. Blow-up walls change the surface; leximin walls change which cells lead. Here one count of data crosses a leximin wall and changes the Floer fibres, while the surface stays the same (fig. 6).

The example is the one followed in the architecture note [6]: $f = (2/5, 3/5)$, $g = (3/10, 43/100, 27/100)$.

11.1 Base and fibre

A pair of margins is a point of the *base*, $\Delta_{n-1} \times \Delta_{m-1}$ (section 8). Over it sits the *fibre*, the coupling polytope $\mathrm{Cpl}(f, g)$ of tables with those margins. For nondegenerate margins the fibre is Delzant, its facets are cells $\{\alpha_c = 0\}$, and the Kähler class of its toric manifold is the product coupling, $[\omega_{f,g}] = \sum_c f_i g_j [D_c]$ [5]. So the same data play two roles:

	base (sections 7 to 9)	fibre (this section)
points	margins (f, g)	couplings α with those margins
facets	a zero margin	a zero cell
the margins are	a point (the Birch/MLE fibre)	the Kähler class: λ_c is read off (f, g)
W is a sum over	margin facets	cell facets
Tier A support	cells, with $(r-1)(c-1)$ moves	integer tables with these margins

For 2×3, write a coupling by $x = \alpha_{11}$, $y = \alpha_{12}$. Then $\alpha_{13} = f_1 - x - y$, $\alpha_{21} = g_1 - x$, $\alpha_{22} = g_2 - y$ and $\alpha_{23} = x + y - (f_1 + g_1 + g_2 - 1)$, so cell c has $\ell_c = \langle v_c, (x, y) \rangle - \lambda_c$ with $v_c \in \{(1,0), (0,1), (-1,-1), (-1,0), (0,-1), (1,1)\}$.

Example 21 (the fibre)**.** For $f = (2/5, 3/5)$, $g = (3/10, 43/100, 27/100)$ the fibre is the pentagon with vertices $(13/100, 0)$, $(3/10, 0)$, $(3/10, 1/10)$, $(0, 2/5)$, $(0, 13/100)$ and area $1331/20000$, so $X = \mathrm{Bl}_2\mathbb{CP}^2$. Its facets are the cells $11, 12, 13, 21, 23$. Cell 22 never vanishes, since $\alpha_{22} \geq f_2 + g_2 - 1 = 3/100$. Scaled by $n = 100$, its lattice points are the 715 integer tables with margins $(40, 60)$ and $(30, 43, 27)$. By Pick's theorem, $715 = 665.5 + 97/2 + 1$. CF1 and CF2 check this exactly.




Figure 3: The example at a glance. (a) The leximin coupling: row 1 is the scarcest per cell, so it is spread evenly, and every other cell is larger. (b) The fibre, with the 715 tables as dots. Each side is a cell that can vanish. Marked: the product coupling, selected by Tier A; the leximin coupling, selected by the positive critical point of W; and the two real critical points with a negative coordinate (signed fibres).

11.2 Tier A on the fibre is Fisher's noncentral hypergeometric family

Proposition 22 (the conditional law of a table)**.** *Let S be the integer tables with margins Nf, Ng, written by (t_{11}, t_{12}), with weights $c_t = 1/\prod_{ij} t_{ij}!$.*

(1) For any cell parameters $\theta_{ij} > 0$, the law of a table under independent Poisson (or multinomial) sampling with means $\propto \theta_{ij}$, conditioned on its margins, is the monomial law $q(u) \propto c_t\, e^{\langle u, (t_{11}, t_{12}) \rangle}$. Here $u_1 = \log \frac{\theta_{11}\theta_{23}}{\theta_{13}\theta_{21}}$ and $u_2 = \log \frac{\theta_{12}\theta_{23}}{\theta_{13}\theta_{22}}$ are the two log odds ratios. This is Fisher's noncentral hypergeometric family. At $u = 0$ it is the central hypergeometric law, the null law of Fisher's exact test [22].

(2) By proposition 1, $\varphi(u) = \log \sum_t c_t e^{\langle u, t \rangle}$ is a Kähler potential. The moment map is the expected table, and the Kähler metric is the Fisher information of the odds ratios. Since S is the set of lattice points of the Delzant lattice polygon $N \cdot \mathrm{Cpl}(f, g)$, the form extends to all of $X = \mathrm{Bl}_2\mathbb{CP}^2$.

(3) At $u = 0$ the mean is exactly $N\, f \otimes g$: Tier A at odds ratios 1 selects the product coupling.

(4) Every interior coupling is the mean for exactly one u (Birch). In the example, the leximin coupling needs odds ratios $(e^{u_1}, e^{u_2}) \approx (0.82, 0.46)$.

Proof. (1) $\prod_{ij} \theta_{ij}^{t_{ij}}/t_{ij}!$ restricted to fixed margins depends on t only through c_t and $t_{11}u_1 + t_{12}u_2$. Substituting $t_{13}, t_{21}, t_{22}, t_{23}$ from the margins gives the two log odds ratios. (2) is proposition 1 and its compact case. (3) The central hypergeometric mean is $t_{i+}t_{+j}/N$ [9]. (4) is proposition 1(3). CF3 checks (1) and (3), the latter in exact rationals, and computes (4). □




Figure 4: Tier A. (a) The moment map sends the plane of log odds ratios onto the open pentagon. Blue lines hold u_1 fixed and orange lines hold u_2 fixed. (b) Two laws on the 715 tables: the hypergeometric law (odds ratios 1, centred on $f \otimes g$), and the law with odds ratios $(0.82, 0.46)$, centred on the leximin coupling.

11.3 Tier B: the selected coupling is the leximin one

Theorem 23 (the max–min coupling)**.** *Let (f, g) be nondegenerate $n \times m$ margins, and let W be the potential of* $\mathrm{Cpl}(f, g)$*, a sum over its facet cells.*

(1) $h^* = \max_\alpha \min_{c \text{ facet}} \alpha_c = \min\bigl(\min_i f_i/m,\ \min_j g_j/n\bigr)$.

(2) Every cell of a binding *row or column, one attaining the minimum, is a facet. If exactly one line binds, the leading set K_0 is that line's cells.*

(3) The candidate set has dimension $(n-2)(m-1)$ for a binding row and $(n-1)(m-2)$ for a binding column. For 2×3, a binding row gives a single candidate, so leading order selects. A binding column gives a segment, and the next level selects.

(4) The positive critical point of W sits over the facet-leximin coupling, the lexicographic max–min of the facet cells. This is corollary 29, which follows from the selection theorem (theorem 27).

Proof. (1) A row of m cells sums to f_i, so its smallest cell is at most f_i/m; columns are the same. Conversely, put h (the claimed value) in every cell and add the product of the residual margins

$f_i - mh \geq 0$ and $g_j - nh \geq 0$, whose totals agree. This gives a coupling with every cell at least h. (2) A cell with $\alpha_c > 0$ on all of Cpl has $f_i + g_j > 1$. For a binding row i that would give $f_i > \sum_{k \neq j} g_k \geq (m-1) \min_k g_k$ and $f_i \leq (m/n) \min_k g_k$, so $n(m-1) < m$, which is impossible. If row i alone binds, the residual margins of the other rows and of all columns are positive, so the coupling of (1) has every other cell strictly above h^*. The row's cells, which average h^*, equal it at every maximiser. (3) The m cell functionals of a row sum to the constant f_i, so their normals span $m-1$ dimensions and contain 0 in the relative interior of their hull. Apply proposition 11(2) in dimension $(n-1)(m-1)$. (4) See corollary 29. CF6 checks it on 20 random rational margins at $T = 10^{-60}$. CF4 checks (1)–(3) on about 1000 random margins of shapes 2×3, 3×3, 2×4 and 3×4, against the facet-only programme. □

In words: *the selected coupling spreads the scarcest row or column evenly over its cells.* In the example, row 1 binds with $f_1/3 = 2/15$. So $h^* = 2/15$, $K_0 = \{11, 12, 13\}$, and the positive critical point sits over the leximin coupling $(2/15, 2/15)$. The product coupling $f \otimes g = (0.12, 0.172)$ has the single leading cell 13, at $0.108 < h^*$, so it is not a candidate. That is the Tier-B reason why "the product coupling carries no brane" [4].

Example 24 (the five critical points)**.** $W = y_1 + y_2 + T^{2/5}/(y_1 y_2) + T^{3/10}/y_1 + T^{-13/100} y_1 y_2$ has 5 critical points, twice the area 5/2 of its Newton polygon (Kushnirenko). Exact resultants at $T = 10^{-100}$ and 10^{-200} place them as follows.

- Three lie over the leximin coupling, one of them positive.
- One is real with $y_1, y_2 < 0$, over $(13/100, 13/100)$. Its leading cells are $\{11, 12, 23\}$, whose normals have the signed balance $(1,0) + (0,1) - (1,1) = 0$ but no positive one.
- One is real with $y_2 < 0$, over $(1/5, 1/10)$. Its leading cells are $\{12, 13, 21\}$, also with a signed balance only.

This is theorem 3(4) on the example. By the cited theorem [11], three torus fibres have non-vanishing Floer cohomology. This agrees with [4, Thm. 8.2]. CF5 checks the count, the positions, the signs and the leading sets.

11.4 The positive thimble on couplings

By theorem 8, the positive thimble with $\hbar = T^h$, read in $u = \log y / \log T$, tends to the uniform law on $Q_h = \{\alpha : \text{every facet cell} \geq h\}$ for $h < h^*$, and concentrates on argmax at h^*. On the fibre this has a plain reading. At $h = 0$ it is the uniform law on all couplings, which is Duistermaat–Heckman, and the limit of the uniform law on tables. As h grows it is uniform on the couplings whose every cell is at least h. At $h^* = 2/15$ it is the leximin coupling (fig. 5; CF7). The thimble is thus a one-parameter family of probability laws on couplings, from "all equally likely" to "the fairest one".

11.5 Two kinds of wall, and one count of data

In the base, the margins (f, g) move the Kähler class of the fibre. Two families of walls cut the base.

- *Kähler-chamber walls* $f(I) = g(J)$. Across them the polygon gains or loses a side, which is a toric blow-up or blow-down [5]. These are the viability walls of the ledger.




Figure 5: The positive thimble on the fibre, at four values of h, with $\log T = -400$. Dashed: the shrunk polygon Q_h, every cell at least h. The law is uniform on Q_h, and at $h^* = 2/15$ all its mass is at the leximin coupling (circle). The product coupling (square) is passed over.

- *Leximin walls*, where two of the ratios f_i/m, g_j/n tie. Across them the binding line, and with it the leading cells and the candidate dimension, jumps (theorem 23). These are the outcome walls of corollary 12, here explicit.

They are different walls (fig. 6(a)). The monotone point, with uniform margins, lies on all of them.

Example 25 (one count)**.** In the example, $f_1/3 = 2/15$ and $g_3/2 = 27/200$ differ by only $1/600$. Move one observation of 100 from column 3 to column 2, so that $g = (30, 44, 26)/100$. Then:

- column 3 binds, $h^* = 13/100$, and the leading cells are $\{13, 23\}$;
- the candidates become a segment, and the leximin coupling, selected at the next level, is $(27/200, 27/200)$;
- the real critical point with $y_1, y_2 < 0$ moves onto the leximin fibre, so the distinct valuations drop from three to two. By the cited theorem, so does the number of torus fibres with non-vanishing Floer cohomology.

The surface does not change: the nearest blow-up wall, $f_1 = g_2$, is three counts away. So in this example the least data that changes the perspective is one count, and it changes the leading cells and the Floer fibres, not the manifold (fig. 6(b); CF8, with exact resultants at $T = 10^{-100}$ and 10^{-300}).

11.6 The selection theorem, and a column-binding example

Theorem 3 and proposition 11 put the valuation of the positive critical point in the candidate set. When that set is a point, this is the whole answer. When it is not, as for a binding column, the next level selects. This subsection proves that the levels continue all the way down, for any configuration.

Definition 26 (lexicographic max–min)**.** Let $\ell_k(u) = \langle v_k, u\rangle - \lambda_k$ with $v_1, \dots, v_m$ positively spanning $\mathbb{R}^n$. Put $A_0 = \mathbb{R}^n$ and $S_0 = \emptyset$. For $j \geq 1$, let h_j be the maximum over A_{j-1} of the least ℓ_k with k not yet fixed. Let A_j be the set where it is attained, and let S_j be the not-yet-fixed k with $\ell_k \equiv h_j$ on A_j. Stop when every k is fixed. Then A_J is a single point, the *lexicographic max–min point* u^{lex}.




Figure 6: Margin space, in the slice $g_1 = 0.3$. (a) Colours show which row or column binds. Dashed lines are the leximin walls and solid lines the blow-up walls $f_i = g_j$. The example (circle) sits almost on a leximin wall, and one count (cross) crosses it. (b) Along $f_1 = 0.4$: the three ratios $f_1/3$, $g_3/2$, $g_1/2$, whose minimum is h^*. The leximin wall is a third of a count from the example, while the blow-up wall $f_1 = g_2$ is three counts away.

Each S_j is nonempty (proposition 11(2), applied on A_{j-1}), and each A_j is bounded, so the recursion stops. The fixed normals span $\mathbb{R}^n$, so A_J is a point. On $\mathbb{CP}^2$ and on F_1 with $\alpha < 1/3$ there is one level. On F_1 with $\alpha > 1/3$ there are two, as in example 9.

Theorem 27 (selection)**.** *Let $a_k(L) \in [c, C] \subset (0, \infty)$, and let u_L minimise $F_L(u) = \sum_k a_k(L) e^{-L\ell_k(u)}$. Then $|u_L - u^{\mathrm{lex}}| = O(1/L)$ as $L \to \infty$. In particular, the valuation of the positive critical point of W_T, with $L = \log(1/T)$ and $u_L = \log y_T / \log T$, is the lexicographic max–min point of the facet functions, and the error at finite T is $O(1/\log(1/T))$.*

Proof. F_L is strictly convex and coercive, because the v_k positively span, so u_L is unique. We argue by induction on n; the case $n = 0$ is empty.

Level one is pinned to $O(1/L)$. For $\bar{u} \in A_1$, $F_L(u_L) \le F_L(\bar{u}) \le mCe^{-Lh_1}$. So each term satisfies $ce^{-L\ell_k(u_L)} \le mCe^{-Lh_1}$, that is, $\ell_k(u_L) \ge h_1 - \kappa/L$ with $\kappa = \log(mC/c)$. By proposition 11(2) there is a balanced law q supported exactly on S_1, and then $\sum_{k \in S_1} q_k \ell_k \equiv h_1$. Hence $\ell_k(u_L) \le h_1 + \kappa/(Lq_{\min})$ for $k \in S_1$.

The first level drops out. Let $D_1 = \{d : \langle v_k, d\rangle = 0,\ k \in S_1\}$ be the direction space of $E_1 = \operatorname{aff} A_1$, and P the orthogonal projection onto D_1. Then $Pv_k = 0$ for $k \in S_1$. So projecting $\nabla F_L(u_L) = 0$ gives, exactly,

$$\sum_{k \notin S_1} a_k(L)\, e^{-L\ell_k(u_L)}\, Pv_k = 0.$$

Write $u_L = p_L + w_L$, with $p_L \in E_1$ and $w_L \perp D_1$. The v_k, $k \in S_1$, span $D_1^{\perp}$, and $\langle v_k, w_L\rangle = \ell_k(u_L) - h_1 = O(1/L)$, so $|w_L| = O(1/L)$. Hence p_L is a critical point, and so the minimiser, of $F_L^{(1)}(p) = \sum_{k \notin S_1} a'_k(L) e^{-L\ell_k(p)}$ on E_1. Here $a'_k = a_k e^{-L\langle v_k, w_L\rangle}$ again lies in a fixed compact subset of $(0, \infty)$.

Induction. On E_1 the remaining ℓ_k have projected normals Pv_k that positively span D_1, because $A_1 = \{p \in E_1 : \ell_k(p) \geq h_1,\ k \notin S_1\}$ is bounded. Their lexicographic max–min point is u^{lex}. Indeed, off A_1 some remaining ℓ_k is below h_1, while on the relative interior of A_1 all exceed it, so the reduced recursion is levels $2, 3, \ldots$ of the original. Since $\dim D_1 < n$, the induction hypothesis gives $|p_L - u^{\mathrm{lex}}| = O(1/L)$, and so $|u_L - u^{\mathrm{lex}}| = O(1/L)$. LX1–LX3 corroborate the statement and the mechanism, on 30 random configurations in multiprecision arithmetic. □

Remark 28 (provenance)**.** The procedure is the one Fukaya, Oh, Ohta and Ono use to locate a non-displaceable fibre [23]. Judd and Rietsch construct the tropical critical point recursively, and prove that it maximises the tropicalisation [32]. A related hierarchical selection governs the exponential-penalty trajectory in linear programming [15]. The statement is therefore adapted; the proof above is direct, and it gives the rate. The constant in $O(1/L)$ depends on the gaps between levels. That is why the near tie of example 25 converged slowly.

Corollary 29 (the selected coupling)**.** *For nondegenerate $n \times m$ margins, the positive critical point of the potential of* $\mathrm{Cpl}(f, g)$ *sits over the facet-leximin coupling. In* 292 *random margins of shapes* 2×3*,* 3×3 *and* 2×4*, this equals the leximin coupling over all cells (LX4). We have not proved that the two always agree.*

Example 30 (a column-binding fibre)**.** Repurpose Example 3.4(b) of [5]: $f = (2/5, 3/5)$, $g = (1/2, 3/10, 1/5)$. The fibre is the pentagon with vertices $(1/5, 0)$, $(2/5, 0)$, $(1/10, 3/10)$, $(0, 3/10)$, $(0, 1/5)$ and area $11/200$. Its facets are the cells $11, 12, 13, 22, 23$. Cell 21 never vanishes, since $f_2 + g_1 = 11/10$. The ratios are $f_i/3 = (2/15, 1/5)$ and $g_j/2 = (1/4, 3/20, 1/10)$, so column 3 binds. The levels are:

level	value	cells fixed	what is decided
1	$h_1 = 1/10$	$13, 23$	column 3 is spread evenly; the candidates form the open segment $x + y = 3/10$, $1/10 < x < 1/5$
2	$h_2 = 3/20$	$11, 12, 22$	along the segment, the least remaining cell is largest at the midpoint

So $u^* = (3/20, 3/20)$ (fig. 7(a),(b)). The endpoints are maximisers but not candidates: their leading sets, $\{11, 13, 22, 23\}$ and $\{12, 13, 23\}$, have no positive balance. At finite T,

$$x_L = \tfrac{3}{20} + \frac{\ln 2}{2L} + o(1/L),$$

because cells 11 and 22 coincide on the segment (both equal x) and balance cell 12 alone: $2e^{-Lx} = e^{-L(3/10-x)}$ (fig. 7(c)). Of the five critical points, the positive one and a real one with $y_1, y_2 < 0$ sit over $(3/20, 3/20)$, both with this correction. The other three (one real, two complex) lie over the fibre $(2/15, 1/6)$ on the candidate segment, off the leximin point. Section 12.7 derives this exactly. The slow approach seen at finite T is the $1/\log(1/T)$ effect of their leading moduli $2^{\pm 1/3}$. The positive thimble shows the two levels as h grows: it is uniform on the segment at $h = 1/10$, uniform on $[h, 3/10 - h]$ between the levels, and concentrated at $3/20$ above (fig. 7(d)). The one-count example of example 25 is column-binding too, with levels $(13/100;\ 13, 23)$ and $(27/200;\ 11, 12)$. LX5–LX9 check every number here, the first exactly.




Figure 7: The selection theorem on a column-binding fibre. (a) Level 1 fixes cells 13 and 23 at $1/10$ and leaves a segment of candidates, whose endpoints are not candidates. Level 2 picks its midpoint. (b) Along the segment, the least remaining cell is largest at $x = 3/20$. (c) At finite T the selected point is offset by $\ln 2/(2L)$, because two cells pull against one. (d) The positive thimble along the segment shows both levels as h grows.

11.7 The example through every layer

layer	in the example	where
fibre	pentagon, area 1331/20000, facets $11, 12, 13, 21, 23$; 715 tables (Pick)	example 21; CF1–2
Tier A	Fisher's noncentral hypergeometric family; odds ratios 1 give $f \otimes g = (0.12, 0.172)$; the leximin needs $(0.82, 0.46)$	proposition 22; CF3
Tier B	$h^* = f_1/3 = 2/15$; row 1 binds; one candidate, the leximin coupling $(2/15, 2/15)$	theorem 23; CF4, CF6
selection	the valuation is the lexicographic max–min point, error $O(1/\log(1/T))$; a binding column needs a second level	theorem 27 and example 30; LX1–LX9
sign split	five critical points; signed fibres over $(13/100, 13/100)$ and $(1/5, 1/10)$	example 24; CF5
Floer (cited)	three fibres with non-vanishing Floer cohomology; none over $f \otimes g$	[11, 4]
thimble	uniform on couplings at $h = 0$; uniform on Q_h; the leximin coupling at h^*	theorem 8; CF7
walls	leximin wall $1/600$ away in ratio; blow-up wall $f_1 = g_2$ three counts away	fig. 6
one count	column 3 binds; candidates a segment; three Floer fibres become two; same surface	example 25; CF8
not claimed	other thimbles, Stokes data, stability conditions, the category	section 14

Part III
Towards morphisms

12 Objects, morphisms and a categorical comparison

What this section shows, in plain words. So far we have found *which* torus fibres matter. A category also needs the arrows between them.

- Every critical point of W is an *object*: a torus fibre with a local system. Distinct objects have no arrows between them, and each object's arrows to itself form a Clifford algebra. For the probabilistic object, its quadratic form is the Fisher information of the conditioned law (fig. 8).
- Up to equivalence, an outcome wall does *not* change the category. It moves objects between fibres.
- It does leave two traces. The Clifford form of the probabilistic object, reduced modulo T, drops rank exactly when the candidate set stops being a point (fig. 9). And the Stokes walls, where the basis of thimbles mutates, crowd into a window of width $O(1/\log(1/T))$ around the outcome wall (fig. 10).

The computations are all on the critical points, values and Hessians of W. The Floer reading of those data is cited, and its status is marked in each statement.

12.1 Objects

By [11], a critical point c of W with valuation u_c gives the torus fibre $L(u_c)$ with a rank-one local system. Its holonomy has leading term y_c/T^{u_c}, and $L(u_c)$ has non-vanishing Floer cohomology. We call the pair an *object*. In our examples W has $\chi(X)$ critical points, twice the area of its Newton polygon: 4 for F_1 and 5 for $\mathrm{Bl}_2\mathbb{CP}^2$.

example	objects: fibre, type, leading holonomy
F_1, $\alpha = 1/5$	over the barycentre: positive $(1,1)$, complex (ζ,ζ), (ζ^2,ζ^2), with $\zeta^3=1$; over $(1/5,3/5)$: real-signed $(-1,1)$
F_1, $\alpha = 3/5$	all four over $(2/5,1/5)$: positive $(1,1)$, real-signed $(-1,1)$, complex $(i,-1)$, $(-i,-1)$
2×3 example	over the leximin coupling $(2/15,2/15)$: positive and two complex; over $(13/100,13/100)$: real $(-1,-1)$; over $(1/5,1/10)$: real $(1,-1)$
one count later	over $(27/200,27/200)$: positive, real $(-1,-1)$, two complex; over $(1/5,1/10)$: real $(1,-1)$

MO1 and MO2 check this table, the first part from the quartic and the second from exact resultants at $T = 10^{-200}$.




Figure 8: Objects over their torus fibres (crosses). Each box is a fibre; each marker is an object, labelled by its leading holonomy; each loop is its endomorphism algebra Cl_2. There are no arrows between distinct objects. (a)→(b): crossing the F_1 wall merges two fibres into one. (c)→(d): one count of data merges the leximin fibre with the nearby real fibre. In both cases the objects persist; only their fibres change.

12.2 Morphisms

Proposition 31 (morphisms between the objects)**.** *(1) Objects over distinct fibres have no morphisms: distinct moment fibres are disjoint, so their Floer complex is zero (exact).*

(2) Distinct objects over one fibre have distinct critical values $W(c) = m_0$. Morphisms are defined

only between objects with equal m_0, so the category splits into summands $\mathcal{F}_\lambda$ [3, 23], and they have no morphisms. This uses distinct m_0 only. Their holonomies also differ, but we do not use that to separate objects: an eigenvalue decomposition cannot by itself separate two objects with the same m_0. That these objects generate their summands is proved in the monotone case and conditionally otherwise [40]; our examples are not monotone.

(3) The Floer cohomology ring of an object is the Clifford algebra of the Hessian $Q = \sum_k c_k v_k v_k^\top$ of W at c, where c_k are the values of its terms [12, Thm. 5.6]. So it has dimension 2^n.

(4) For the positive object, $Q = W \cdot \mathrm{Cov}_w(v)$, where w is the law of the terms of W at y_T (mean exponent 0). Its quadratic form is the Fisher information of the conditioned law, scaled by W. It is real and positive definite.

Proof. (1) $L(u) = \mu^{-1}(u)$. (2) and (3) are the cited theorems; MO3 checks that the critical values (and, for the record, the holonomies) are distinct in every example. (4) $Q = \sum_k c_k v_k v_k^\top$ with $c_k = W w_k$, and $\sum_k w_k v_k = 0$. MO4 and MO5 check the identity, and the Clifford relations with explicit generators that span M_2, for all 18 objects. □

12.3 The A_∞ products

Proposition 32 (no higher products survive)**.** *Over the algebraically closed Novikov field Λ, let Q be nondegenerate. The $\mathbb{Z}/2$-graded algebra $\mathrm{Cl}(Q)$ is separable. For $n = 2$ it is the super matrix algebra $\mathrm{End}(\Lambda^{1|1})$. Its Hochschild cohomology vanishes in positive degrees. So every A_∞ structure on it, with $m_1 = 0$ and this m_2, is A_∞-isomorphic to the formal one: it is intrinsically formal [34, 38]. On the subcategory generated by these objects, the higher products m_k for $k \geq 3$ can be removed by an A_∞ isomorphism.*

Proof. A separable algebra has vanishing Hochschild cohomology in positive degrees. The obstructions to formality lie in $HH^{k,2-k}$ for $k \geq 3$, so they vanish. MO5 exhibits $\mathrm{Cl}(Q) \cong M_2$ for every object. The statement rests on the cited m_2 of proposition 31(3); we do not compute the A-side higher products themselves. □

So, within what the cited theorems give, the category generated by the objects is $\bigoplus_c \mathrm{Cl}(Q_c)$. It is semisimple, with one summand per critical point, matching $\dim QH^*(X) = \chi(X)$. On the mirror side, matrix factorisations at a nondegenerate critical point have the same Clifford description [19].

12.4 The Clifford shadow of selection

Over Λ, all Clifford algebras of nondegenerate forms of rank n are isomorphic, so (2)–(3) cannot see an outcome wall. The integral form can.

Theorem 33 (the Clifford shadow)**.** *Write the Clifford form of the positive object in the integral basis of $H^1(L;\mathbb{Z})$ and divide it by W. Then $Q/W = \mathrm{Cov}_w(v)$ has entries of non-negative valuation. Its reduction modulo T is $\mathrm{Cov}_{q^*}(v)$, where q^* is the limit law of theorem 3(2). Its rank is $\dim \sigma_0 = n - \dim(\text{candidate set})$. So the reduced Clifford algebra is semisimple exactly when the candidate set is a single point, that is, exactly when leading order selects.*

Proof. w_k has valuation $\ell_k(u^*) - h^* \geq 0$, and $w \to q^*$ by theorem 3(2). $\mathrm{Cov}_{q^*}(v) = \sum_k q_k^* v_k v_k^\top$, because q^* has mean 0. Its rank is the dimension of the span of $\{v_k : k \in K_0\}$, which is $\dim \sigma_0$ because $0 \in \mathrm{relint}\, \sigma_0$. Then apply proposition 11(2). MO6 checks the convergence and the ranks. □

In the examples, the other eigenvalues have valuation equal to the gap between selection levels. On F_1 above the wall the small eigenvalue is $\sim T^{(3\alpha-1)/4}$, where $(3\alpha-1)/4 = h_2 - h_1$; it closes at the wall (fig. 9(a)). On the 2×3 fibre, one count turns rank 2 into rank 1, with valuation $1/200 = h_2 - h_1$ (fig. 9(b)). In the column-binding example the valuation is $1/20$. So the hierarchy of selection reappears as the hierarchy of scales of the probabilistic object's Clifford form.




Figure 9: The Clifford shadow. (a) F_1: eigenvalues of Q/W for the positive object (solid: the small one; dotted: the large one) at three values of T. Above the wall the small eigenvalue tends to 0 like $T^{(3\alpha-1)/4}$, so the form reduced mod T has rank 1. (b) The 2×3 fibre, before and after one count. After the move the small eigenvalue decays like $T^{1/200}$.

12.5 Comparing the two sides of a wall

Three mechanisms could relate the two sides of a wall: an A_∞ functor, a continuation map, or a mutation. Here is what each does.

Theorem 34 (an A_∞ equivalence across the F_1 wall)**.** *On F_1, for every $T \in (0,1)$ and every $\alpha \in (0,1)$, the four critical points are distinct and nondegenerate. So they form four continuous branches in α, and the semisimple categories $\bigoplus_c \mathrm{Cl}(Q_c(\alpha))$ over Λ are related by algebra isomorphisms that depend continuously on α. For the positive object these are real: $A^\top Q(\alpha')A = Q(\alpha)$. In particular they are A_∞ equivalences across $\alpha = 1/3$. What changes at the wall is the* support*: which fibre carries which object. That is the merging of fibres in fig. 8.*

Proof. The discriminant $-27T^2 - 256T^{3-3\alpha}$ never vanishes (example 14(d)), so the critical points never collide, and nondegeneracy follows. A nondegenerate complex symmetric form has a symmetric square root, which gives the isomorphisms. MO7 checks the path $\alpha \in [1/5, 3/5]$ at $T = 10^{-24}$. The A-side reading is that of propositions 31 and 32, so the statement inherits their status. □

Continuation maps compare Floer theory of Hamiltonian-isotopic Lagrangians in one symplectic manifold. Here the evidence moves the Kähler class itself, and distinct fibres are not Hamiltonian isotopic. So there is no continuation map in the strict sense: the family comparison of theorem 34 takes its place.

Mutations belong to the category of thimbles, the Fukaya–Seidel category of the Landau–Ginzburg model. For a phase θ of $\hbar$, a distinguished basis of thimbles changes by a mutation when two critical values align along θ [39].

Proposition 35 (Stokes walls crowd the outcome wall)**.** *On F_1, the Stokes walls for the phases $\theta = 0.3$, 1 and $\pi/2$ lie within $4/\log(1/T)$ of $\alpha = 1/3$, for $T = 10^{-12}, 10^{-24}, 10^{-48}$, and $\log(1/T)\,(\alpha_\theta - 1/3)$ does not depend on T. Away from the window there are none. The mechanism is that $|W|$ is $\sim T^{\alpha}$ for the signed object and $\sim T^{1/3}$ for the barycentre objects. Outside a window of width $O(1/\log(1/T))$ one of them dominates, and the phases of the differences are frozen; inside it their ratio sweeps $(0, \infty)$, and every phase is crossed. For real $\hbar$ ($\theta = 0$) the positive thimble never mutates: it is the positive orthant for every λ (proposition 7).*

This is a computation, MO8 and MO9, with the mechanism sketched; we have not proved the window for general configurations.




Figure 10: Mutations and the outcome wall on F_1. (a) The sizes of the four critical values at $T = 10^{-24}$: the signed object's value dominates below the wall, and the barycentre objects' values above it. (b) The Stokes walls for three phases, at three values of T, lie on lines through $(0, 1/3)$ in the variable $1/\log(1/T)$.

12.6 What sees the wall

invariant	across the outcome wall	status
which fibres carry objects	changes ($2 \to 1$; $3 \to 2$)	cited [11]; MO1–2
the category over Λ, $\bigoplus \mathrm{Cl}(Q_c)$	unchanged, up to A_∞ equivalence	theorem 34, on cited m_2
higher products	none survive	proposition 32, on cited m_2
the positive object's Clifford form mod T	rank drops ($2 \to 1$)	theorem 33, new
the thimble basis at phase $\theta \neq 0$	mutates within $O(1/\log(1/T))$ of the wall	proposition 35, computed
the positive thimble, real $\hbar$	never changes	proposition 7, proved

So a data-induced outcome wall is not a wall of the Fukaya category over Λ. It is a wall of its *integral* structure for the probabilistic object, and the place where, at finite T, the thimble bases mutate. The full A_∞ category over Λ_0, and split generation in the non-monotone case, are left open.

12.7 The column-binding example, classified

The column-binding fibre of example 30 is the first example whose objects are not all over the selected fibre, and whose holonomies are not all of modulus one. Stage IV (section 12.8) shows why: the example lies on a level-two leximin wall, and both features belong to the wall. We classify its five critical points exactly, then compute the morphisms between the resulting objects. Wall-crossing from this example is deferred.

The Floer convention. We use the convention of [23]. An object is a pair $(L(u), b)$ with $b \in H^1(L(u); \Lambda_0)$, whose holonomy coordinates are $y_i = T^{u_i} e^{b_i}$. Their leading coefficients $e^{b_{i,0}}$ are arbitrary nonzero complex numbers, not only those of modulus one, so local systems with non-unit leading modulus are included. On a compact toric Fano manifold the potential is W [11]. The pair has non-vanishing Floer cohomology iff y is a critical point of W, and $m_0(L(u), b) = W(y) \cdot 1$ [23]. For the Clifford description of the endomorphism ring we use the formula $Q = \operatorname{Hess} W$ at y [12, Thm. 5.6]. Cho states it in the setting of [11], and its use with non-unit leading coefficients is the one place where our citation goes beyond the original setting.

With $\tau = T^{1/10}$, the potential of this fibre is $W = y_1 + y_2 + \tau^4/(y_1 y_2) + \tau^3/y_2 + y_1 y_2/\tau^2$ (cells $11, 12, 13, 22, 23$).

Proposition 36 (the five critical points)**.** *(1)* Reduction. $y_1 \partial_1 W - y_2 \partial_2 W = y_1 - y_2 + \tau^3/y_2$. *So $y_1 = s - \tau^3/s$ with $s = y_2$, and the remaining equation is exactly*

$$q(s) = s^5 + \tau^2 s^4 - 2\tau^3 s^3 - 2\tau^5 s^2 + \tau^8 = 0.$$

Since $q(0) \neq 0$ and q has no common root with $s^2 - \tau^3$, the critical points correspond one to one to the roots of q.

(2) Newton polygon. *The lower hull of q has vertices $(0,8)$, $(3,3)$, $(5,0)$. The edge of slope $-5/3$, with edge polynomial $1 - 2c^3$, gives three roots $s \approx 2^{-1/3} \omega\, \tau^{5/3}$ with $\omega^3 = 1$. The edge of slope $-3/2$, with edge polynomial $c^2 - 2$, gives two roots $s \approx \pm\sqrt{2}\, \tau^{3/2}$.*

(3) Distinct and nondegenerate for every $T > 0$. $\operatorname{disc}_s q = -\tau^{31}(256\tau^2 - 837\tau + 3456)$, *and the resultant of q with the Hessian determinant is $\tau^{54}(256\tau^2 - 837\tau + 3456)$. The quadratic has discriminant $-2838375 < 0$, so neither vanishes for $\tau > 0$.*

(4) Distinct critical values for every $T \in (0, 1)$. $W = 2s + \tau^4/(s^2 - \tau^3) + (s^2 - \tau^3)/\tau^2$ *at a critical point. The polynomial whose roots are the five critical values has discriminant $-\tau^{225}(\tau - 1)^2(256\tau^2 - 837\tau + 3456)^3$, which vanishes on $(0, \infty)$ only at $T = 1$.*

Proof. Each statement is an exact symbolic computation (CB1–CB4). In (2) the edge polynomials are read off the coefficients on each edge: $\tau^8 - 2\tau^3 s^3$ and $-2\tau^3 s^3 + s^5$. □

So there are five critical points, two fibres and five objects:

object	fibre	leading $(y_1, y_2)/T^u$	leading W	leading $\det(Q/W)$
P positive	$(\frac{3}{20}, \frac{3}{20})$	$(\frac{\sqrt{2}}{2},\ \sqrt{2})$	$2\tau + 2\sqrt{2}\,\tau^{3/2}$	$\sqrt{2}\,\tau^{1/2}$
N real $(-,-)$	$(\frac{3}{20}, \frac{3}{20})$	$(-\frac{\sqrt{2}}{2},\ -\sqrt{2})$	$2\tau - 2\sqrt{2}\,\tau^{3/2}$	$-\sqrt{2}\,\tau^{1/2}$
R real $(-,+)$	$(\frac{2}{15}, \frac{1}{6})$	$(-2^{1/3},\ 2^{-1/3})$	$-2\tau + 2^{2/3}\tau^{5/3}$	$-c\,\tau^{2/3}$
C_ω, $C_{\bar\omega}$ complex	$(\frac{2}{15}, \frac{1}{6})$	$(-2^{1/3}\bar\omega,\ 2^{-1/3}\omega)$	$-2\tau + 2^{2/3}\omega\,\tau^{5/3}$	$-c\,\bar\omega^2\tau^{2/3}$

Here $c = 3 \cdot 2^{-4/3}$, $\omega = e^{\pm 2\pi i/3}$, and $C_{\bar{\omega}}$ is the conjugate of C_ω. CB5 and CB6 check every entry against the roots at $\tau = 10^{-10}, 10^{-20}, 10^{-40}$, with relative errors $1.5 \cdot 10^{-4}$, $6.8 \cdot 10^{-8}$ and $1.5 \cdot 10^{-14}$. The two groups have opposite leading critical value, $\pm 2\tau$. Within a group the values separate at order $\tau^{3/2}$, or at order $\tau^{5/3}$ through the cube roots of unity (fig. 11).




Figure 11: The column-binding example classified. (a) The Newton polygon of the quintic: one edge gives the three branches over $(2/15, 1/6)$, the other the two over the leximin fibre. (b) The five roots against τ follow the two predicted lines; the vertical offsets are the moduli of the leading coefficients.

Remark 37 (the drift explained)**.** At finite T a branch with leading coefficient ρ has apparent valuation $u - \log|\rho|/\log(1/T)$. For P, $\log(\sqrt{2}/2)$ gives the offset $\ln 2/(2L)$ of example 30. For the other three, $\log 2^{\pm 1/3}$ gives $\pm \ln 2/(3L)$. This was the slow "drift" seen there: the limits are exactly $(2/15, 1/6)$. The three sit there because their level-one terms have $y_1 y_2 \approx -\tau^3$. This sign makes the level-two terms (cells 11 and 22) cancel exactly, so the position is set by the next balance, of cell 12 against the constant term: $-2\tau^3 s^3 + \tau^8 \approx 0$. In the row-binding examples every leading coefficient has modulus one. Non-unit moduli appear here because a level of selection cancels.

Theorem 38 (reduction of the Clifford form)**.** *Let c be a nondegenerate critical point with valuation u_c. Let K be its leading set, the k with $\ell_k(u_c)$ minimal, equal to h_c. Write the term values as $c_k = \rho_k T^{h_c}(1 + o(1))$ for $k \in K$, and assume*

(H1) $\sum_{k \in K} \rho_k \neq 0$.

Then Q/W has entries of non-negative valuation and reduces modulo T to $M_c = \sum_K \rho_k v_k v_k^\top / \sum_K \rho_k$. Always $\operatorname{rank} M_c \leq \dim \operatorname{span}\{v_k : k \in K\}$. *Equality holds under*

(H2) the ρ_k, $k \in K$, are real and of one sign,

in particular for the positive object (theorem 33). Without (H2), cancellation can lower the rank.

Proof. By (H1), $W = (\sum_K \rho_k) T^{h_c}(1 + o(1))$, and every c_k has valuation at least h_c. Under (H2), $\sum_K \rho_k v_k v_k^\top$ is a semidefinite Gram matrix, whose rank is that of $\{v_k\}$. CB8 tests the theorem on 23 objects: F_1 at $\alpha = 1/5$ and $3/5$, the 2×3 example, its one-count neighbour, and this example. (H1) holds for all 23, and (H2) for 13. The rank equals the span dimension in all 23, including the 10 without (H2). An example is the signed object of F_1 at $\alpha = 1/5$, where $\rho = (-1, -1, 1)$ and $M_c = \left(\begin{smallmatrix} 2 & 1 \\ 1 & 0 \end{smallmatrix}\right)$. □

In the column-binding example $K = \{13, 23\}$ for all five objects, with $\rho = (1, 1)$ on the leximin fibre and $(-1, -1)$ on the other. (H2) holds, and $M_c = J = \left(\begin{smallmatrix} 1 & 1 \\ 1 & 1 \end{smallmatrix}\right)$, of rank 1, for every object. The reduced Clifford algebras are therefore all non-semisimple. The objects are told apart one order further:

$$P, N: \quad \frac{Q}{W} = J \pm \frac{\tau^{1/2}}{2} \operatorname{diag}\Big(\tfrac{\sqrt{2}}{2}, \tfrac{3\sqrt{2}}{2}\Big) + \dots,$$
$$R, C: \quad \frac{Q}{W} = J + 2^{-2/3}\bar{\omega}\,\tau^{1/3} \operatorname{diag}(1, -1) + \dots$$

For P and N the correction is first order in the small direction, so the small eigenvalue is $\pm(\sqrt{2}/2)\tau^{1/2}$, of valuation $1/20$ in T. The form is positive definite for P and indefinite for N. For R and C the first-order correction vanishes in the small direction, which is the cancellation the reduction theorem warns about. The small eigenvalue appears only at second order, $-\frac{3}{2}\,2^{-4/3}\bar{\omega}^2\tau^{2/3}$, of valuation $1/15$.

Proposition 39 (the Hom table)**.** *In the convention above, the five objects have the following morphisms.*

(1) Between the two fibres: none. The fibres are disjoint, so the unperturbed Floer complex is zero; this is a legitimate model, since the pair is transverse.

(2) Within a fibre: none. The critical values are distinct (proposition 36(4)), and morphisms are defined only between objects with equal m_0 [3, 23]. (Their holonomies are also distinct, but the argument does not use this.)

(3) On the diagonal: $\mathrm{Cl}(Q_c) \cong M_2$, with Q_c as above (cited, in the convention stated). No higher products survive (proposition 32).

So the category these objects generate is $\bigoplus_{c=1}^{5} \mathrm{Cl}(Q_c)$ over Λ. Over Λ_0, all five reduce to the rank-one form J. Generation of the Fukaya summands remains conditional, because the example is not monotone [40].


Proof. (1) $L(u) = \mu^{-1}(u)$. (2) and (3) are the cited statements together with proposition 36. CB7 checks the counts, the distinct values and holonomies within each fibre, and that explicit Clifford generators span M_2. □

12.8 Stage IV: a path through the column-binding example

Data so far moved margins by one count (a point) or along one direction of evidence (a line). Here we take a sheet of margins and one path across it, and we keep four questions separate along the path: do critical points collide, do critical values coincide, which fibres carry the objects, and what do the Clifford forms and the Stokes alignments see?

The sheet and the path. Fix $f = (2/5, 3/5)$ and let the column margins vary: $g = (1 - g_2 - g_3,\ g_2,\ g_3)$. Figure 13 shows this sheet with its blow-up walls $f_1 = g(J)$, which bound the Kähler chambers, and its leximin walls. The path is

$$g(\gamma) = \Big(\tfrac{7}{10} - \gamma,\ \tfrac{3}{10},\ \gamma\Big), \qquad \gamma \in \big[\tfrac{3}{20}, \tfrac{11}{40}\big].$$



Hom between the five objects ($c = 3 \cdot 2^{-4/3}$): only the diagonal survives.
First two: the leximin fibre; last three: the fibre (2/15, 1/6)

	P positive	*N* real (−, −)	*R* real (−, +)	C_ω complex	$C_{\bar{\omega}}$ complex
P positive	$\mathrm{Cl}(Q) \cong M_2$, $\det \sim \sqrt{2}\,\tau^{1/2}$, reduced: rank 1	0, same fibre, distinct m_0	0, disjoint fibres	0, disjoint fibres	0, disjoint fibres
N real (−, −)	0, same fibre, distinct m_0	$\mathrm{Cl}(Q) \cong M_2$, $\det \sim -\sqrt{2}\,\tau^{1/2}$, reduced: rank 1	0, disjoint fibres	0, disjoint fibres	0, disjoint fibres
R real (−, +)	0, disjoint fibres	0, disjoint fibres	$\mathrm{Cl}(Q) \cong M_2$, $\det \sim -c\,\tau^{2/3}$, reduced: rank 1	0, same fibre, distinct m_0	0, same fibre, distinct m_0
C_ω complex	0, disjoint fibres	0, disjoint fibres	0, same fibre, distinct m_0	$\mathrm{Cl}(Q) \cong M_2$, $\det \sim -c\,\bar{\omega}^2\tau^{2/3}$, reduced: rank 1	0, same fibre, distinct m_0
$C_{\bar{\omega}}$ complex	0, disjoint fibres	0, disjoint fibres	0, same fibre, distinct m_0	0, same fibre, distinct m_0	$\mathrm{Cl}(Q) \cong M_2$, $\det \sim -c\,\omega^2\tau^{2/3}$, reduced: rank 1

Figure 12: Top: the five objects over their two fibres, with exact leading holonomies; the leximin fibre carries P and N, the fibre $(2/15, 1/6)$ carries R, C_ω and $C_{\bar{\omega}}$. Bottom: the Hom table. Grey cells vanish because the fibres are disjoint, orange cells because the values of m_0 differ. The diagonal is $\mathrm{Cl}(Q_c) \cong M_2$, and every one reduces to the rank-one form J modulo T.

It stays in the pentagon chamber (facets $11, 12, 13, 22, 23$; no blow-up wall meets $\gamma \in (1/10, 3/10)$), and only the exponent of cell 23 moves. With $A = T^{2/5}$, $B = T^{3/10}$ and $C = T^{\gamma - 2/5}$,

$$W_\gamma = y_1 + y_2 + \frac{A}{y_1 y_2} + \frac{B}{y_2} + C\, y_1 y_2.$$

The column-binding example of section 12.7 is $\gamma = 1/5$.




Figure 13: The sheet of column margins at $f = (2/5, 3/5)$. Black lines are blow-up walls. The stepped rose lines are leximin walls computed on a 120×120 grid (colours: leximin types), and the dashed lines are the two walls the path crosses, drawn exactly: $g_2 + g_3/2 = 2/5$ (level two) and $g_3 = 4/15$ (level one). Grey numbers count the facets of the fibre. The path (blue) runs inside one Kähler chamber, the pentagon, and crosses two leximin walls: the column-binding example (black dot) sits on a level-two wall, and the level-one wall (orange square) is at $\gamma = 4/15$.

Proposition 40 (the path, exactly)**.** *(1)* Two leximin walls. *For $\gamma < 1/5$ the levels are $\gamma/2$ (column 3), $3/20$ (column 2) and $1/4 - \gamma/2$ (cell 11). For $1/5 < \gamma < 4/15$ they are $\gamma/2$ and $(4/5 - \gamma)/4$, where the residual of row 1 binds. For $\gamma > 4/15$ row 1 binds at once, at $2/15$. At $\gamma = 1/5$ the cells $11, 12, 22$ tie at the second level, and at $\gamma = 4/15$ the cells $11, 12, 13, 23$ tie at the first. So* the column-binding example lies on a level-two leximin wall.

(2) Reduction. *The critical points are $y_1 = (s^2 - B)/s$, $y_2 = s$, where*

$$(s^2 - B)^2(1 + Cs) = As. \tag{1}$$

At $\gamma = 1/5$, where $C = A/B^2$, this is the quintic of proposition 36.

(3) Where the objects sit. *The Newton polygon of* (1) *changes exactly at the two leximin walls, and gives*

γ	*over the leximin fibre*	*elsewhere*
$(3/20, 1/5)$	4, *at* $(1/4 - \gamma/2, 3/20)$*:* $s = \pm T^{3/20}$, $y_1 = \pm T^{1/4-\gamma/2}$	R, *at* $(\gamma - 1/10, 2/5 - \gamma)$, *signs* $(+,-)$
$1/5$	2, *at* $(3/20, 3/20)$	3, *at* $(2/15, 1/6)$ *(proposition 36)*
$(1/5, 4/15)$	4, *at* $\frac{4/5-\gamma}{4}(1,1)$*:* $y_1 \approx y_2 = i^k T^{(4/5-\gamma)/4}$	R, *at* $(1/10, 1/5)$, *signs* $(-,+)$
$(4/15, 11/40]$	3, *at* $(2/15, 2/15)$*:* $y_1 \approx y_2 = \omega^k T^{2/15}$	N, *at* $(2/5 - \gamma)(1,1)$, *signs* $(-,-)$*; and* R

In every regime the leximin fibre carries the multiple root, as theorem 27 requires of the positive object.

Proof. (1) By theorem 23 and the recursion of definition 26. After column 3 binds, the segment has $a_{11} + a_{12} = r = 2/5 - \gamma/2$, and the facet cells on it are $11, 12, 22$. The residual row binds if $r < g_2 = 3/10$, that is, if $\gamma > 1/5$, and column 2 binds otherwise; row 1 binds at the first level once $\gamma/2 > 2/15$. (2) As in proposition 36: $y_1\partial_1 W - y_2\partial_2 W = y_1 - y_2 + B/y_2$. (3) Three dominant balances in (1) decide everything. Whether Cs or 1 dominates on the leximin roots switches at $\gamma = 4/15$. Whether C or A/B^2 dominates on the isolated root switches at $\gamma = 1/5$. And below $1/5$ the leximin roots have $s^2 = B(1 + o(1))$: writing $s^2 - B = \delta$ gives $\delta^2 \approx A/C$, so $y_1 = \delta/s$ has valuation $1/4 - \gamma/2$. PP2–PP5 check the levels, the Newton polygons at 10 rational γ, and the valuations and phases at $L = 200, 400$. □

So the peculiarities found in section 12.7 belong to the wall, not to column binding. The grouping $2+3$, the fibre $(2/15, 1/6)$ and the leading moduli $2^{\pm 1/3}$ occur only at $\gamma = 1/5$. On either side of it, four objects sit over the leximin fibre with leading moduli one. This is what the remark on the drift meant by a level of selection that cancels: there the linear coefficient $B^2C - A$ of (1) vanishes identically.




Figure 14: The five objects over the pentagon at four points of the path; the marker size and the number give how many objects sit over each fibre. Away from the wall $\gamma = 1/5$ the leximin fibre (blue, star) carries four objects, then three after the level-one wall.

Proposition 41 (what crosses each wall)**.** *Along the path, at fixed small* T*:*

(1) The level-two wall forces a collision. *At* $\gamma = 0.15$ *and* 0.18 *all five critical points are real; at* $\gamma \geq 0.22$*, three are. The coefficients of* (1) *are real and its leading coefficient* C *never*

vanishes, so the number of real roots can change only through a double root. Numerically the discriminant has exactly one zero on the path, at $\gamma_c(T)$, for $T = e^{-50}, e^{-100}, e^{-200}, e^{-400}$, with $\gamma_c = 0.19235,\ 0.19918,\ 0.199985,\ 0.19999999$: it approaches the wall from below, faster than $1/\log(1/T)$. There the two objects of signs $(+,-)$ meet and leave as the complex pair $I_\pm$. At that one parameter a critical point is degenerate.

(2) The level-one wall does not. *The discriminant has no zero on* $(0.21, 0.275]$. *The pair* $I_\pm$ *continues into* $C_\omega, C_{\bar\omega}$, *and* N *leaves the leximin fibre along the diagonal.*

(3) Critical values *never coincide on the path away from the collision, and the order of the real values is constant on each side (PP7). So outside* $\gamma_c(T)$ *the hypotheses of proposition 31 hold at every point: every off-diagonal Hom vanishes, and each endomorphism algebra is* $\mathrm{Cl}(Q_c) \cong M_2(\Lambda)$.

(4) The Clifford shadow sees both walls *(fig. 15(b)). The positive object's scale is* $h_2 - h_1 = 3/20 - \gamma/2$ *below* $1/5$ *and* $1/5 - 3\gamma/4$ *above it, so the scale has a kink at the level-two wall and closes at the level-one wall, where the reduced rank jumps from* 1 *to* 2 *(theorem 33). The four objects over the leximin fibre share this scale;* R *has rank* 2 *throughout.*

Proof. (1) Real-root counting, together with PP5 for the counts and PP6 for the zero. (2) PP6. (3) PP7, and proposition 31(2)–(3). (4) PP8 compares the slopes at $L = 400 \to 800$ with $h_2 - h_1$ from proposition 40(1). □

What (3) does and does not give needs saying plainly. At every parameter away from $\gamma_c(T)$ the five objects generate $\bigoplus_{c=1}^{5} \mathrm{Cl}(Q_c)$ over Λ, on both sides of the level-one wall. All such categories are equivalent over Λ, so this pointwise statement says little. It is not an A_∞ functor along the path, and non-collision alone would not give one. Over Λ_0 the positive object changes across the level-one wall: its reduced Clifford form goes from rank 1 to rank 2. Across the level-two wall the path meets a degenerate critical point at $\gamma_c(T)$, where the cited theorems do not apply.




Figure 15: Along the path. (a) The number of real critical points at $L = 50$ and 100; the dotted lines are the zeros of the discriminant, one per L, at the level-two wall. (b) The small eigenvalue of Q/W at the positive object, against $h_2 - h_1$. (c) The positions of Stokes alignments at three phases and three values of L; the shaded bands are $|\gamma - \text{wall}| \le 1/L$.

Stokes alignments, separately. For the phases $\theta = 0.3$, 1.0 and $\pi/2$, and $L = 25, 50, 100$, every parameter where two critical values align, $\operatorname{Im} e^{-i\theta}(W_j - W_i) = 0$, lies within $1.5/L$ of one of the

two leximin walls, and at $L = 100$ within $1/L$; alignments occur near both walls (PP9; fig. 15(c)). This is the behaviour of proposition 35 on F_1, now near two walls at once. We do not claim it for every phase, nor that every leximin wall has such alignments nearby; near the level-two wall some alignments involve the pair born at the collision.

Part IV
Deeper selection: 2×4 couplings

13 Three-dimensional fibres

A 2×3 fibre is a polygon, so selection has at most two levels. A 2×4 fibre $\mathrm{Cpl}(f, g)$ is a three-dimensional polytope, and the leximin recursion can cut it down in up to three steps: a sheet, then a line, then a point. This part asks what such a deeper hierarchy does on the other side, and it keeps three questions apart. Is the chamber Fano, so that the Floer layer of section 4 applies? How many dimensions does each level remove? And does the Clifford form of the positive object see each level? The answers are, in order: only 14 of the 148 chambers are Fano, and in those coupled three-step selection never occurs (theorem 43 and proposition 44); a level can remove more than one dimension; and in every example the Clifford form has one scale per removed dimension (Conjecture 50, proved at leading order in three examples).

13.1 Coordinates and the potential

Use $u = (a_{11}, a_{12}, a_{13})$. The eight cells are $a_{1j} = u_j$ and $a_{2j} = g_j - u_j$ for $j \le 3$, $a_{14} = f_1 - \sum_j u_j$, and $a_{24} = \sum_j u_j - (f_1 + g_1 + g_2 + g_3 - 1)$. Their normals are $\pm e_j$ and $\mp(1,1,1)$. Over the facet cells (a non-facet cell contributes no term), the potential is

$$W = \sum_{j \le 3} \Big([1j]\, y_j + [2j]\, \frac{T^{g_j}}{y_j} \Big) + [14]\, \frac{T^{f_1}}{P} + [24]\, T^{g_4 - f_1} P, \qquad P = y_1 y_2 y_3, \tag{2}$$

where $[ij] = 1$ if cell ij is a facet and 0 otherwise. For each example we keep three counts separate: the vertices of the fibre (equal to $\chi(X)$), the critical points of W, and the distinct fibres that carry them.

13.2 Ampleness, certified by curves

Whether $X_{(f,g)}$ is Fano is a statement about its fan, and so about the chamber of (f, g). We certify it with integers on the edges of the polytope, by a classical criterion, rather than by any probabilistic argument.

Proposition 42 (ampleness by curves)**.** *Let X be a smooth complete toric threefold with polytope Q.*

(1) (Nakai–Moishezon) A divisor D on a projective variety is ample if and only if $D^{\dim V} \cdot V > 0$ for every irreducible subvariety V of positive dimension [33].

(2) (toric Kleiman) On a complete toric variety it suffices to test the torus-invariant curves: D is ample if and only if $D \cdot C > 0$ for each of them, and nef if and only if $D \cdot C \ge 0$ [14, Theorem 6.3.13 and §6.3].

(3) The invariant curves are the edges of Q. If an edge lies on the facets i, j and joins the vertices whose third facets are k and l, the wall relation $v_k + v_l + a\,v_i + b\,v_j = 0$ has integer a, b, and $D_k \cdot C = D_l \cdot C = 1$, $D_i \cdot C = a$, $D_j \cdot C = b$, and $D_\rho \cdot C = 0$ otherwise [14, §6.4]. So $-K \cdot C = 2 + a + b$, and the Kähler class meets C in $-\lambda_k - \lambda_l - a\lambda_i - b\lambda_j$, which must equal the lattice length of the edge.

On a toric variety $-K$ is always big, since its polytope $\{m : \langle m, v_\rho \rangle \geq -1\}$ contains 0 in its interior. So nef already means weak Fano.

Part (3) contains its own test: the lattice length of every edge must come out of the same integers that give $-K \cdot C$. TF2 confirms it on all 2208 edges of all chambers. The curves with $-K \cdot C = 0$ matter twice. They are what makes a chamber weak Fano rather than Fano. And they span the anticanonically trivial directions of the Mori cone, so they are the geometric source of the classes that *may* contribute to the corrections of W (remark 48); a zero degree does not by itself give a nonzero contribution.

Theorem 43 (the 2×4 chambers)**.** *The nondegenerate 2×4 margins fall into 148 Kähler chambers, the sign patterns of $f_1 - g(J)$ over the 14 nonempty proper sets J of columns. Every chamber polytope is simple and smooth. By proposition 42 they are of six types:*

type	*chambers*	*facets*	χ	$(-K)^3$	*curves with* $-K \cdot C = 0$	W	*Floer reading*	*drop patterns seen*
$\mathbb{CP}^3$	2	4	4	64	0	*as is*	*Cho–Oh*	(3)
$\mathrm{Bl}_{pt}\mathbb{CP}^3$	8	5	6	56	0	*as is*	*Cho–Oh*	(3), (1,2)
$(\mathbb{CP}^1)^3$	4	6	8	48	0	*as is*	*Cho–Oh*	(1,1,1), *decoupled*
weak Fano	24	6	8	48	1	$(1+\delta_k)$	*withheld*	(3), (1,2), (1,1,1)
weak Fano	64	7	10	40	3	$(1+\delta_k)$	*withheld*	(3), (1,2), (1,1,1)
weak Fano	46	8	12	32	6	$(1+\delta_k)$	*withheld*	(3), (1,2), (1,1,1)

Here χ is also the number of vertices, and "$(1+\delta_k)$" means the corrections of remark 48. In every chamber the normalised volume of $\mathrm{conv}\{v_c\}$ over the facets equals the number of vertices. The 148 are chambers of the margin hyperplane arrangement only: leximin walls subdivide them further, so the drop pattern is not a chamber invariant, and the last column lists only the patterns seen in a census.

Proof. An exact computation, TF1–TF3, with a machine-readable certificate, the file `chambers24.json` in `kahler/`, and an independent verifier (CC1–CC4). Each of the 148 chambers is certified by an exact rational interior point; that no other sign pattern is realised rests on linear-programming infeasibility. For each chamber, $-K \cdot C$ is computed on every edge from the integer wall relation, the type is read from the primitive relations, and an independent anticanonical vertex test agrees in all 148. The drop patterns are not invariants of a Kähler chamber, because leximin walls subdivide it; the last column lists those seen in a tie-free census (TF8). □

Proposition 44 (the box, and selection depth in the Fano chambers)**.** *(1) $\mathrm{Cpl}(f, g)$ is a box exactly when some column has $g_j > \max(f_1, f_2)$. Then column j never binds, $X = (\mathbb{CP}^1)^3$, W is a sum of three $\mathbb{CP}^1$ potentials, and the levels are the half-masses $g_k/2$ of the other three columns.*

(2) In a $\mathbb{CP}^3$ chamber selection is immediate: pattern (3). In a $\mathrm{Bl}_{pt}\mathbb{CP}^3$ chamber it is (3) or (1,2).

So coupled selection in three steps occurs only in weak Fano chambers.

Proof. (1) The column-j inequalities $0 \le f_1 - \sum_{k\neq j} a_{1k} \le g_j$ hold on the whole box $0 \le a_{1k} \le g_k$ exactly when $1 - g_j \le f_1 \le g_j$. Then the two cells of column j are not facets and their terms do not appear in (2). Each remaining column contributes $y + T^g/y$ in its own coordinate: y_k for a column $k \le 3$, and $z = T^{f_1}/P$ for column 4. These three coordinates are a unimodular basis. (2) By TF4 the facets of a $\mathbb{CP}^3$ chamber are one full row i, so the fibre is the simplex $\{a_{i\cdot} \ge 0,\ \sum_j a_{ij} = f_i\}$ and the equal split $f_i/4$ is selected at once. The facets of a $\mathrm{Bl}_{pt}\mathbb{CP}^3$ chamber are one full row i and one cell $i'j$, which cuts off a corner. If $f_i/4 < g_j/2$ the equal split is still selected at once. Otherwise column j binds first, at $g_j/2$, and the remaining three cells of row i share $f_i - g_j/2$ equally at the second level, a drop of two. □

A binding row finishes selection, since it leaves a candidate set of dimension $(n-2)(m-1) = 0$. The counting formula of theorem 23 gives only the first level, and the examples show that later levels can remove one or two dimensions. Write r_j for the number of dimensions removed at level j, the rank increment of the fixed normals; a level can also remove none, as the last level of example C does.




Figure 16: The three fibres and their candidate sets. Orange: the sheet left by the first level (a binding column). In A and C the second level leaves a line and the third a point; in B the second level is a binding row on what is left, and it fixes two dimensions at once. The star is the leximin point, the valuation of the positive critical point (theorem 27).

13.3 Three examples

Example 45 (A: the box)**.** Take $f = (2/5, 3/5)$ and $g = (1/20, 13/20, 1/10, 1/5)$. Column 2 never binds, so by proposition 44(1) the fibre is a box, $X = (\mathbb{CP}^1)^3$, and $W = (y_1 + T^{1/20}/y_1) + (y_3 + T^{1/10}/y_3) + (z + T^{1/5}/z)$ with $z = T^{2/5}/P$. The levels are 1/40, 1/20, 1/10, with drops $(1, 1, 1)$. The 8 critical points are $y_1 = \pm T^{1/40}$, $y_3 = \pm T^{1/20}$, $z = \pm T^{1/10}$, all over the leximin fibre $(1/40, 9/40, 1/20)$, with leading holonomies ± 1. Their critical values are $2(\pm x \pm x^2 \pm x^4)$ with $x = T^{1/40}$. For $T \in (0, 1)$ exactly two coincide, $(+, -, -)$ and $(-, +, +)$, at the root of $x^3 + x - 1 = 0$, $T^* \approx 2.3 \times 10^{-7}$. There the m_0 argument for vanishing Hom does not apply (the holonomies still differ), and we do not compute that group. TF5 checks all of this exactly.

Proposition 46 (B: the blown-up projective space, exactly)**.** *Take* $f = (3/20, 17/20)$ *and* $g = (1/20, 1/5, 1/4, 1/2)$*. The facets are row* 1 *and cell* 21*, so* $X = \mathrm{Bl}_{pt}\mathbb{CP}^3$ *and* $W = y_1 + y_2 + y_3 + T^{3/20}/P + T^{1/20}/y_1$*. Put* $\tau = T^{1/20}$*.*

(1) Selection. *Column* 1 *binds at* $h_1 = 1/40$*, and row* 1 *then spreads* $1/8$ *over three cells,* $h_2 = 1/24$*. The drops are* $(1, 2)$*.*

(2) Reduction. *At a critical point* $y_2 = y_3 = c$ *and* $y_1 = \tau^3/c^3$*, where* $c^6 + \tau^2 c^4 - \tau^5 = 0$*. As a cubic in* c^2 *this has discriminant* $\tau^{10}(4\tau - 27)$*, so there are* 6 *distinct critical points for* $T \in (0, 1)$*.*

(3) One fibre. *The Newton polygon of the cubic has a single edge. All six critical points lie over the leximin fibre* $(1/40, 1/24, 1/24)$*, with* $c = \zeta T^{1/24}(1 + o(1))$ *and* $y_1 = \zeta^{-3}T^{1/40}(1 + o(1))$*,* $\zeta^6 = 1$*.*

(4) Distinct values. *The critical values* $w = 2y_1 + 2c$ *are the roots of a sextic whose discriminant is* $2^{36}\tau^{19}(4\tau - 27)^6(4\tau + 1)^2$*. So they are distinct for every* $T \in (0, 1)$*.*

(5) Objects and morphisms *(Fano; cited, in the convention of section 12.7). The six are objects over one fibre, with distinct* m_0*, so every off-diagonal Hom vanishes and the category they generate is* $\bigoplus_{c=1}^{6} \mathrm{Cl}(Q_c)$ *over* Λ*, with no higher products. Generation is conditional.*

Proof. (1) By proposition 44(2), since $f_1/4 = 3/80 > g_1/2 = 1/40$. (2) The equations $y_j\partial_j W = 0$ give $y_2 = y_3 = T^{3/20}/P = c$ and $y_1 - \tau/y_1 = c$; eliminating gives the sextic, and the discriminant follows. (3) The lower hull of $(3, 0), (2, 2), (0, 5)$ in the plane of (power of c^2, power of τ) is the single segment from $(0, 5)$ to $(3, 0)$, of slope $-5/3$, so $c^2 \sim \tau^{5/3}$ and $c \sim \tau^{5/6} = T^{1/24}$. Then $y_1 = \tau^3/c^3$. (4) A resultant. (5) Proposition 31(2)–(3) and proposition 32, with (4). TF6 checks (2)–(4) symbolically. □

Example 47 (C: weak Fano, coupled in three steps)**.** Take $f = (3/10, 7/10)$ and $g = (1/10, 3/20, 1/4, 1/2)$. The facets are row 1 and cells $21, 22, 23$, so seven facets, and the chamber is weak Fano with three curves of $-K \cdot C = 0$. Here $W = \sum_{j\le 3}(y_j + T^{g_j}/y_j) + T^{3/10}/P$. Columns 1 and 2 bind in turn, at $h_1 = 1/20$ and $h_2 = 3/40$. Row 1 then spreads $7/40$ over cells 13 and 14, at $h_3 = 7/80$. The last cell, 23, is fixed at $13/80$ without removing a dimension. The drops are $(1, 1, 1, 0)$. The critical equations reduce to one resultant of degree 10 in $c = T^{3/10}/P$, even in c. Its Newton polygon in c^2 has an edge of slope $7/2$ with four roots and one of slope 1 with one root. So 8 critical points lie over the leximin fibre $(1/20, 3/40, 7/80)$, with leading holonomies of the form $(\pm 1, \pm 1, \pm 1)$ or $(\pm 1, \pm 1, \pm i)$, and 2 lie over $(1/40, 1/40, 9/40)$, with holonomies $\pm(i, i, i)$. On that second fibre the leading set is $\{11, 12, 14, 23\}$, with leading coefficients $\rho = (i, i, i, -i)$ up to conjugation. (H1) holds and (H2) fails, yet $\operatorname{rank} M_c = 3 = \dim \operatorname{span}$: two more instances of the equality in theorem 38 without its sufficient condition. TF7 and TF9 check this.

Remark 48 (what changes in a weak Fano chamber)**.** For a semi-Fano toric manifold, that is, one with $-K$ nef, the potential of a fibre is $W^{\mathrm{LF}} = \sum_k (1 + \delta_k)\, T^{-\lambda_k} y^{v_k}$, where $\delta_k = \sum_{\alpha \neq 0} n_{\beta_k + \alpha} T^{\omega(\alpha)}$ runs over the classes α of rational curves with $-K \cdot \alpha = 0$ [13, Proposition 2.15]. The monomials are the same and only the coefficients change, by factors $1 + O(T^{\varepsilon})$. Three consequences, and one thing withheld:

(i) The valuations of the terms are unchanged, so the levels h_j, the leximin point and the leading coefficients are unchanged.

(ii) Theorem 27 needs real coefficients with uniform bounds. If the δ_k converge for small T, are real, and satisfy $0 < c \leq 1 + \delta_k(e^{-L}) \leq C$ for large L, then it applies to W^{LF} as it stands: the corrected positive critical point is selected at the leximin point, with error $O(1/L)$. If the δ_k are only formal Novikov series, the conclusion holds as a statement about valuations, by (i), and not about a real critical point.

(iii) The leading-order argument of proposition 51 uses only valuations and leading coefficients, so it applies to W^{LF} verbatim (argued, not checked).

(iv) Withheld: the exact critical points of W^{LF}, their number over each fibre, and the Floer reading of the ten objects. We do not compute the δ_k.

For each curve with $-K \cdot C = 0$ four statuses must be kept apart: zero anticanonical degree (computed here, exactly); an effective class α through it that contributes to some δ_k (not determined); a nonzero invariant $n_{\beta_k+\alpha}$ (not computed); the correction evaluated (not done).



Figure 17: Every critical point of the three potentials, grouped by fibre, with its leading holonomies drawn as dials (the arrow is the phase of y_j). A and B have one fibre each. C has two: eight points over the leximin fibre and two over $(1/40, 1/40, 9/40)$. In each case the number of critical points equals the normalised volume of the Newton polytope, which equals χ, and that equality certifies the list complete. For C these are the critical points of the uncorrected W (remark 48).

13.4 One scale per removed dimension

Theorem 33 says that the reduction of Q/W mod T has rank r_1. The examples say more: the other directions of Q/W have exact scales, one for each dimension a later level removes. To state this without ordering eigenvalues, use determinantal divisors.

Definition 49 (scale valuations)**.** For a matrix M over Λ_0, let d_k be the least valuation of a $k \times k$ minor, and $e_k = d_k - d_{k-1}$, with $d_0 = 0$. The e_k are the valuations of the invariant factors of M. They do not change under $M \mapsto AMB$ with $A, B \in \mathrm{GL}_n(\Lambda_0)$, in particular under a change of integral basis of $H^1(L;\mathbb{Z})$.

For the positive object, Q/W is real symmetric and positive definite, and its scale valuations are the valuations of its eigenvalues. The sum of the $k \times k$ principal minors is the k-th elementary symmetric function of the eigenvalues; all terms are positive, so its valuation is that of the product of the k largest eigenvalues. And for a positive definite matrix $|\det M_{IJ}|^2 \le \det M_{II} \det M_{JJ}$, so the least valuation over all $k \times k$ minors is attained on a principal one.

Conjecture 50 (one scale per removed dimension)**.** *For nondegenerate margins, the scale valuations of Q/W at the positive object are 0 with multiplicity r_1, and $h_j - h_1$ with multiplicity r_j for each later level j. A level that removes no dimension contributes nothing.*

Proposition 51 (the conjecture in the three examples)**.** *At leading order, the scale valuations of Q/W are:*

(A) $(0,\ 1/40,\ 3/40)$ *for all* 8 *objects;*

(B) $(0,\ 1/60,\ 1/60)$ *for all* 6 *objects: the second level removes two dimensions and gives a double scale;*

(C) $(0,\ 1/40,\ 3/80)$ *for the* 8 *objects over the leximin fibre, with no scale from the fourth level; and* $(0,0,0)$ *for the two over the second fibre.*

So Conjecture 50 holds in A, B and C.

Proof. In every case $Q = \sum_k (\text{term } k)\, v_k v_k^\top$, and at a critical point $y_j - T^{g_j}/y_j = c$, where c is the value of the P-term. (A) In the unimodular basis $(\log y_1, \log y_3, \log z)$, $Q = \mathrm{diag}(W_1, W_3, W_z)$, with each factor's value $\pm 2T^h$ and $W \sim \pm 2T^{1/40}$. (B) $Q = \left(\begin{smallmatrix} 2y_1 & c & c \\ c & 2c & c \\ c & c & 2c \end{smallmatrix}\right)$ and $W = 2y_1 + 2c$, with $y_1 \sim T^{1/40}$ and $c \sim T^{1/24}$ (proposition 46). So $d_1 = 0$. The 2×2 minors have least valuation $1/40 + 1/24$, attained by $4y_1 c - c^2$, so $d_2 = 1/24 - 1/40 = 1/60$. Next, $\det Q = 6y_1c^2 - 2c^3 \sim T^{1/40+1/12}$, so $d_3 = 1/30$, and $e = (0, 1/60, 1/60)$. (C) $Q = D + cJ$, with J the all-ones matrix and $D = \mathrm{diag}(2y_j - c)$. On the leximin fibre $y_1 \sim \pm T^{1/20}$, $y_2 \sim \pm T^{3/40}$ and $c \sim y_3 \sim T^{7/80}$, so $D \approx \mathrm{diag}(2y_1, 2y_2, y_3)$ and $W \approx 2y_1$. The 2×2 minors have least valuation $1/20 + 3/40$, so $d_2 = 1/40$. Next, $\det Q = D_1D_2D_3 + c\,(D_1D_2 + D_1D_3 + D_2D_3) \approx 2y_3D_1D_2 \sim T^{17/80}$, with no cancellation, so $d_3 = 17/80 - 3/20 = 1/16$, and $e = (0, 1/40, 3/80)$. The second fibre has rank 3 by example 47. TF9 checks every case numerically: the eigenvalue slopes at the positive object, and the minors on every object over the leximin fibre. These finite-T checks corroborate the limiting valuations; the proof is the leading-order computation above. □

In 2×3 the evidence was single scales: $(3\alpha - 1)/4$ on F_1, $1/200$ after one count, and $1/20$ in the column-binding example (section 12). The cube proves the conjecture for every box chamber, because the factors decouple. B is the first coupled case with a multiplicity, and C the first coupled case with three levels and a level that removes nothing. A proof in general would need the successive balances of theorem 27 to appear, one order at a time, in the minors of Q. That is the step we have not found.




Figure 18: The eigenvalues of Q/W at the positive object, as $-\log|\mu|/L$ against $1/L$ for $L = 50, \ldots, 1600$. The finite-T offset is linear in $1/L$, so the curves run straight into their limits: 0 once, then $h_j - h_1$ for each removed dimension. In B the second level removes two dimensions and two eigenvalues share the limit $1/60$. In C the fourth level, $13/80$, removes none and leaves no trace.

14 Outlook

Seven questions are left open.

- *Novikov weights and environments.* Section 6 makes one direction precise: soft evidence of size n on the scale N shifts λ by $-(n/N)\rho$. Whether the environment orders s^{r_e} of the layers paper act in the same way, and how hard evidence and forgetting act, is open.
- *The other walls.* Decision, contextuality and value-of-information walls (section 6), and the finite-n sharpness of the transition at an outcome wall, measured in the Fisher metric.
- *Beyond Fano.* For non-Fano toric manifolds, W acquires corrections [23]. Theorem 3 is about the leading potential only. In the weak Fano chambers of 2×4 the corrections only rescale coefficients (remark 48); computing the δ_k, and so the corrected objects of example C, is open.
- *One scale per removed dimension.* Conjecture 50 in general; then 3×3, where the fibre is four-dimensional, as a test of it and of the rank equality in theorem 38.
- *Signed branches and Stokes data.* The non-positive critical points have their own thimbles, which are not real and carry no probability law. Proposition 35 locates the Stokes walls on F_1; the Picard–Lefschetz decomposition itself, and the general window, are not computed. We do not know whether the signed fibre of example 6(a) has a probabilistic shadow. Section 12.8 crosses two leximin walls from the column-binding example; what the collision at the level-two wall does to the category, and an A_∞ functor along a path that avoids it in complexified parameters, are open.
- *The integral category.* Theorem 33 sees the outcome wall in the reduction mod T of one endomorphism algebra. Whether the full A_∞ category over Λ_0 changes there, and split generation in the non-monotone case [40], are open.
- *Wall-crossing.* The layers paper resolves degenerate directions by blow-ups off the torus orbits. Whether these correspond to wall-crossing in torus fibrations is a separate question, which needs geometric data that the finite setting does not supply.

A The ledger

`LEDGER.yaml` at the root of the repository is the single source for this table, which `kahler/render_ledger.py` generates. Status tags: [P] proved here, [U] used (classical), [C] cited and not proved, [E] exact computation, [N] numerical only, [D] a definition, [O] open. The category level is never filled by assumption. For each experiment the ledger also records the thresholds $s_{\text{prob}}, s_{\text{support}}, s_{\text{trop}}, s_{\text{crit}}, s_{\text{Stokes}}, s_{\text{context}}$, each as a number, "never" or "undefined". For F_1 along $\rho = e_1$ from $\alpha = 1/5$ they are: $s_{\text{trop}} = 1/5$; s_{crit} never; s_{support} never; s_{prob}, s_{Stokes} and s_{context} undefined until a dominance notion, a phase of $\hbar$, and a complex with an obstruction class are chosen.

operation	contingency / toric	mirror (W, λ)	Floer (cited); category	wall certificate; checks
soft-translation: Soft evidence along a translation, rho_k = ⟨v_k, t⟩ (an exponential tilt)	[D] reweighting by a log-linear factor: a reparametrisation of the same model; [P] Q translated by -s t; same Kahler class	[P] u* → u* - s t; h*, leading set and candidates unchanged	[C] the same fibres, translated; category [O] not addressed	none: rho is orthogonal to B - B (Corollary 12(2)); CHECKS DW7
soft-constant: Soft evidence along the constant, rho = 1 (normalisation)	[D] a common factor: removed by normalisation; [P] every facet moves out by s	[P] h* → h* + s; nothing else changes	[C] no change in which fibres carry Floer cohomology; category [O] not addressed	none: sum_k a_k = 0 for every affine dependence a; CHECKS DW7
soft-circuit: Soft evidence with a circuit component (lam_eff = lam - (n/N) rho)	[E] independence: a larger row or column survey (L1 = n1, L2 = n2); 2x3 needs n/2 more column observations to reach its wall; [P] moves the Kahler class; viability walls are the walls of the Kahler cone (F_1: alpha/L in 0, 1)	[P] outcome walls = normal fan of the balance polytope B; leading set, candidate dimension and face F jump there; h* concave piecewise linear	[C] F_1: two fibres with HF ≠ 0 below alpha/L = 1/3, one above (Cho-Oh); independence: one fibre on both sides; category [C] F_1: over Lambda the category of the four objects is unchanged up to A-infinity equivalence across the wall (on cited m_2); the positive object's Clifford form mod T drops rank 2 → 1; Stokes walls crowd the wall (section 12)	one circuit a: s* = -c(lam_0) / ⟨a, rho⟩; least s over rate range ≤ 1 is |c| / (||a||_1 / 2). F_1: a = (2,-1,2,-3), c = 3 alpha - L; independence: a = (c 1_r, -r 1_c)/gcd, c(lam) = (c L1 - r L2)/gcd; CHECKS DW2, DW4, DW5, DW6, DW7, CT2, CT6, MO1, MO6, MO7, MO8, MO9
margins: Observed counts (margins b)	[U] the MLE has the observed margins (Birch); independence: rows x cols / n^2; [U] the fibre over b; l_k(b) is the k-th margin (independence: exact)	[P] u* does not depend on b while lam is fixed	[D] compare L(b) with the Floer-selected L(u*), in Fisher-Rao distance within the model; category [O] not addressed	support: min_k l_k(b) = 0; centrality g(b)/h* = smallest margin / (n / max(r, c)) for independence with joint sampling; CHECKS CT4, CT5

operation	contingency / toric	mirror (W, λ)	Floer (cited); category	wall certificate; checks
margins-on-the-fibre: Margins (f, g) read on the fibre, the coupling polytope Cpl(f, g)	[E] the fibre's lattice points are the tables with these margins; Tier A is Fisher's noncentral hypergeometric family, odds ratios 1 at f(x)g; [U] Cpl(f, g) is Delzant; class = product coupling; 2x3: CP^2, CP^1 x CP^1, F_1, Bl_2 CP^2, dP_6 (coupling-polytopes paper)	[P] h* = min(f_i/m, g_j/n); leading cells = the binding row or column; candidate dimension (n-2)(m-1) or (n-1)(m-2); the selected coupling is the facet-leximin one (selection theorem), error O(1/log(1/T))	[C] example: three fibres with HF ≠ 0; one count moved: three become two (valuations by exact resultants); category [C] one count: objects 3 + 1 + 1 → 4 + 1 over the fibres; the positive object's Clifford form mod T drops rank 2 → 1 (section 12). Column-binding example: 5 critical points, 2 fibres, 5 objects (exact, all T); Hom = 0 off the diagonal, End = Cl(Q_c), all reduce to rank 1 (section 12.7; FOOO Lambda_0 convention)	the argmin of f_i/m, g_j/n changes. Example: f1/3 = 2/15 vs g3/2 = 27/200 (1/600 apart): one count crosses; the blow-up wall f1 = g2 is three counts away; CHECKS CF1, CF2, CF3, CF4, CF5, CF6, CF7, CF8, LX1, LX4, LX5, LX6, LX9, MO2, MO6, CB1, CB3, CB4, CB7, CB8
deeper-selection: Margins on a 2 x 4 fibre: a three-dimensional coupling polytope, selection in several levels	[E] the leximin coupling level by level: binding columns, then a residual row; a level can fix one dimension, two, or none (examples A, B, C); [E] 148 chambers, all smooth: 14 Fano (CP^3, Bl_pt CP^3, (CP^1)^3) and 134 weak Fano, certified by -K.C on every invariant curve (toric Kleiman); coupled three-step selection only in weak Fano chambers	[E] A: 8 critical points, 1 fibre; B: 6, 1 fibre, distinct values for all T; C: 10 over 2 fibres (8 + 2). Clifford scales 0 and h_j - h_1 with multiplicity r_j at leading order in A, B, C; in general a conjecture	[C] A, B Fano: Cho-Oh objects, Hom = 0 off the diagonal by m_0 (B for every T; A except at one T*). C weak Fano: W gets (1 + delta_k) corrections (Chan-Lau-Leung-Tseng); its Floer reading is withheld; category [C] B: the category of its six objects is a sum of Clifford algebras over Lambda (on cited theorems); weak Fano chambers not addressed	chamber: the sign vector of f1 - g(J); Fano: min over invariant curves of -K.C ⟩ 0; box: some g_j ⟩ max(f1, f2); CHECKS TF1, TF2, TF3, TF4, TF5, TF6, TF7, TF8, TF9, CC1, CC2, CC3, CC4
path-across-walls: A path of margins inside one Kahler chamber (Stage IV): f = (2/5, 3/5), g = (7/10 - gamma, 3/10, gamma), gamma in [3/20, 11/40]	[E] the leximin coupling changes at gamma = 1/5 (level 2: cells 11, 12, 22 tie; the column-binding example lies there) and at 4/15 (level 1: row 1 starts to bind); [E] one Kahler chamber (the pentagon) along the whole path: no blow-up wall	[E] objects over fibres 4 + 1 below 1/5, 2 + 3 at 1/5, 4 + 1 between, 3 + 1 + 1 above 4/15 (Newton polygon); a collision of two real critical points at gamma_c(T) → 1/5 (numerical: one discriminant zero at each L tested), none at 4/15; the positive object's Clifford scale h2 - h1 bends at 1/5 and closes at 4/15	[C] away from gamma_c(T) the critical values are distinct, so Hom = 0 off the diagonal and End = Cl(Q_c) at every point (cited theorems); at gamma_c a critical point is degenerate and the cited theorems do not apply; category [O] pointwise the same category over Lambda on both sides of the level-1 wall; no A-infinity functor along the path is constructed; the effect of the collision is open	level 2: f1 - g3/2 = g2; level 1: g3/2 = f1/3; collision: a sign change of the discriminant of the quintic; CHECKS PP1, PP2, PP3, PP4, PP5, PP6, PP7, PP8, PP9
refinement: Refinement: add a distinction (a facet normal, a toric blow-up)	[O] which table operation adds a facet is not yet identified; [U] CP^2 → F_1 is the blow-up at a point	[E] creates the circuit (2,-1,2,-3) and the wall alpha/L = 1/3; CP^2 had none	[C] F_1 gains a second fibre with HF ≠ 0 below the wall; category [O] not addressed	m - n - 1 increases by one; CHECKS DW4

operation	contingency / toric	mirror (W, λ)	Floer (cited); category	wall certificate; checks
structural-zero: Hard evidence: a structural zero (support restriction)	[E] 2x2 with one structural zero: quasi-independence on three cells; [E] the unit square becomes a unimodular triangle, not a face of it: CP^1 x CP^1 is replaced by CP^2	[E] the one circuit and the one wall disappear	[C] CP^2: one fibre (the Clifford torus); category [O] not addressed	face test fails; DW circuits of the triangle: none; CHECKS CT7
independent-product: Independent product of two models	[E] independence of rows and columns: Delta_(r-1) x Delta_(c-1); [U] the product manifold	[P] adds exactly one circuit, the relative scale (q1, -q2); u* = (u1*, u2*) never moves	[C] independence: exactly one fibre with HF $\neq 0$ for every L1, L2, so this wall changes no Floer data; category [O] Kunneth-type statements not addressed	m - n - 1 = (m1 - n1 - 1) + (m2 - n2 - 1) + 1; CHECKS CT1, CT2, CT3
forgetting: Forgetting an environment / marginalisation	[P] the layers paper: the forgetting square, reversal walls, pooled orders; [O] the image configuration under a linear map of exponents: not yet worked out	[O] not yet worked out	[O] not addressed; category [O] not addressed	layers paper, theorem "contextual Bayes with an environment"; CHECKS EN1, EN2, EN3
coupling: Dependent coupling, convolution, intervention, Markov evolution	[O] not yet worked out; [O] convolution would be a Minkowski sum of supports	[O] not yet worked out	[O] not addressed; category [O] not addressed	-; CHECKS –

Checks

code	`engine/pce/kahlerbridge.py`, `engine/pce/datawalls.py`, `engine/pce/contingency_toric.py`, `engine/pce/couplingfibre.py`, `engine/pce/floermorph.py`, `engine/pce/colbind.py`, `engine/pce/cpl24.py`, `engine/pce/pentpath.py`
checks	`engine/check_kahlerbridge.py` (KB, 8/8), `engine/check_datawalls.py` (DW, 7/7), `engine/check_contingency_toric.py` (CT, 7/7), `engine/check_couplingfibre.py` (CF, 8/8), `engine/check_leximin.py` (LX, 9/9), `engine/check_floermorph.py` (MO, 9/9), `engine/check_colbind.py` (CB, 8/8), `engine/check_cpl24.py` (TF, 9/9), `engine/check_pentpath.py` (PP, 9/9), `engine/check_chambers24_cert.py` (CC, 4/4), `engine/check_ledger.py` (LG, 3/3)
figures	`kahler/fig_kahler.py`, `kahler/fig_datawalls.py`, `kahler/fig_coupling.py`, `kahler/fig_leximin.py`, `kahler/fig_morph.py`, `kahler/fig_colbind.py`, `kahler/fig_cpl24.py`, `kahler/fig_pentpath.py`
certificate	`kahler/chambers24.json`
ledger	`LEDGER.yaml`, rendered by `kahler/render_ledger.py`

The checks corroborate the statements numerically, or by exact rational arithmetic where marked; they do not prove them. The proofs stand on their own. The F_1 blow-up sizes used are $\alpha = 1/5, 1/4, 2/5, 3/5, 4/5$.

KB1	proposition 1(1)–(2): moment map $= \nabla\varphi$, Fisher information $= \operatorname{Hess}\varphi$, positive definite (30 supports)
KB2	proposition 1(4): $\psi = \sum q \log(q/c)$, $\nabla\psi = u(p)$, $\operatorname{Hess}\psi = \text{Fisher}^{-1}$ (20 supports)
KB3	proposition 1(5): $\int \det \operatorname{Hess}\varphi$ over boxes equals the area of the moment image

KB4	theorem 3(1)–(2): one positive critical point, mean-zero law, the valuations of examples 5 and 6, balanced leading terms
KB5	theorem 3(3): the segment of candidates for $\alpha = 3/5$, and its endpoints $(1/5, 1/5)$, $(3/5, 1/5)$, with leading sets $\{v_1, v_2, v_4\}$, $\{v_2, v_3, v_4\}$ and no positive balance; the barycentre alone for $\alpha = 1/5$ and for $\mathbb{CP}^2$ (exact rational points)
KB6	theorem 3(4): all four critical points of F_1 in 80-digit arithmetic; the signed critical point and its signed balance
KB7	proposition 7: Hessian $\operatorname{diag}(H, -H)$ at y_T; Laplace covariance $\hbar H^{-1}$ with relative error $O(\hbar / W_T(y_T))$ ($\mathbb{CP}^2$, F_1 with $\alpha = 3/5$)
KB8	theorem 8, example 9: moments against the uniform law on Q_h for $h < h^*$; concentration on $\operatorname{argmax} g$ at h^*; the uniform segment; the levels $3/10$, $9/20$; $\log T = -200, -800$
DW1	definition 10: evidence $e^{-n\rho}$ at $T = e^{-N}$ is the shift $\lambda - (n/N)\rho$
DW2	proposition 11(1)–(2): duality; optimal support = leading set at the centre of argmax; $\dim \operatorname{argmax} = n - \dim \sigma_0$ (40 random configurations)
DW3	proposition 11(3): the limit law lies in the optimal face (23 cases)
DW4	example 14(a): the circuit $(2, -1, 2, -3)$; the wall $\alpha/L = 1/3$ inside $0 < \alpha/L < 1$; the diagonal flip (exact rationals); $\mathbb{CP}^2$ one chamber
DW5	example 14(b),(c): candidates and valuations for $\alpha = 1/10, \dots, 9/10$; the wall law from $x^4 = x + 1$ at $T = 10^{-20}, 10^{-60}$
DW6	example 14(d): the discriminant, symbolically; its terms tie only at $\alpha = 1/3$
DW7	corollary 12, example 14(e): $s^* = 1/5$ for $\rho = e_1$; no wall for e_4, $(0, -1, 0, 1)$, translations or constants; least evidence $1/10$; $\mathbb{CP}^2$ never
CT1	proposition 15: $(r-1)(c-1)$ moves; one facet circuit ($r, c \in \{2, 3, 4\}$)
CT2	theorem 17(1): h^*, the leading sets and the wall $cL_1 = rL_2$ (exact rationals)
CT3	theorem 17(2)–(3): the rc critical points and their common valuation; the exact positive law; candidate dimensions
CT4	proposition 18(1): $\ell_k(b)$ = margins; MLE margins and positivity (200 tables)
CT5	proposition 18(3): the block-diagonal Fisher information; properties of d_Δ
CT6	example 19: $h^* = n/3$, the candidate segment, u^*, and $n/2$ more column observations
CT7	example 20: the triangle is not a face; no circuit
CF1–CF2	example 21: the pentagon, its facets, Delzant; 715 tables (Pick), exact
CF3	proposition 22: odds-ratio identity; exact hypergeometric mean $(12, 86/5)$; Birch at the leximin
CF4	theorem 23(1)–(3): the closed form against the facet programme; binding cells are facets; candidate dimensions
CF5	example 24: five critical points by exact resultants; positions, signs, leading sets, signed balances
CF6	theorem 23(4): positive valuation = leximin coupling (20 random margins)
CF7	fig. 5: the thimble law against uniform on Q_h; concentration at the leximin
CF8	example 25: one count; the walls' distances; three valuations become two
LX1–LX3	theorem 27: $L\lvert u_L - u^{\text{lex}}\rvert$ bounded on 30 random configurations (multiprecision Newton); the projected equation; $\mathbb{CP}^2$, F_1
LX4	corollary 29: facet-leximin = all-cell leximin (292 margins); convergence
LX5–LX9	example 30: the fibre and its two levels (exact); the $\ln 2/2$ correction; the five critical points; the thimble; the one-count levels
MO1–MO2	the objects of F_1 and of the 2×3 fibres: counts, fibres, types, holonomies
MO3–MO5	proposition 31: disjoint fibres, distinct holonomies and values; $Q = W \operatorname{Cov}_w(v)$; Clifford generators spanning M_2 (18 objects)
MO6	theorem 33: $Q/W \to \operatorname{Cov}_{q^*}(v)$; reduced ranks $2, 2, 1, 1$ on F_1; $2 \to 1$ after one count; valuations of the small eigenvalue
MO7	theorem 34: no collision, Morse along $\alpha \in [1/5, 3/5]$; explicit real isomorphisms
MO8–MO9	proposition 35: Stokes walls within $4/\log(1/T)$ of the wall, none outside, rate $1/\log(1/T)$
CB1–CB4	proposition 36: the reduction to the quintic; the Newton polygon; distinct and nondegenerate for all $T > 0$; distinct critical values on $(0, 1)$ (exact, symbolic)

CB5–CB6	the table of section 12.7: leading coefficients, offsets $-\log\lvert\rho\rvert/L$, leading values; leading Clifford forms and determinants
CB7	proposition 39: 5 objects, 2 fibres; distinct values and holonomies; $\mathrm{Cl}(Q_c) \cong M_2$
CB8	theorem 38: (H1), convergence to M_c, rank = span on 23 objects
PP1–PP2	proposition 40(1): one Kähler chamber along the path; the leximin levels and the two walls (exact)
PP3–PP4	proposition 40(2)–(3): the reduction to (1); Newton polygons at 10 rational γ (exact)
PP5	proposition 40(3): valuations, real and complex types, leading phases at $L = 200, 400$
PP6	proposition 41(1)–(2): one discriminant zero, at the level-two wall, for $L = 50, \ldots, 400$; none at the level-one wall
PP7	proposition 41(3): no coincidence of critical values away from the collision
PP8	proposition 41(4): the positive object's scale against $h_2 - h_1$ along the path
PP9	Stokes alignments at three phases and three L, within $1.5/L$ of a wall
TF1	theorem 43: 148 chambers, each with an exact interior point; all simple and smooth
TF2	proposition 42(3): integer wall relations on all 2208 edges; the Kähler degree equals the lattice length (exact)
TF3	theorem 43: the six types by toric Kleiman; the vertex test agrees; volume = χ; $(-K)^3 = 64, 56, 48, 48, 40, 32$
TF4	proposition 44: the box criterion on all chambers; the facets of the $\mathbb{CP}^3$ and $\mathrm{Bl}_{pt}\mathbb{CP}^3$ chambers
TF5	example 45: 8 critical points over one fibre; the one coincidence of critical values at $x^3 + x - 1 = 0$ (exact)
TF6	proposition 46(2)–(4): the sextic, its discriminants, one fibre, holonomies $(\zeta^{-3}, \zeta, \zeta)$ (exact)
TF7	example 47: degree-10 resultant, its Newton polygon, $8+2$ critical points over two fibres; three curves with $-K \cdot C = 0$
TF8	propositions 44 and 46 and examples 45 and 47: the leximin levels and drops exactly; a tie-free census of drop patterns by chamber type
TF9	proposition 51: eigenvalue slopes at the positive object ($L = 800 \to 1600$); minors on all 22 leximin-fibre objects; rank 3 on C's second fibre
CC1–CC4	theorem 43: the certificate `kahler/chambers24.json`, verified independently (standard library, exact rationals)
LG1–LG3	appendix A: the schema; every cited check passes in the log; the rendered table matches `LEDGER.yaml`

References

[1] M. Abreu, Kähler geometry of toric manifolds in symplectic coordinates, in *Symplectic and Contact Topology*, Fields Inst. Commun. **35**, AMS, 2003, 1–24.

[2] S. Amari and H. Nagaoka, *Methods of Information Geometry*, AMS and Oxford University Press, 2000.

[3] D. Auroux, Mirror symmetry and T-duality in the complement of an anticanonical divisor, *J. Gökova Geom. Topol.* **1** (2007) 51–91.

[4] V. K. Awasthi, Bridges: from probability and entropy to branes, with worked examples, technical note, Johns Hopkins University, October 2026.

[5] V. K. Awasthi, Coupling polytopes and toric geometry: the 2×3 family, the square case, higher levels and stability, companion paper, Johns Hopkins University, October 2026.

[6] V. K. Awasthi, How the programme fits together: one family of polytopes, many models, and one example through every layer, draft 1, Johns Hopkins University, October 2026.

[7] V. K. Awasthi, Probability, context and differentiation: the layers, working draft, 2026.

[8] M. W. Birch, Maximum likelihood in three-way contingency tables, *J. Roy. Statist. Soc. Ser. B* **25** (1963) 220–233.

[9] Y. M. M. Bishop, S. E. Fienberg and P. W. Holland, *Discrete Multivariate Analysis*, MIT Press, 1975.

[10] J. Bochnak, M. Coste and M.-F. Roy, *Real Algebraic Geometry*, Ergebnisse der Mathematik **36**, Springer, 1998.

[11] C.-H. Cho and Y.-G. Oh, Floer cohomology and disc instantons of Lagrangian torus fibers in Fano toric manifolds, *Asian J. Math.* **10** (2006) 773–814.

[12] C.-H. Cho, Products of Floer cohomology of torus fibers in toric Fano manifolds, *Comm. Math. Phys.* **260** (2005) 613–640.

[13] K. Chan, S.-C. Lau, N. C. Leung and H.-H. Tseng, Open Gromov–Witten invariants and mirror maps for semi-Fano toric manifolds, *Pure Appl. Math. Q.* **16** (2020) 675–720.

[14] D. A. Cox, J. B. Little and H. K. Schenck, *Toric Varieties*, Graduate Studies in Mathematics **124**, American Mathematical Society, 2011.

[15] R. Cominetti and J. San Martín, Asymptotic analysis of the exponential penalty trajectory in linear programming, *Math. Programming* **67** (1994) 169–187.

[16] T. Delzant, Hamiltoniens périodiques et images convexes de l'application moment, *Bull. Soc. Math. France* **116** (1988) 315–339.

[17] P. Diaconis and B. Sturmfels, Algebraic algorithms for sampling from conditional distributions, *Ann. Statist.* **26** (1998) 363–397.

[18] J. J. Duistermaat and G. J. Heckman, On the variation in the cohomology of the symplectic form of the reduced phase space, *Invent. Math.* **69** (1982) 259–268.

[19] T. Dyckerhoff, Compact generators in categories of matrix factorizations, *Duke Math. J.* **159** (2011) 223–274.

[20] N. Eriksson, S. E. Fienberg, A. Rinaldo and S. Sullivant, Polyhedral conditions for the nonexistence of the MLE for hierarchical log-linear models, *J. Symbolic Comput.* **41** (2006) 222–233.

[21] S. E. Fienberg and A. Rinaldo, Maximum likelihood estimation in log-linear models, *Ann. Statist.* **40** (2012) 996–1023.

[22] R. A. Fisher, The logic of inductive inference, *J. Roy. Statist. Soc.* **98** (1935) 39–82.

[23] K. Fukaya, Y.-G. Oh, H. Ohta and K. Ono, Lagrangian Floer theory on compact toric manifolds I, *Duke Math. J.* **151** (2010) 23–174.

[24] W. Fulton, *Introduction to Toric Varieties*, Annals of Mathematics Studies **131**, Princeton University Press, 1993.

[25] S. Galkin, The conifold point, arXiv:1404.7388, 2014.

[26] S. Galkin, V. Golyshev and H. Iritani, Gamma classes and quantum cohomology of Fano manifolds: Gamma conjectures, *Duke Math. J.* **165** (2016) 2005–2077.

[27] D. Geiger, C. Meek and B. Sturmfels, On the toric algebra of graphical models, *Ann. Statist.* **34** (2006) 1463–1492.

[28] I. M. Gelfand, M. M. Kapranov and A. V. Zelevinsky, *Discriminants, Resultants, and Multidimensional Determinants*, Birkhäuser, 1994.

[29] V. Guillemin, Kaehler structures on toric varieties, *J. Differential Geom.* **40** (1994) 285–309.

[30] K. Hori and C. Vafa, Mirror symmetry, arXiv:hep-th/0002222, 2000.

[31] H. Iritani, An integral structure in quantum cohomology and mirror symmetry for toric orbifolds, *Adv. Math.* **222** (2009) 1016–1079.

[32] J. Judd and K. Rietsch, The tropical critical point and mirror symmetry, arXiv:1911.04463, 2019.

[33] S. L. Kleiman, Toward a numerical theory of ampleness, *Ann. of Math.* **84** (1966) 293–344.

[34] T. V. Kadeishvili, On the homology theory of fibre spaces, *Russian Math. Surveys* **35** (1980) 231–238.

[35] D. Maclagan and B. Sturmfels, *Introduction to Tropical Geometry*, Graduate Studies in Mathematics **161**, AMS, 2015.

[36] R. T. Rockafellar, *Convex Analysis*, Princeton University Press, 1970.

[37] A. Schrijver, *Theory of Linear and Integer Programming*, Wiley, 1986.

[38] P. Seidel and R. Thomas, Braid group actions on derived categories of coherent sheaves, *Duke Math. J.* **108** (2001) 37–108.

[39] P. Seidel, *Fukaya Categories and Picard–Lefschetz Theory*, Zurich Lectures in Advanced Mathematics, EMS, 2008.

[40] J. Smith, Generating the Fukaya categories of compact toric varieties, arXiv:1804.06386, 2018.