Hom between the five objects ($c = 3 \cdot 2^{-4/3}$): only the diagonal survives.
First two: the leximin fibre; last three: the fibre (2/15, 1/6)

	P positive	*N* real $(-,-)$	*R* real $(-,+)$	C_ω complex	$C_{\bar\omega}$ complex
P positive	$\mathrm{Cl}(Q) \cong M_2$ $\det \sim \sqrt{2}\,\tau^{1/2}$ reduced: rank 1	0 same fibre, distinct m_0	0 disjoint fibres	0 disjoint fibres	0 disjoint fibres
N real $(-,-)$	0 same fibre, distinct m_0	$\mathrm{Cl}(Q) \cong M_2$ $\det \sim -\sqrt{2}\,\tau^{1/2}$ reduced: rank 1	0 disjoint fibres	0 disjoint fibres	0 disjoint fibres
R real $(-,+)$	0 disjoint fibres	0 disjoint fibres	$\mathrm{Cl}(Q) \cong M_2$ $\det \sim -c\,\tau^{2/3}$ reduced: rank 1	0 same fibre, distinct m_0	0 same fibre, distinct m_0
C_ω complex	0 disjoint fibres	0 disjoint fibres	0 same fibre, distinct m_0	$\mathrm{Cl}(Q) \cong M_2$ $\det \sim -c\,\bar\omega^2\tau^{2/3}$ reduced: rank 1	0 same fibre, distinct m_0
$C_{\bar\omega}$ complex	0 disjoint fibres	0 disjoint fibres	0 same fibre, distinct m_0	0 same fibre, distinct m_0	$\mathrm{Cl}(Q) \cong M_2$ $\det \sim -c\,\omega^2\tau^{2/3}$ reduced: rank 1