

Conditionals, Couplings and Toric Geometry: Established Results and a Fukaya Programme

Revised draft

Vinay K. Awasthi
Department of Applied Mathematics and Statistics
Johns Hopkins University*

October 4, 2026

Abstract

We separate the conditional/coupling formulation of the finite distribution monad into what is proved, what is imported from symplectic topology, and what is conjectural. The proved layer has four parts. (1) Markov kernels are the 1-cells of the Kleisli category; couplings are 2-cells whose vertical composition is the gluing lemma; the natural whiskerings are not functorial, and the strict 2-categorical structure is a graded metric shadow. (2) The base-relative posterior tower is a fibre over this base: its coherence, canonical section, strict mixed distributive law, admissibility data and obstructions are all transported along Markov kernels. (3) Coupling polytopes are moment polytopes: for nondegenerate marginals the polytope of couplings is Delzant, so it is the image of the moment map of a compact toric symplectic manifold, and when every cell is a facet the Fisher metric on the polytope is twice the Hessian of Guillemin's canonical symplectic potential. (4) The marginal space is cut by walls $\sum_I f = \sum_J g$; the combinatorial type, and hence the toric manifold, is locally constant off the walls and changes across them; the manifold is monotone exactly for uniform marginals, which are nondegenerate exactly for coprime sizes, and the smallest example is the degree-6 del Pezzo surface. The imported layer uses known Floer theory to locate a distinguished Lagrangian torus fibre in the monotone case over the *product* coupling, not the maximal coupling. The conjectural layer is restated with correct types: a family of toric manifolds over the chamber complex with wall-crossing in place of monodromy, and the whiskering defect as the second component of an A_∞ -functor rather than an m_3 . We also list the corrections to the first draft.

Contents

1	Introduction	2
2	The established layer: kernels and couplings	2
3	The fibre: stochastic change of base for the tower	3
4	Coupling polytopes are moment polytopes	4
4.1	The polytope and its lattice	4
4.2	Nondegeneracy implies Delzant	4
4.3	Walls, chambers and monotone cases	5

*Verification scripts: `check_glue.py`, `check_stochastic_basechange.py`, `check_conditionals.py`; see [section 9](#).

5	The imported layer: Floer theory of the special fibre	6
6	Corrections to the first draft	6
7	The conjectural layer, restated	7
8	Next computations	7
9	Computational checks	8

1 Introduction

Two structures sit on top of the finite distribution monad $\mathbf{D}$. The base-relative posterior tower [2] fixes an outcome base B and solves a *strict integrability* problem: it carries a mixed distributive law satisfying all four Beck axioms. The conditional/coupling formulation [1] lets the base vary along Markov kernels and compares kernels by couplings; it solves a *transport* problem. The relation is

$$\begin{aligned} \text{base-relative tower} &= \text{fibre over a fixed base,} \\ \text{conditional/coupling structure} &= \text{the varying-base global structure.} \end{aligned}$$

The first draft of this note left one gap: the tower’s change of base had been established only for functions $f: B \rightarrow B'$, while the global structure uses kernels $k: B \rightarrow \mathbf{D}(B')$. That gap is closed in [2, §9.7]; we state the result in [section 3](#).

The second purpose of this note is the geometry. The first draft conjectured a symplectic structure whose restriction to a fibre “is the Fisher metric”, and that the critical points of a projection are the maximal couplings. The correct statements turn out to be theorems about coupling polytopes ([section 4](#)), and they point in a different direction: the distinguished point is the product coupling.

Status of the results.

Layer	Content	Where
Proved	kernels, coupling 2-cells, graded shadow; stochastic base change of the tower; coupling polytopes are Delzant; Fisher = 2 Hess of Guillemin’s potential; walls and chambers; monotone cases	sections 2 to 4
Imported	Lagrangian torus fibres; nondisplaceability and Floer cohomology of the special fibre in the monotone case	section 5
Conjectural	families over the chamber complex; whiskering defects as A_∞ -functor data; a Fukaya-type functor	section 7

2 The established layer: kernels and couplings

$\mathbf{D}$ is the finitely supported distribution monad on **Set** and $\mathbf{Kl}(\mathbf{D})$ its Kleisli category. A 1-cell $k: B \rightsquigarrow B'$ is a Markov kernel $k: B \rightarrow \mathbf{D}(B')$, acting on states by $k^\#(p)(b') = \sum_b p(b)k(b)(b')$. The results of this section are proved in [1, §7]; we restate them for reference.

Proposition 2.1 (1-cells [1, Props. 7.2, 7.3]). $k^\#$ is affine, $(h \circ k)^\# = h^\# k^\#$, $k^\# \circ \mu = \mu \circ D(k^\#)$, and $\text{TV}(k^\# p, k^\# q) \leq \text{TV}(p, q)$. Attribution sets are transported along k , and precision can only be lost.

Definition 2.2 (Coupling 2-cells). For 1-cells $f, g: X \rightsquigarrow Y$, a coupling 2-cell $\alpha: f \Rightarrow g$ is a kernel $\alpha: X \rightarrow D(Y \times Y)$ with marginals $f(x)$ and $g(x)$. Its cost is $c(\alpha) = \max_x \Pr_{\alpha(x)}(y \neq y')$; the identity is the diagonal coupling; the vertical composite glues along the common marginal, $(\beta \cdot \alpha)(x)(y, y'') = \sum_{y'} \alpha(x)(y, y') \beta(x)(y', y'') / g(x)(y')$.

Theorem 2.3 (Witnesses and shadow [1, Thms. 7.6, 7.8]).

- (i) For fixed X, Y , 1-cells and coupling 2-cells form a category under gluing, strictly associative and unital. Cost is subadditive and $\min_\alpha c(\alpha) = d(f, g) = \max_x \text{TV}(f(x), g(x))$.
- (ii) Whiskering by a deterministic precomposition is strictly functorial. Whiskering by a stochastic precomposition (mixing couplings) or by a postcomposition (copying on equal pairs, independent on unequal pairs) preserves identities, marginals and the cost bound, but is not functorial.
- (iii) Graded 2-cells $f \Rightarrow_\varepsilon g$, $\varepsilon \geq d(f, g)$, with grades adding, form a strict 2-category on kernel 1-cells, and d makes $\text{Kl}(D)$ a Lawvere-enriched category.

3 The fibre: stochastic change of base for the tower

The base-relative tower over B has context algebra $A_B = DB \times D^2 B \times D^3 B \times \text{Sh}(B)$, reader comonad $\mathbb{C}^B(Y) = Y \times A_B$ and strict law λ^B [2, §9]. With $\text{Hyp} = \Theta \times D(-)$, a kernel $k: B \rightarrow D(B')$ acts on hypotheses by $\text{Hyp}(k)(\theta, \rho) = (\theta, k^\# \rho)$, and we put $A_k = k^\# \times D(k^\#) \times D^2(k^\#) \times D(\text{Hyp}k)$.

Theorem 3.1 (The tower is a functor on the Kleisli category [2, Prop. 9.14, Cor. 9.15, Prop. 9.16]).

- (i) A_k is a morphism of D -algebras, preserves coherent contexts, commutes with the canonical section σ , and $A_{h \circ k} = A_h A_k$; for deterministic k it is the earlier base change.
- (ii) $\text{id} \times A_k$ is a comonad morphism $\mathbb{C}^B \Rightarrow \mathbb{C}^{B'}$ commuting with the laws $\lambda^B, \lambda^{B'}$.
- (iii) Admissibility data, coherent lifts and the effective sets are transported; certificates pull back along $f' \mapsto k \triangleright f'$; failure levels can only rise and the secondary obstruction can only fall.
- (iv) Conditioning on an outcome observed after the channel equals conditioning on the pulled-back predicate $k \triangleright 1_e$, then changing base.

Graded 2-cells transport to every level: for kernels f, g , the level-one and level-two Kantorovich distances between $A_f(c)$ and $A_g(c)$ are at most $d(f, g)$.

The proof uses a single fact beyond naturality: Kleisli extension is a morphism of free D -algebras. All other components of A_k are pushforwards along functions. So the strict integrability of the fibre and the transport of the global structure are compatible, and the fibre is functorial on all of $\text{Kl}(D)$, not only on **Set**.

4 Coupling polytopes are moment polytopes

4.1 The polytope and its lattice

Let $f \in D(Y)$ and $g \in D(Y')$ have full support, with $|Y| = n$, $|Y'| = m$. (For coupling 2-cells $Y = Y'$; rectangular polytopes arise when gluing contexts [1, Prop. 7.11].) The *coupling polytope* is

$$\text{Cpl}(f, g) = \left\{ \alpha \in \mathbb{R}_{\geq 0}^{Y \times Y'} : \sum_{y'} \alpha(y, y') = f(y), \sum_y \alpha(y, y') = g(y') \right\}.$$

Its tangent space is the space of *circulations*, matrices with zero row and column sums. The integer circulations form a lattice Λ of rank $(n-1)(m-1)$; a basis is $E_{ij} - E_{i,m} - E_{n,j} + E_{n,m}$ for $i < n$, $j < m$.

Proposition 4.1 (Interior and Fisher metric). *Cpl(f, g) has dimension $(n-1)(m-1)$, its relative interior is the set of strictly positive couplings, and on it the Fisher metric $\mathcal{I}_\alpha(v, w) = \sum v(y, y')w(y, y')/\alpha(y, y')$ is positive definite on circulations.*

Proof. The product coupling $f \otimes g$ is strictly positive, so no inequality $\alpha \geq 0$ is an implicit equality, and the relative interior of the polytope in its affine hull is $\{\alpha > 0\}$. There $\mathcal{I}_\alpha(v, v) = \sum v^2/\alpha > 0$ for $v \neq 0$. $\square$

Remark 4.2. The first draft argued nondegeneracy through a radical and then distinguished “the Fisher metric” from “the Fisher–Rao metric”. These are the same metric, the Fisher information metric. The square-root embedding is a way of computing its geodesic distance [1, Prop. 8.1].

4.2 Nondegeneracy implies Delzant

Definition 4.3 (Walls). The pair (f, g) is *degenerate* if $\sum_{y \in I} f(y) = \sum_{y' \in J} g(y')$ for some nonempty proper subsets $I \subsetneq Y$ and $J \subsetneq Y'$. These equations define hyperplanes, the *walls*, in the space of pairs of marginals; the *chambers* are the connected components of the complement of the walls.

Theorem 4.4 (Coupling polytopes are Delzant). *If (f, g) is nondegenerate, then $\text{Cpl}(f, g)$ is a Delzant polytope with respect to Λ : it is simple, its facet normals are rational, and at each vertex the primitive edge vectors form a basis of Λ . Its edges at a vertex are the fundamental cycles of a spanning tree.*

Proof. Regard α as a weighting of the edges of the complete bipartite graph on $Y \sqcup Y'$. Vertices are forests. If the support of a feasible α contains a cycle, adding $\pm \epsilon$ alternately around it stays feasible, so α is not a vertex. If the support is a forest, the marginal equations determine α on it uniquely (peel off leaves), so α is a vertex. *Nondegeneracy makes them spanning trees.* Every node has positive marginal and hence an incident edge of the support. If the support forest had two or more components, take one with row set I and column set J . Then $\sum_I f = \sum_J g$; I and J are nonempty, and if $I = Y$ then $\sum_J g = 1$ forces $J = Y'$ and the component is everything. So I and J are proper, contradicting nondegeneracy. *Simplicity and smoothness.* At a vertex with spanning tree T , the zero cells are the $nm - (n + m - 1) = (n-1)(m-1)$ non-tree cells c . Each closes a unique cycle in $T \cup \{c\}$, giving a circulation C_c with entries in $\{0, \pm 1\}$ and $C_c(c) = 1$. A circulation w equals $\sum_{c \notin T} w(c)C_c$, because the difference is a circulation supported on the tree, hence zero. So the feasible directions at the vertex form the cone $\{\sum_c t_c C_c : t \geq 0\}$ on exactly $\dim \text{Cpl}(f, g)$ independent generators (simplicity), and the C_c are integral and form a $\mathbb{Z}$ -basis of Λ , since integer coefficients $w(c)$ suffice (smoothness). The facet normals are the coordinate functionals $\alpha \mapsto \alpha(c)$, which are rational. $\square$

By Delzant's theorem [5] there is a compact connected toric symplectic manifold $(X_{f,g}, \omega)$ of real dimension $2(n-1)(m-1)$, unique up to equivariant symplectomorphism, whose moment map $\mu: X_{f,g} \rightarrow \text{Cpl}(f,g)$ has image the coupling polytope. The fibre over an interior coupling is a Lagrangian torus $T^{(n-1)(m-1)}$, and the fibre over a vertex is a point.

Proposition 4.5 (Fisher metric and Guillemin's potential). *Suppose every inequality $\alpha(c) \geq 0$ defines a facet of $\text{Cpl}(f,g)$. The canonical Kähler metric of $X_{f,g}$ constructed by Guillemin [8] has, in action-angle coordinates over the interior, the form $\text{Hess } G$ in the polytope directions, with symplectic potential*

$$G(\alpha) = \frac{1}{2} \sum_c \alpha(c) \log \alpha(c).$$

Hence the Fisher metric on $\text{Cpl}(f,g)$ is twice the restriction of the canonical Kähler metric to the real section, and the product coupling $f \otimes g$, the coupling of maximal entropy, is the unique critical point of G on the polytope.

Proof. Guillemin's potential is $\frac{1}{2} \sum_i \ell_i \log \ell_i$ over the facets, where $\ell_i \geq 0$ are the facet inequalities written with primitive inward normals. Here $\ell_c(\alpha) = \alpha(c)$, and the coordinate functional is primitive on Λ because some integer circulation has entry 1 at c . Then $\text{Hess } G(v, v) = \frac{1}{2} \sum_c v(c)^2 / \alpha(c)$. The critical-point condition is that $\log \alpha$ is orthogonal to all circulations, that is $\log \alpha(y, y') = r(y) + s(y')$, which together with the marginals gives $\alpha = f \otimes g$. Uniqueness follows from strict convexity. $\square$

If some cell is not a facet, for instance when $f(y) + g(y') > 1$ forces $\alpha(y, y') > 0$, the potential omits that term and the identity holds only up to those terms.

4.3 Walls, chambers and monotone cases

Proposition 4.6 (Chambers). *The set of spanning trees that support vertices, and hence the combinatorial type of $\text{Cpl}(f,g)$ and the toric manifold $X_{f,g}$ up to equivariant diffeomorphism, is constant on each chamber. The type can change across a wall. For $f = (\frac{1}{2}, \frac{1}{2})$ and $g = (t, \frac{2}{3} - t, \frac{1}{3})$ the polytope is a hexagon for $t < \frac{1}{2}$ and a quadrilateral for $t > \frac{1}{2}$, both Delzant: two facets become redundant across the wall $g_1 = f_1$.*

Proof. The tree solution on a spanning tree T assigns to each tree edge the value $\sum_I f - \sum_J g$ (up to sign) for the bipartition (I, J) obtained by deleting that edge. So feasibility of T depends only on the signs of such differences, which are constant on chambers. The example is checked in `check_conditionals.py`. $\square$

Proposition 4.7 (Monotone cases). *Assume (f, g) is nondegenerate and every cell defines a facet. Then $X_{f,g}$ is monotone if and only if f and g are uniform, and uniform marginals are nondegenerate exactly when $\gcd(n, m) = 1$. In particular, in the square case $Y = Y'$ of coupling 2-cells no nondegenerate pair is monotone. On the wall, the uniform 2×2 pair still gives a Delzant segment, $X = \mathbb{CP}^1$, with the product coupling as its midpoint; the uniform $n \times n$ pair with $n \geq 3$ gives a polytope that is not simple.*

Proof. For a Delzant polytope written with primitive normals, the toric manifold is monotone exactly when some interior point has equal lattice distance to all facets: the classes of the toric divisors span H^2 subject to the linear relations given by the normals, c_1 is the sum of the divisor classes, and $[\omega]$ is the sum weighted by the facet distances. Here the distances are the entries $\alpha(c)$, so they are equal exactly when α is constant, which forces f and g to be uniform; then $\alpha = f \otimes g$. Uniform marginals are nondegenerate iff $a/n \neq b/m$ for $0 < a < n$, $0 < b < m$, iff $\gcd(n, m) = 1$, which fails when $n = m$. The two wall cases are checked in `check_conditionals.py` (T3). $\square$

Example 4.8 (The degree-6 del Pezzo surface). For f uniform on two points and g uniform on three, $\text{Cpl}(f, g)$ is a hexagon. In the basis of Λ above, its facet normals are $\pm e_1$, $\pm e_2$ and $\pm(e_1 + e_2)$. This is the fan of $\mathbb{CP}^2$ blown up at three points, the del Pezzo surface of degree 6. All six entries of the product coupling equal $\frac{1}{6}$, so it is the equidistant point. The case 3×4 (uniform) gives a Delzant polytope of dimension 6 with 96 vertices and all 12 cells facets, again monotone.

5 The imported layer: Floer theory of the special fibre

We now quote results from symplectic topology. They are not proved or checked here.

Nondisplaceability. For a monotone toric manifold, the torus fibre over the point at equal distance from all facets (the *special fibre*) is a stem in the sense of Entov and Polterovich, hence nondisplaceable [6]. For toric Fano manifolds, Cho and Oh show that a torus fibre with a flat $U(1)$ connection has nonvanishing Floer cohomology when it corresponds to a critical point of the disc potential [4]; see [3] for the Clifford torus and [7] for general compact toric manifolds.

Consequence for couplings. In the monotone cases of [proposition 4.7](#), the special fibre lies over the product coupling $f \otimes g$. For [example 4.8](#), the disc potential of the monotone special fibre is, up to the factor $T^{1/6}$, the Laurent polynomial

$$W(x, y) = x + y + \frac{1}{x} + \frac{1}{y} + xy + \frac{1}{xy},$$

read off from the facet normals, and it has a critical point at $(x, y) = (1, 1)$ (checked exactly), so the special fibre with trivial holonomy has nonzero Floer cohomology by [4].

The maximal coupling, by contrast, has zero entries whenever $f \neq g$, so it lies on the boundary of $\text{Cpl}(f, g)$, where the torus fibre collapses (`check_conditionals.py`, T6). The first draft’s conjecture that the critical points are the maximal couplings should therefore be replaced. The geometrically distinguished coupling is the product coupling, which is also the maximum-entropy coupling and the critical point of Guillemin’s potential ([proposition 4.5](#)).

6 Corrections to the first draft

1. *Candidate disc potential.* $W_\beta(f) = \sum_g \int_{\text{Cpl}(f, g)} e^{-\beta c(\alpha)} d\nu$ sums over a continuum of 1-cells g and is not defined as written. Its Laplace limit $\min_{g, \alpha} c(\alpha)$ is always 0, attained at $g = f$, $\alpha = \Delta_f$, so it carries no information. The toric disc potential of $X_{f, g}$ is the correct object ([section 5](#)).
2. *Symplectic structure.* A symplectic form is antisymmetric and the Fisher metric is symmetric, so the restriction of the one cannot be the other. The correct statement is [proposition 4.5](#): the Fisher metric is twice the canonical Kähler metric on the real section of $X_{f, g}$.
3. *Critical points.* “Critical points of $\pi: E \rightarrow \text{Kl}(\mathbf{D})$ ” is undefined, since π is a functor, not a smooth map. The replacement is the special fibre over the product coupling in the monotone cases, and the walls of [definition 4.3](#) as the locus where the family degenerates.
4. *Monodromy and Dehn twists.* The family $(f, g) \mapsto X_{f, g}$ is constant on chambers and changes by toric birational modifications across walls ([proposition 4.6](#)); no vanishing cycle or Dehn twist has been identified. [Conjecture 7.1](#) restates the question.

5. *The m_3 conjecture.* Gluing is strictly associative ([theorem 2.3\(i\)](#)), so each hom-category, regarded as an A_∞ -category, has $m_k = 0$ for $k \geq 3$. The whiskering defect compares $u \cdot (\beta \cdot \alpha)$ with $(u \cdot \beta) \cdot (u \cdot \alpha)$. Its inputs are two 2-cells and a 1-cell, and it measures the failure of whiskering to be a functor. In A_∞ language that is the second component $\mathcal{F}_2$ of an A_∞ -functor, not a ternary composition ([conjecture 7.3](#)).
6. *Homotopy category.* The graded shadow has hom-sets $[d(f, g), \infty)$ under addition. These are not the degree-zero cohomology of cochain complexes, so the shadow is not a homotopy category of an A_∞ -category without further structure.
7. *Fisher versus Fisher–Rao.* These are the same metric ([proposition 4.1](#)).
8. *References.* The companion tower is now version 9; its posted earlier version has the title “... and Weak Distributive Laws”, a claim withdrawn in the revisions.

7 The conjectural layer, restated

Conjecture 7.1 (Family over the chamber complex). *The toric manifolds $X_{f,g}$ form a family over each chamber, and crossing a wall is a toric birational transformation (a blow-up, blow-down or flip) determined by the pair (I, J) defining the wall. On a wall, $X_{f,g}$ degenerates to a singular toric variety whose moment polytope is the non-simple $\text{Cpl}(f, g)$.*

Question 7.2 (Location of Floer-nontrivial fibres). Outside the monotone cases, the fibres with nonvanishing Floer cohomology (possibly after bulk deformation) are located by the potential of [7]. Is the product coupling always among them, and do they have a statistical meaning?

Conjecture 7.3 (Whiskering as an A_∞ -functor). *After a linearization of coupling 2-cells (for instance chains on the coupling polytopes), whiskering by a 1-cell u extends to an A_∞ -functor whose first component is the whiskering of [theorem 2.3\(ii\)](#) and whose second component is the whiskering defect. The A_∞ -functor equation in arity two then reads $\mathcal{F}_1(m_2(\beta, \alpha)) - m_2(\mathcal{F}_1\beta, \mathcal{F}_1\alpha) = m_1\mathcal{F}_2(\beta, \alpha) \pm \mathcal{F}_2(m_1\beta, \alpha) \pm \mathcal{F}_2(\beta, m_1\alpha)$. Finding a linearization in which this holds is the first concrete target.*

Conjecture 7.4 (Fukaya-type functor). *There is a functor from a category built from the Fukaya categories of the $X_{f,g}$ (with special fibres as distinguished objects) to a linearization of the coupling 2-cells, compatible with gluing on one side and composition of Lagrangian correspondences on the other.*

8 Next computations

- (1) *The 2×3 family.* Describe the toric surfaces in every chamber of $\{(f, g)\}$ for $|Y| = 2$, $|Y'| = 3$, and the transformations across each wall ([conjecture 7.1](#)).
- (2) *Square 2×2 .* Here Cpl is a segment and $X = \mathbb{CP}^1$. Compute the coupling 2-cells, the whiskerings and their defects exactly for all four deterministic 1-cells and a rational grid of stochastic ones, as the base case of [conjecture 7.3](#).
- (3) *Level two.* Extend [theorem 4.4](#) to couplings of second-order states, using the level-two gluing of [1, Prop. 7.16].

9 Computational checks

All checks pass.

- `check_glue.py` (30 checks): [proposition 2.1](#) and [theorem 2.3](#).
- `check_stochastic_basechange.py` (13 checks): [theorem 3.1](#); the hexagon of [example 4.8](#); the Delzant property for random nondegenerate 3×3 marginals; non-simplicity of the uniform 3×3 case; the Hessian identity and critical point of [proposition 4.5](#).
- `check_conditionals.py` (7 checks): [theorem 4.4](#) by a geometric test (vertex degrees from the face lattice and unimodularity of primitive edge vectors) on random 2×3 , 3×3 and 2×4 cases; [proposition 4.6](#); the degenerate 2×2 and 3×3 cases; [proposition 4.7](#) and [example 4.8](#), including the 3×4 case; the critical point of W ; the boundary position of maximal couplings; strict associativity of gluing.

References

- [1] V. K. Awasthi. Precise attribution of model interventions: attribution sets, mechanism reach, path defects and gluing. Technical report, Johns Hopkins University, October 2026. Draft 4.
- [2] V. K. Awasthi. The three-level posterior tower: hierarchical Bayesian inference and mixed distributive laws. Technical report, Johns Hopkins University, October 2026. Revised version 9. An earlier version is posted as DOI 10.13140/RG.2.2.29743.50088.
- [3] C.-H. Cho. Holomorphic discs, spin structures, and Floer cohomology of the Clifford torus. *International Mathematics Research Notices*, 2004(35):1803–1843, 2004.
- [4] C.-H. Cho and Y.-G. Oh. Floer cohomology and disc instantons of Lagrangian torus fibers in Fano toric manifolds. *Asian Journal of Mathematics*, 10(4):773–814, 2006.
- [5] T. Delzant. Hamiltoniens périodiques et images convexes de l’application moment. *Bulletin de la Société Mathématique de France*, 116(3):315–339, 1988.
- [6] M. Entov and L. Polterovich. Quasi-states and symplectic intersections. *Commentarii Mathematici Helvetici*, 81(1):75–99, 2006.
- [7] K. Fukaya, Y.-G. Oh, H. Ohta, and K. Ono. Lagrangian Floer theory on compact toric manifolds I. *Duke Mathematical Journal*, 151(1):23–174, 2010.
- [8] V. Guillemin. Kaehler structures on toric varieties. *Journal of Differential Geometry*, 40(2):285–309, 1994.