
(a) the race: $\Lambda(1, A) = ts$, $\Lambda(2, A) = (1 - t)s^2$
Newton polyhedron N
ts
(source 1)
$-ts^2$
normal (1, 1)
s^2 (source 2)
exponent of t
exponent of s


(b) the order fan: two cones and a ray;
iterated limits sit at the corners
corner $w_t \gg w_s$:
$t \to 0$ first, limit 0
source 2 explains A
$\pi_1^A \to 0$
ray $t = cs$:
$\pi_1^A \to c/(1 + c)$
source 1 explains A
$\pi_1^A \to 1$
corner $w_s \gg w_t$:
$s \to 0$ first, limit 1
weight w_t
weight w_s (how fast $s \to 0$)


(c) the race along four arcs: the limit set
is all of [0, 1]
$t = s^{1/2}$: $\to 1$
$t = 3s$: $\to 3/4$
$t = s$: $\to 1/2$
$t = s^{3/2}$: $\to 0$
π_1^A
$s \to 0$ along each arc


(d) lines are not enough: every line agrees,
but curved arcs do not
parabola $y = 1x^2$: $\to 1/2$
parabola $y = 3x^2$: $\to 1/4$
lines $y = vx$: $\to 0$
share of the cell x^2 against y
$x \to 0$