
(a) the statistical fan, $s = N^{-1/2}$: Theorem J is one point on the exposure line
invisible: exposure → 0, impact → 0
Theorem J: the band, impact → B SE
exposure line $w_D + w_\pi/2 = 1$
exposed, impact → ∞
exposed, impact → 0
w_D (its shift $D \sim s^{w_D}$)
w_π (adversary's share $\pi \sim s^{w_\pi}$)


(b) Monte Carlo along three arcs ($\pi = 0.3$, $D = 1.5\,s^{w_D}$)
$w_D = \frac{1}{2}$: exposed
$w_D = 1$: the band
$w_D = \frac{3}{2}$: invisible
false-alarm rate
exposure rate $P(Z > c_s)$
records N


(c) Gaussian bands are valid only where event counts grow: $P(A) = 3N^{-w_A/2}$
$w_A = 1$: counts grow
$w_A = 2$: Poisson ray
$w_A = 3$: empirical wall
fixed $P(A) = 0.2$
coverage of the nominal 95% interval
records N


(d) one Newton polyhedron, two decorations: limit sets [0, 1] and [½, 1]
decoration $x \mid y$: $\pi_1^A = \frac{1}{1+c}$
decoration $x + y \mid x$: $\pi_1^A = \frac{1+c}{2+c}$
limiting π_1^A
arc coefficient c (along $y = c\,x$)