(a) step 0-1: the eight cells; the hidden channel adds $s\chi$.
$s \le 0.027$ (rarest orange cell, 110) and $s \ge -0.013$ (rarest green cell, 111)




(b) steps 7-8: before the probe the set rule picks the pair (0.332 > 0.33075);
at the truth the triple is better (0.333); Savage's regret already picks the triple




(c) steps 1, 2, 10, 11: one line; the decision boundary (0.668) is nearer the truth
than the detection boundary (0.66925), so a stealthy takeover exists




(d) step 6: the Stanley-Reisner complex of the fibre over an independent law




two tetrahedra on d1d2d3
(an actor absent)

three pages on the axis w1w2
(both present, differing in one variable)

glue equal labels: the complex has
facets {w1,d1,d2,d3}, {w2,d1,d2,d3},
{w1,w2,d1}, {w1,w2,d2}, {w1,w2,d3};
minimal non-faces {w1,w2,di,dj};
Groebner basis w1(1 - w1) di dj (i<j);
ideal = (w1) ∩ (w2) ∩ (d1,d2) ∩ (d1,d3) ∩ (d2,d3)

the truth: w1w2 = 0.24,
d = (0.45, -0.40, 0.30), all non-zero:
it lies on no facet, so the fibre is
flat there (a segment over the pairs)