

Q1 to Q10 are MCQs with only one correct answer. Choose the correct option.

- 1. Using a goodness of fit, we can assess whether a set of obtained frequencies differ from a set of frequencies.
- a) Mean
- b) Actual
- c) Predicted
- d) Expected

Answer: d) Expected

- 2. Chisquare is used to analyze
- a) Score
- b) Rank
- c) Frequencies
- d) All of these

Answer: c) Frequencies

- 3. What is the mean of a Chi-Square distribution with 6 degrees of freedom?
- a) 4
- b) 12
- c) 6
- d) 8

Answer: b) 12

- 4. Which of these distributions is used for goodness of fit testing?
- a) Normal distribution
- b) Chisqared distribution
- c) Gamma distribution



d) Poisson distribution

Answer: b) Chisqared distribution

5. Which of the following distributions is Continuous

- a) Binomial Distribution
- b) Hypergeometric Distribution
- c) F Distribution
- d) Poisson Distribution

Answer: c) F Distribution

- 6. A statement made about a population for testing purpose is called?
- a) Statistic
- b) Hypothesis
- c) Level of Significance
- d) TestStatistic

Answer: b) Hypothesis

- 7. If the assumed hypothesis is tested for rejection considering it to be true is called?
- a) Null Hypothesis
- b) Statistical Hypothesis
- c) Simple Hypothesis
- d) Composite Hypothesis

Answer: a) Null Hypothesis

- 8. If the Critical region is evenly distributed then the test is referred as?
- a) Two-tailed
- b) One tailed
- c) Three tailed



d) Zero tailed

Answer: a) Two-tailed

- 9. Alternative Hypothesis is also called as?
- a) Composite hypothesis
- b) Research Hypothesis
- c) Simple Hypothesis
- d) Null Hypothesis

Answer: b) Research Hypothesis

- 10. In a Binomial Distribution, if 'n' is the number of trials and 'p' is the probability of success, then the mean value is given by
- a) np
- b) n

Answer: a) np

