Homework 2: Basis for a Topology

Assignments should be **stapled** and written clearly and legibly.

- 1. §1.1, #1.4.
- 2. §1.2, #1.11(a),(b).
- 3. Let \mathcal{B} and \mathcal{B}' be two bases on a set X, and let $\mathcal{T}_{\mathcal{B}}$ and $\mathcal{T}_{\mathcal{B}'}$ be the corresponding topologies generated by the bases. Consider the following two statements:
 - (i) For every $B' \in \mathcal{B}'$ and $x \in B'$, there exists $B \in \mathcal{B}$ such that $x \in B \subseteq B'$.
 - (ii) For every $B \in \mathcal{B}$ and $x \in B$, there exists $B' \in \mathcal{B}'$ such that $x \in B' \subseteq B$.
 - (a) Prove that $\mathcal{T}_{\mathcal{B}}$ is finer than $\mathcal{T}_{\mathcal{B}'}$ if and only if (i) holds.
 - (b) Prove that $\mathcal{T}_{\mathcal{B}} = \mathcal{T}_{\mathcal{B}'}$ if and only if (i) and (ii) both hold. (Hint: use part (a).)
 - (c) Use the Union Lemma to express (i) in terms of unions.
- 4. §1.2 #1.12, 1.13.