

Homework 10: Kernel, Range

Assignments should be **stapled** and written clearly and legibly. Problems 3 and 4 are optional.

1. §8.1, #10, 11, 24.
2. Let V and W be vector spaces, and let $T : V \rightarrow W$ be linear. Prove that if $\{\mathbf{v}_1, \dots, \mathbf{v}_p\}$ spans V , then $\{T(\mathbf{v}_1), \dots, T(\mathbf{v}_p)\}$ spans $R(T)$.
3. Let $T : V \rightarrow W$ be a linear transformation. Let w be an element of W , and let v_0 be an element of V which T maps to w . Prove that the set of all vectors which T maps to w is equal to $\{v_0 + u : u \in \ker T\}$. (Note that this is not, in general, a subspace of V – why not?)
4. (Differential Equations) Let $V = W = C^\infty(-\infty, \infty)$ be the vector space of infinitely differentiable functions. Let $T : V \rightarrow W$ be the linear transformation

$$T(f) = a_m f^{(m)} + a_{m-1} f^{(m-1)} + \dots + a_1 f,$$

where $f^{(i)}$ denotes the i -th derivative of f , and a_m, a_{m-1}, \dots, a_1 are constants.

- (a) Recall that $\ker T = \{f \in V : T(f) = 0\}$. In terms of differential equations, what does $\ker T$ represent?
- (b) Let $g \in W$, and let $H = \{h \in V : T(h) = g\}$. In terms of differential equations, what does H represent?
- (c) Let h_0 be any element of H . Use Problem 3 to prove that $H = \{h_0 + f : f \in \ker T\}$.