

Homework 15: The Mean Value Theorem (Due May 12, 2023)

Directions. Assignments should be **stapled** and written clearly and legibly. Problem 6 is optional.

1. §6.2, #16.
2. Find a twice differentiable function $f(x)$ such that $f'(1) = -1$, $f'(4) = 7$, and $f''(x) > 3$ for all x , or prove that such a function cannot exist.
3. Suppose that $f : [a, b] \rightarrow \mathbb{R}$ has continuous derivatives f' and f'' , and assume there exists $c \in (a, b)$ such that $f(a) = f(c) = f(b)$. Prove that there exists d in (a, b) such that $f''(d) = 0$.
Hint. First prove that there exist x_1 and x_2 such that $a < x_1 < x_2 < b$ and $f'(x_1) = f'(x_2) = 0$.
4. Recall that a number c is called a **fixed point** of a function f if $f(c) = c$. Prove that if f is a differentiable function and $f'(x) \neq 1$ for all real numbers x , then f has at most one fixed point.
5. Suppose that f and g are differentiable functions such that $f' = g$ and $g' = -f$. Prove that $h(x) = (f(x))^2 + (g(x))^2$ is a constant function. (Hint: Use the chain rule.)
6. Let h be **Thomae's function**:

$$h(x) = \begin{cases} 1 & \text{if } x = 0 \\ 1/n & \text{if } x = m/n \text{ and } x = m/n \text{ in lowest terms with } n > 0 \\ 0 & \text{if } x \notin \mathbb{Q} \end{cases}$$

Prove that h is continuous at every $x \notin \mathbb{Q}$ and discontinuous at every $x \in \mathbb{Q}$.