

Applications of Chatterjee's Correlation in MCMC

(UGP, 21-22 Even Semester)

Vivek Kumar Singh, Dootika Vats

Department of Mathematics and Statistics
Indian Institute of Technology, Kanpur

April 17, 2022



Table of Contents

- 1 Markov chain Monte Carlo
- 2 Pearson's Correlation Coefficient
- 3 Chatterjee's Correlation Coefficient
- 4 Chatterjee's Autocorrelation Coefficient
- 5 Sketch of proof of consistency
- 6 Simulations



Markov Chain Monte Carlo

- A Markov chain is a discrete time stochastic process X_1, X_2, \dots such that the next state of the process depends solely on the present state, i.e.

$$\Pr(X_n | X_1, \dots, X_{n-1}) = \Pr(X_n | X_{n-1}).$$

- A Markov chain can be specified by two things
 - 1 The initial distribution, i.e. the marginal of X_1 .
 - 2 The transition probabilities, i.e. the conditional of $X_{n+1} | X_n$



Markov chain Monte Carlo

- A Markov chain is said to be **stationary** if the marginal of X_n is independent of n . This invariant distribution is called the stationary distribution.
- A Markov chain is **ergodic** if the distribution of X_n converges to the invariant distribution.
- A stationary Markov chain is **time-reversible** w.r.t. the stationary distribution if X_n and X_{n+1} are exchangeable.



Pearson's Correlation Coefficient



References I

