

Direct photo-production of narrow ρK resonance in CLAS

Moskov Amaryan

In collaboration with
Chandra Nepali and Gagik Gavalian

CLAS Collaboration Meeting
February 24, 2012

Outline

Introduction

Review of current status


CLAS Data analysis

Direct production of a resonance in pK system

Conclusions

Constituent Quark Model

Light Mesons



$$\underline{J^{PC} = 0^-}$$

(pseudoscalar
nonet)

$$\pi^0 = (u\bar{u} - d\bar{d})/\sqrt{2}$$

$$\eta^0 = (u\bar{u} + d\bar{d} - 2s\bar{s})/\sqrt{6}$$

$$\eta' = (u\bar{u} + d\bar{d} + s\bar{s})/\sqrt{3}$$

All are $q\bar{q}$ states

Why?


$$Q = I_3 + (B+S)/2$$

Gell-Mann-Nishijima

$$Y = B + S$$

B-baryon number

S-strangeness



$$\underline{J^{PC} = 1^-}$$


(vector nonet)

$$\rho^0 = (u\bar{u} - d\bar{d})/\sqrt{2}$$

$$\omega^0 = (u\bar{u} + d\bar{d})/\sqrt{2}$$

$$\phi = s\bar{s}$$

Light Baryons



octet


$$J^P = \frac{1}{2}^+$$

All are qqq states
Why?

We then refer to the members $u^{\frac{2}{3}}$, $\bar{d}^{-\frac{1}{3}}$, and $s^{-\frac{1}{3}}$ of the triplet as "quarks" q and the members of the anti-triplet as anti-quarks \bar{q} . Baryons can now be constructed from quarks by using the combinations (qqq) , $(qqq\bar{q})$, etc., while mesons are made out of $(q\bar{q})$, $(qqq\bar{q})$, etc. It is assuming that the lowest baryon configuration (qqq) gives just the representations 1, 8, and 10 that have been observed, while the lowest meson configuration $(q\bar{q})$ similarly gives just 1 and 8.

M.Gell-Mann, Phys. Lett.V8, Num.3, 214 (1964)


But where are multiquark states?



decuplet

$$J^P = \frac{3}{2}^+$$

Anti-decuplet



Already anticipated by Gell-Mann
in the same 1964 paper

For any value of z and of triplet spin, we can construct baryon octets from a basic neutral baryon singlet b by taking combinations $(bt\bar{t})$, $(btt\bar{t}\bar{t})$, etc. **. From $(bt\bar{t})$, we get the representations 1 and 8 , while from $(btt\bar{t}\bar{t})$ we get 1 , 8 , 10 , 10 , and 27 . In a similar way, meson singlets and octets can be made out of $(t\bar{t})$, $(t\bar{t}t\bar{t})$, etc. The quantum num-

Revitalized in the paper

D.Diakonov, V.Petrov, and M.Polyakov
Z.Phys.A359, 305(1997)

Based on χ QSM

Predicted narrow ($\Gamma < 15$ MeV), Θ^+ state at $M \sim 1530$ MeV

Where Θ^+ was searched for?

Since 2003 almost all particle and nuclear physics collaborations have been involved,
About few thousand physicists !
~1000 (or more) papers are published!

Japan:
SPRING-8
BELLE

USA: **CLAS,** ?
BABAR,D0,
HyperCP,
STAR, PHENIX

Europe:
SAPHIR, COSY,
HERMES, ZEUS,
H1,HERA-B,
NOMAD,ALEPH


Russia:
DIANA,
SVD,
Dubna,
SPHINX

China:
BES

A lot of negative results: Pentaquark is really elusive!

CLAS at Tampa 2005

nK^+ Mass Spectrum



- ▶ the nK^+ mass spectrum is smooth
- ▶ no structure is observed at a mass of ~ 1540 MeV

Since then the fate of Θ^+ was decided, but was it justified?
This is the question.

Where we stand?

- **The CLAS has set up an upper limit for the cross section**
- Many experiments do not see a signal, but should they see it?
- Some previous positive results still hold
- Is the case closed?
- Can we increase sensitivity to the tiny cross section ?
- What must be done in order to convince ourselves in existence or in absence of the resonance ?
- *We reported the signal via interference with ϕ (arXiv:110.3325)*
- **Can we observe it in the direct production?**

Reanalysis of CLAS Data


Reaction of interest:

$$\gamma + p = p K_S (\pi^+ \pi^-) X(K_L)$$



Experimental Strategy:

- a) In final state detect $p \pi^+ \pi^-$
- b) reconstruct K_S from $\pi^+ \pi^-$
- c) reconstruct K_L in missing mass of $p \pi^+ \pi^-$
- d) search for a resonance in $p K$ system
- e) invoke Dalitz plot distribution

Reconstructing K_S



Reconstruction of Ks and K_L



CLAS published paper
Phys. Rev. D74, 032001(2006)

$M_x(p) > 1.04 \text{ GeV}$

$\text{Doca1}(Kp) < 1 \text{ cm}$

$\text{Doca2} < 1 \text{ cm}$

$d > 3 \text{ cm}$

CLAS new analysis

$M_x(p) > 1.04 \text{ GeV}$



$\text{Doca1} < 1.0 \text{ cm}$

$\text{Doca2} < 0.5 \text{ cm}$


$d > 3 \text{ cm}$

$\cos\theta > 0.98$

Present Analysis:
 close # of signal events.
 But !




Missing Mass of Proton



$M=1.021$ GeV

$\sigma=7$ MeV

Dalitz Plot with no ϕ



No ϕ











$$M_X(p) > 1.035 \text{ GeV}$$

2σ cut


Missing Mass of Ks cutting out M(pKs)

Counts/20 MeV




$M_x(K_s) \text{ [GeV]}$

$M_x(K_s) \text{ [GeV]}$



Fit: Gauss1+Gauss2



Apply Vertex Cuts:

$\text{Doca1} < 1.0$ cm

$\text{Doca2} < 0.5$ cm

$\cos\theta > 0.98$

$d > 3.0$ cm

Select 2 kaons

$M(\pi^+\pi^-) = 0.497 \pm 0.004$ GeV

$M_x(pK_s) = 0.497 \pm 0.020$ GeV

Reject ϕ meson

$M_x(p) > 1.035$ GeV





Apply Dalitz Cut

$M(pK_s) < 1.52$ GeV

And



How stable is a peak vs ϕ -cut ?



very weak $M_x(p)$
cut dependence

peek one

No tuning!

$\text{Doca1} < 1.0 \text{ cm}$

$\text{Doca2} < 0.5 \text{ cm}$

$\cos\theta > 0.98$





$d > 3.0 \text{ cm}$

$M(pKs) < 1.52 \text{ GeV}$

$M_x(Ks) \text{ [GeV]}$


$M_x(Ks) \text{ [GeV]}$

How stable is a peak vs $\cos\Theta$ -cut?



$M_x(K_s)$ [GeV]

$M_x(K_s)$ [GeV]







$DOCA1 < 1.0\text{cm}$
 $DOCA2 < 0.5\text{cm}$
 $d > 3\text{cm}$

$M(pK_s) < 1.52\text{GeV}$


$M_x(p) > 1.035\text{ GeV}$

How stable is a peak vs DOCAI-cut?



$M_x(K_s)$ [GeV]

$M_x(K_s)$ [GeV]



DOCA2 < 0.5 cm





$d > 3$ cm

$\cos\Theta > 0.98$

$M_x(p) > 1.035$ GeV


$M(pK_s) < 1.52$ GeV

How stable is a peak vs DOCA2-cut?



$M_x(K_s)$ [GeV]

$M_x(K_s)$ [GeV]



DOCA1 < 1.0 cm





$d > 3$ cm

$\cos\Theta > 0.98$


$M_x(p) > 1.035$ GeV

$M(pK_s) < 1.52$ GeV

How stable is a peak vs Decay distance-cut?



$M_x(K_s)$ [GeV]



$M_x(K_s)$ [GeV]

$\text{DOCA1} < 1.0\text{cm}$

$\text{DOCA2} < 0.5\text{cm}$



$\cos\Theta > 0.98$

$M_x(p) > 1.035\text{GeV}$

$M(pK_s) < 1.52\text{GeV}$



How stable is a peak vs $\Delta M(\pi^+\pi^-)$ cut around Ks?

Counts/20 MeV



Don't panic!

Counts/20 MeV



No tuning!


all other cuts
the same

$M_x(K_s)$ [GeV]

$M_x(K_s)$ [GeV]

*“Le doute n'est pas une condition agréable,
mais la certitude est absurde.” Voltaire*

What if we change the binning?




And Finally:

10σ !

From our previous paper

arXiv: 1110.3325v1

PHYSICAL REVIEW C 85,
035209 (2012)





Interference with ϕ at low $t < 0.45 \text{ GeV}^2$

We hypothesized strong t dependence

Can we see it now?



How peak changes vs t-cut?



For illustration purposes
only:

**now we do not
need this cut**


However:




We see **sharper t-dependence**
for the peak at ~ 1.55 GeV
(it is less affected by the t-cut)

compared to (higher mass region) of
 Σ^* 's
with shallower t-dependence

The Effect of the CLAS Acceptance





No acc. correction:
sharp decrease near
threshold



Acceptance corrected
Peak at ~ 1.55 GeV
is very robust

How about invariant mass $M(pK_s)$?



Same cuts:



$DOCA1 < 1.0$ cm

$DOCA2 < 0.7$ cm

$\cos\theta > 0.95$


$d > 3$ cm

$M_x(p) > 1.035$ GeV



We see a signal in both channels


CLAS Acceptance for observed signals



It is very different for
 pK_S and pK_L

Therefore huge peak
at higher masses
(see previous slide)

Cross Section



in a 4MeV bin CLAS
upper limit **was 2nb**
(for $E_\gamma \sim 2\text{GeV}$)

now per 4MeV we have
(if calculated similarly)

$$\sigma_{\text{max}} = 1 \text{ nb}$$

$$\text{Br}[K_S(\pi^+\pi^-)] = 69\%$$

$$\text{Br}[K_S(K^0)] = 50\%$$

$$\text{Br}[\Theta^+(pK^0)] = 50\%$$

total cross section

over Θ peak:

$$\sigma = 20 \pm 7 \text{ nb}$$

Conclusions

- Narrow resonance in pK system is observed for the first time in a direct photo-production in CLAS
- Statistical significance of the peak is $\sim 10\sigma$
- Strangeness of the state is not fixed
- Observed peak could be either due to pentaquark Θ^+
- Or previously unobserved excited hyperon state
- **Observation** of narrow peak at $M(pK) \sim 1.545$ GeV


via interference **with ϕ**

and in a direct production **in both pK_s and pK_L channels**

Provides very strong confidence that the observed signal is real

- Preliminary estimation of the cross section is consistent with CLAS upper limit and integrated over the peak is
$$\sigma = 20 \pm 7 \text{ nb}$$

Backup: Classic Dalitz Plots




$\gamma p \rightarrow 1+2+3 \rightarrow K_S K_L p$

$$Q = T_1 + T_2 + T_3$$

T_i - KE in CM

$$Y = \frac{T_3}{Q} - \frac{1}{3}$$

$$X = \frac{T_2 - T_1}{\sqrt{3} Q}$$



it is insufficient
to cut out ϕ only