


1.5<Q²<2.5, 0.15<x_B<0.3, 0.2<−t<0.6; ε = 0.857; pseudo-data: 1180 rad events (even half)
model (generated Born)
corrected with δ_B (rad/Born MC)
corrected with δ_A, v_cut = 0.06
corrected with δ_A, v_cut = 0.2
dσ in the bin (pb per 30°)
corrected / model − 1 (%)
φ = φ_p′ + 180° (deg, Trento)
Structure functions: shift from the model, in % of σ_U
σ_U = σ_T + εσ_L: model 2.63 pb
σ_TT: model -1.17 pb = -0.447 σ_U
σ_LT: model -0.05 pb = -0.019 σ_U
(fit − model) / σ_U(model) (%)
δ_B (rad/Born MC)
δ_A, v_cut = 0.06
δ_A, v_cut = 0.2


1.0<Q²<2.5, 0.1<x_B<0.3, 0.09<−t<1.0; ε = 0.889; pseudo-data: 6013 rad events (even half)
model (generated Born)
corrected with δ_B (rad/Born MC)
corrected with δ_A, v_cut = 0.06
corrected with δ_A, v_cut = 0.2
dσ in the bin (pb per 30°)
corrected / model − 1 (%)
φ = φ_p′ + 180° (deg, Trento)
Structure functions: shift from the model, in % of σ_U
σ_U = σ_T + εσ_L: model 22.08 pb
σ_TT: model -10.22 pb = -0.463 σ_U
σ_LT: model -0.47 pb = -0.021 σ_U
(fit − model) / σ_U(model) (%)
δ_B (rad/Born MC)
δ_A, v_cut = 0.06
δ_A, v_cut = 0.2