Fermi motion in untagged J/ψ photoproduction on the deuteron

Bremsstrahlung beam from the 5 cm target · spectator model of the deuteron · yield ratio to a free neutron at rest

V. Kubarovsky, Jefferson Lab · 27 September 2026 · generator exclurad_py, branch jpsi-fermi, module generator/photo.py

0.994 ± 0.0003

yield per neutron, deuteron / free neutron, $E_\gamma > 8.6$ GeV: Fermi motion changes it by less than 1 %

2.55 %

of the J/ψ yield on the deuteron comes from photons below the free threshold 8.20 GeV

60 M events

three 20 M-event samples (deuteron, free n, free p), 35 minutes on ten cores, zero majorant violations

The generator: untagged photoproduction $\gamma N \to J/\psi N$

No scattered electron in the event. The beam radiates in the target, the photon strikes a nucleon, the final state is written to LUND.

1. **Photon energy** E_γ drawn from the bremsstrahlung spectrum, Tsai thin-target shape × Schiff endpoint factor, 6.5–10.6 GeV (tabulated inverse CDF)
2. **Target nucleon** at rest, or AV18 spectator momentum p_s: struck nucleon ($M_D - E_s$, $-p_s$) with off-shell mass m^*; draw rejected if $m^* < 0.10$ GeV
3. **Accept** $s = (k + p_N)^2$; accept with $\sigma_{tot}(s) \cdot (1 - \beta_z) / w_{max}$. σ_{tot} tabulated once; w_{max} is the strict head-on maximum, so no overweights
4. **t and ϕ** t from the exact inverse cumulative of $d\sigma/dt$ at this s (401 points, physical limits), ϕ uniform
5. **Kinematics** J/ψ and recoil built in the γN centre of mass, boosted with the actual photon and nucleon four-vectors: $k + p_D = J/\psi + N' + p_s$ to 10^{-9} GeV
6. **Output** LUND: beam pid 22 with E_γ in the header, target 2112 / 1000010020, $J/\psi \to e^+e^-$, recoil, spectator, vertex z along the cell

What is assumed on the hadron side

- the electroproduction model at $Q^2 = 0$: in the Hand convention $\sigma_T(Q^2=0)$ is the real-photon $d\sigma/dt$, so nothing new is assumed
- neutron model jpsin.roberts2026 = proton model relabelled, $\sigma_n = \sigma_p$
- deuteron machinery of the fermi branch reused unchanged (generator/fermi.py)

Validation


- four-momentum conservation 5.5×10^{-9} GeV (the LUND precision), spectator on shell to 3×10^{-16}
- free-proton E_γ spectrum vs $n_\gamma(E_\gamma)\,\sigma_{tot}(W)$: $\chi^2/ndf = 22.1/23$
- t′ slope at 9.5 GeV: 1.64 vs the model's 1.67 GeV^{-2}
- speed 0.09 ms/event free nucleon, 0.65 ms/event deuteron (one core)


Run


```
python3 -m exclurad_py.generator.photo card.txt --jobs 10
```

- key = value card: model, ebeam, kmin/kmax, t range, nev, seed, fermi, fermi_mstar_min, target length and X_0, vz, maj_safety, outroot
- config.json per run: photons per electron, ⟨σ⟩, yield per electron per nucleon

Photon beam: thin-target bremsstrahlung and the endpoint







Spectrum (Tsai, thin target)

- dN/dE_γ = (X/X₀)(1/E_γ)[4/3(1−γ)+γ²]·S(E_γ), γ = E_γ/E₀
- X = L/2: vertex density ∝ upstream thickness; LD_2 3.25 × 10^{-3}, LH_2 2.81 × 10^{-3}
- 1.44 × 10^{-3} photons/e in 6.5–10.6 GeV, 0.77 × 10^{-3} above threshold


Endpoint: complete screening is wrong there

- q_min = m_e²E_γ/2EE′ outgrows 1/a_s as E′ → 0: screening switches off and Schiff's bracket ln M + 1 − (2/b)arctan b falls through zero
- S(E_γ) = Schiff intermediate screening / complete-screening limit; screening radius = Bohr radius, 137 λ_C, for H and D

Size of the correction

- S = 0.97 / 0.91 / 0.77 / 0.58 / 0.36 at 300 / 100 / 30 / 10 / 3 MeV below E₀; zero at 0.42 MeV
- −1.24 % of the photons in 6.5–10.6 GeV; the last 100 MeV of the spectrum reshaped
- not included, each ≈ 1 MeV: dE/dx before radiating, beam spread, e–e endpoint

Fermi motion: plane-wave spectator model with the AV18 deuteron




E_γ [GeV]	configurations above threshold	$\langle p_s \rangle$ [MeV/c]	p_s > 300 MeV/c	removed by floor 0.30	floor 0.10
7.0	0.88 %	373	60.8 %	2.9 %	0.11 %
7.5	4.44 %	224	18.3 %	0.7 %	0.04 %
8.0	24.6 %	117	4.4 %	0.17 %	0.01 %
8.2	44.3 %	91	2.6 %	0.10 %	0.01 %
10.0	97.3 %	86	1.9 %	0.06 %	0.00 %


Kinematics of the struck neutron for a photon along +z, AV18 unweighted (200 000 configurations): near threshold the high-momentum tail is the signal, so the m* floor was lowered from the π^0/η value 0.30 GeV to 0.10 GeV.

Prescription

- deuteron at rest; spectator on shell, p_s from AV18 (S + D), isotropic
- struck nucleon $p_N = (M_D - E_s, -p_s)$, off shell: $m^{*2} = p_N^2 < M_n^2$
- $s = m^{*2} + 2E_\gamma(E_N - p_{N,z})$: a neutron running toward the photon raises s, one running away lowers it
- cross section evaluated at the free-nucleon mass and the event's invariants; m* enters only where the event is built
- the free threshold 8.20 GeV is no threshold: $\gamma d \to J/\psi\, n\, p$ opens at 5.66 GeV; generated range starts at 6.5
- floor $m^* \geq 0.10$ GeV: removes 0.01 % of the draws, bounds the boost to the struck-nucleon frame
- not included: FSI, coherent production, off-shell modification of the amplitude, D-wave tensor structure, $\sigma_n \neq \sigma_p$

Cross section model: roberts2026 against the CLAS12 data





Model

- σ_tot = A (W − W_th)^α · exp[−B (W − W_th)]
- A = 1.018 nb, α = 0.707, B = −0.928 GeV^{-1}, W_th = M_N + M_J/ψ = 4.036 GeV
- dσ/dt = σ_tot · b · exp(b t′), b = 1.67 GeV^{-2}, t′ = t − t_min
- d ln σ/dW = 2.7 GeV^{-1} at 10 GeV and diverges at threshold: shifts of s of 10^{-1} GeV² are visible in the yield
- |t|_min = 1.8 / 0.62 / 0.36 GeV² at 8.5 / 9.5 / 10.5 GeV


Against CLAS12 (Chatagnon et al. 2026, arXiv:2602.22128, Appendix A)


- total cross section described within errors from threshold to 9.6 GeV; at 9.7–10.5 GeV the data lie 20–30 % above the curve, 1–2 σ per point
- t dependence: single exponential compatible with the data at $|t| \lesssim 2$ GeV²; the points at 3–4 GeV² are a factor 2–3 above it
- the Fermi-motion ratio depends only on the slope of σ(W) near threshold, not on the normalisation; a refit of A, α, B, b is the next step for absolute yields

Why this shape

Near threshold σ ∝ (W − W_th)^0.7: convex and steep. That is what makes the deuteron yield sensitive to two small effects of opposite sign, binding and the flux-factor correlation (next slide).

Result: J/ψ yield per neutron, deuteron over free neutron at rest





Two effects of 4 %, opposite sign


- **weight = n_γ(E_γ) · n(p_s) · (1 − β_z) · σ(s)**
- binding: E_N = M_D − E_s = 0.933 < M_n, m* < M_n, so ⟨s⟩ at fixed E_γ is lower by 0.16 GeV² (W by 18 MeV): −4 to −5 %
- flux factor (1 − β_z): the Møller relative velocity of the moving neutron; averages to 1 but is correlated with s: +4 %
- net: between 0.99 and 1.00 at every E_γ above 8.6 GeV


E_γ [GeV]	ratio	without flux factor
8.2–8.4	1.776 ± 0.005	1.62
8.4–8.6	1.088 ± 0.002	1.015
8.6–9.0	0.999 ± 0.001	0.949
9.0–9.5	0.990 ± 0.001	0.952
9.5–10.0	0.993 ± 0.001	0.961
10.0–10.6	0.996 ± 0.001	0.968

What it means

- above 8.6 GeV a generator without Fermi motion is as good as one with it, to better than 1 % at every photon energy; mean 0.9944 ± 0.0003
- within 400 MeV of threshold the ratio is 1.09 and 1.78; below 8.20 GeV only the deuteron produces J/ψ: the threshold of σ(W) is smeared over ± 300 MeV and must be unfolded
- the first samples lacked the flux factor and gave 0.96: the cancellation is real and easy to miss

J/ψ below the free threshold





Numbers (20 M deuteron events)

- 2.55 % of the yield below 8.20 GeV, 1.41 % below 8.0, 0.33 % below 7.5
- 1.02×10^{-5} nb × photons/e per neutron below threshold
- 13 282 / 52 273 / 216 927 / 227 595 events in 6.5–7.0 / 7.0–7.5 / 7.5–8.0 / 8.0–8.2 GeV
- falls by an order of magnitude per 500 MeV: AV18 tail folded with the threshold rise of σ(W)


Reliability


- depends on exactly what is least certain: the AV18 tail beyond 300 MeV/c, the off-shell prescription, the absence of FSI
- read the sub-threshold yield as the plane-wave expectation, good to a factor of about two, not to percent


Consequences

- any threshold measurement on deuterium sees a smeared threshold; the generator provides the response to unfold it
- t below threshold: the events sit at the edge of phase space, |t|_min is large and the distribution is compressed toward it; the slope of a sub-threshold sample is not the elementary b
- the observable handle is the spectator proton (next slide)

The spectator proton tags the sub-threshold events







Above threshold

- ⟨p_s⟩ = 99 MeV/c, 4.4 % above 300 MeV/c
- ⟨m*⟩ = 0.915 GeV
- the bulk of AV18; the spectator is mostly undetectable


Below threshold

- all: ⟨p_s⟩ = 229 MeV/c, 26 % above 300 MeV/c, ⟨m*⟩ = 0.84 GeV
- below 8.0 GeV: 278 MeV/c, 36 %; below 7.5 GeV: 400 MeV/c, 68 %, ⟨m*⟩ = 0.71 GeV
- a fast proton at large angle: inside the CLAS12 acceptance

In the sample

- the spectator is written in every deuteron LUND event
- the floor m* ≥ 0.10 GeV removes 0.01 % of the draws, invisible in every distribution

Do we need radiative corrections for photoproduction?



Untagged 10.6 GeV electrons on a thin target: the quasi-real virtual photons of electroproduction with the electron undetected (equivalent-photon flux, Q² below 0.05–0.2 GeV²) outnumber the bremsstrahlung photons from the target by 8 to 11. The real CLAS12 sample is mostly quasi-real electroproduction.

1 · Real-photon process γN → J/ψ N: no

- no lepton in the initial state, hence no initial-state radiation; the hadronic vertex corrections are absorbed in the measured σ
- what remains are flux corrections: endpoint (done), dE/dx before radiating, beam spread, thick-target multi-photon — each ≲ 1 % and only at the tip

2 · J/ψ → e⁺e⁻(γ) final-state radiation: yes, in every mode

- leading-log estimate: 21 % of the decays have m_ee < 3.0 GeV, 16 % below 2.9 GeV (μ⁺μ⁻: 7.5 % and 5.6 %)
- moves events out of a tight mass window and distorts the peak; GEMC does not radiate, so the generator must (PHOTOS-like decay radiation, channel independent)
- this is the radiative correction that matters for photoproduction

3 · The actual CLAS12 configuration: electroproduction with RC

- 90 % of the untagged photons are virtual: use the electro path of exclurad_py (Q² → Q²_min, electron not required) with its radiative tail; the electron ISR/FSR changes the effective flux by a few %
- for the Fermi-motion ratio it is irrelevant: the flux cancels. photo.py remains the right tool for the bremsstrahlung part and for tagged real-photon beams

Summary and next steps

- **New photoproduction mode of exclurad_py:** bremsstrahlung beam with the correct endpoint, AV18 spectator deuteron, exact kinematics, strict majorant; 0.65 ms per event on the deuteron.
- **The flux factor (1 − β_z) of the moving nucleon is essential:** +4 %, and it cancels the −4 % binding effect. The same factor is missing in the electroproduction Fermi path (π^0: 0.2 % on the yield, 1–3 % tilt in W_obs and y, 9 % in the tail below the box edge).
- **Above 8.6 GeV Fermi motion changes the yield per neutron by less than 1 %** (mean 0.9944 ± 0.0003, every 100 MeV bin within 0.99–1.00).
- **Near and below threshold it is everything:** 1.09 and 1.78 in the two bins above 8.20 GeV, 2.55 % of the yield below it, spectator protons of 230–400 MeV/c that CLAS12 can see.
- **roberts2026 vs CLAS12:** agrees to 9.6 GeV, 20–30 % low above; the t slope holds to 2 GeV². Refit A, α, B, b to the CLAS12 tables.
- **Radiative corrections:** none for the real-photon process itself; $J/\psi \to e^+e^-\gamma$ decay radiation (15–20 % below 3.0 GeV) must be added; for absolute CLAS12 yields the quasi-real electro path with RC is the right generator.

Report: docs/jpsi_fermi/report/jpsi_fermi_photoproduction.pdf (branch jpsi-fermi) · OneDrive Work/2026_Jpsi_Fermi_photoproduction · samples /Volumes/wd_14tb/aao_norad_kin/jpsi_fermi/photo20M