



The MC in the bin after topology + DIS cuts, before exclusivity cuts. Dashed: the cut (3σ windows from the Born MC, outbending CD proton shown)
$1.5<Q^2<2.5$ GeV^2, $0.15<x_B<0.3$, $0.2<-t<0.6$ GeV^2; inbending + outbending






Reconstructed MC counts in the bin. 1.5<Q²<2.5 GeV², 0.15<x_B<0.3, 0.2<−t<0.6 GeV²; inbending + outbending
topology+DIS
+MM2(e'pgg), ME: full chain
born MC, N=9246
rad MC, N=8678
born MC, N=6006
rad MC, N=4317
reconstructed events per 30°
200
400
600
800
1000
1200
0
60
120
180
240
300
360
ϕγγ (deg, Trento)

Closure at generator level: same engine, same cut, no detector. A's ϕ is the proton's azimuth + 180°; with that definition B follows A.




The two corrections in one bin. 1.5<Q²<2.5 GeV², 0.15<x_B<0.3, 0.2<−t<0.6 GeV²; inbending + outbending
A, mx2_cut=0.08 (χ²=24/12)
A, mx2_cut=0.17 (χ²=12/12)
A, mx2_cut=0.3 (χ²=21/12)
A, mx2_cut=None (χ²=239/12)
B: rad/Born MC, full chain, φ_YY
B: same, φ_p + 180
δ_RC = N_rad/N_Born (B), σ_obs/σ_Born (A)
φ (deg)
0
60
120
180
240
300
360
0.6
0.8
1.0
1.2
1.4


A = B at
mx2_cut = 0.188
(0.151-0.231)
A: rc_factor, integrated over the bin
A, no cut: 1.155
B at generator level, same cut (closure)
B reconstructed, topology+DIS: 1.173
B reconstructed, +M(gg), MM2(e'p): 0.942
B reconstructed, +M(e'gg), cone, MPT: 0.925
B reconstructed, +MM2(e'pgg), ME: full chain: 0.899
δRC, bin integral
mx2_cut = upper limit on MM2(e'p) (GeV2)


χ2 of A against B (full chain), 12 ϕ bins
ndf = 12
mx2_cut (GeV2)