




cuts as in the note, Sec. 3.5: Q²>1, W>2, p_e'>2; |ME|<0.2, |MM²(e'pγγ)|<0.004, MP_T<0.75, coplanarity θ(X,π⁰)<4°; M_γγ 3σ, opening>1°
A: rc_factor, v_cut = 0.2
A: rc_factor, v_cut = 0.066
B: rad/Born MC, note cuts
δ_RC
B/A − 1 (%)
Q² (GeV²), other two integrated
x_B, other two integrated
−t (GeV²), other two integrated

Two variables at a time, A at v_cut = 0.200. cuts as in the note, Sec. 3.5: $Q^2>1$, $W>2$, $p_{e'}>2$; $|ME|<0.2$, $|MM^2(e'p\gamma\gamma)|<0.004$, $MP_T<0.75$, coplanarity $\theta(X,\pi^0)<4^\circ$; $M_{\gamma\gamma}$ 3σ, opening$>1^\circ$



Two variables at a time, A at v_cut = 0.066. cuts as in the note, Sec. 3.5: $Q^2>1$, $W>2$, $p_{e'}>2$; $|ME|<0.2$, $|MM^2(e'p\gamma\gamma)|<0.004$, $MP_T<0.75$, coplanarity $\theta(X,\pi^0)<4^\circ$; $M_{\gamma\gamma}$ 3σ, opening$>1^\circ$




The note's (Q^2, x_B, $-t$) cells with ≥ 200 events per arm
A, v_cut = 0.2
A, v_cut = 0.066
B: rad/Born MC, note cuts
δ_{RC}
B/A − 1 (%)
v_cut = 0.200: χ^2 = 128 / 35
v_cut = 0.066: χ^2 = 31 / 35
1.0-1.5 | 0.1-0.15 | 0.2-0.3
1.0-1.5 | 0.1-0.15 | 0.3-0.4
1.0-1.5 | 0.1-0.15 | 0.4-0.6
1.0-1.5 | 0.1-0.15 | 0.6-1.0
1.0-1.5 | 0.15-0.2 | 0.15-0.2
1.0-1.5 | 0.15-0.2 | 0.2-0.3
1.0-1.5 | 0.15-0.2 | 0.3-0.4
1.0-1.5 | 0.15-0.2 | 0.4-0.6
1.0-1.5 | 0.15-0.2 | 0.6-1.0
1.0-1.5 | 0.15-0.2 | 1.0-1.5
1.0-1.5 | 0.2-0.25 | 0.15-0.2
1.0-1.5 | 0.2-0.25 | 0.2-0.3
1.0-1.5 | 0.2-0.25 | 0.3-0.4
1.0-1.5 | 0.2-0.25 | 0.4-0.6
1.0-1.5 | 0.2-0.25 | 0.6-1.0
1.0-1.5 | 0.2-0.25 | 1.0-1.5
1.0-1.5 | 0.25-0.3 | 0.2-0.3
1.0-1.5 | 0.25-0.3 | 0.3-0.4
1.0-1.5 | 0.25-0.3 | 0.4-0.6
1.0-1.5 | 0.25-0.3 | 0.6-1.0
1.5-2.0 | 0.2-0.25 | 0.4-0.6
1.5-2.0 | 0.2-0.25 | 0.6-1.0
1.5-2.0 | 0.25-0.3 | 0.2-0.3
1.5-2.0 | 0.25-0.3 | 0.3-0.4
1.5-2.0 | 0.25-0.3 | 0.4-0.6
1.5-2.0 | 0.25-0.3 | 0.6-1.0
1.5-2.0 | 0.25-0.3 | 1.0-1.5
1.5-2.0 | 0.3-0.38 | 0.2-0.3
1.5-2.0 | 0.3-0.38 | 0.3-0.4
1.5-2.0 | 0.3-0.38 | 0.4-0.6
1.5-2.0 | 0.3-0.38 | 0.6-1.0
1.5-2.0 | 0.3-0.38 | 1.0-1.5
2.0-2.5 | 0.3-0.38 | 0.4-0.6
2.0-2.5 | 0.3-0.38 | 0.6-1.0
2.5-3.0 | 0.38-0.48 | 0.6-1.0

cuts as in the note, Sec. 3.5: $Q^2>1$, $W>2$, $p_{e'}>2$; $|ME|<0.2$, $|MM^2(e'p\gamma\gamma)|<0.004$, $MP_T<0.75$, coplanarity $\theta(X,\pi^0)<4^\circ$; $M_{\gamma\gamma}$ 3σ, opening$>1^\circ$

