

The fit against every set in it

run C1_with_compass, 11 sets, 1555 points, $\chi^2/\text{ndf} = 1.4018$

18 September 2026

Data red, model blue. Each panel carries its own kinematics and χ^2 ; the model curve includes the fitted normalisation where the set has one (De Masi 1.0586, Hall A neutron 0.9776, Hall A 2016 1.0583). Only sets that are *in* the fit are shown here — for the blind predictions (CLAS12 cross sections, the digitised De Masi α) see the full report.




Figure 1: All thirteen sets at a glance, fitted and blind.

CLAS6 n^0 BSA, ϕ distributions — $A_{LU}(\phi)$ — IN THE FIT
 $\chi^2 = 620.7 / 663 = 0.94$ per point




Figure 2: De Masi, $A_{LU}(\phi)$, all 60 bins — $A_{LU}(\phi)$. This set: $\chi^2 = 620.7/663 = 0.94$ per point.

CLAS6 π^0 structure functions — σ_{LT} — IN THE FIT
 $\chi^2 = 191.3 / 96 = 1.99$ per point




Figure 3: CLAS6 π^0 off the proton — σ_{LT} . This set: $\chi^2 = 482.4/288 = 1.68$ per point.

CLAS6 π^0 structure functions — σ_{TT} — IN THE FIT
 $\chi^2 = 120.2 / 96 = 1.25$ per point




Figure 4: CLAS6 π^0 off the proton — σ_{TT} . This set: $\chi^2 = 482.4/288 = 1.68$ per point.

CLAS6 π^0 structure functions — $\sigma_U = \sigma_T + \epsilon\sigma_L$ — IN THE FIT
 $\chi^2 = 170.9 / 96 = 1.78$ per point




Figure 5: CLAS6 π^0 off the proton — $\sigma_U = \sigma_T + \epsilon\sigma_L$. This set: $\chi^2 = 482.4/288 = 1.68$ per point.

CLAS6 η structure functions — σ_{LT} — IN THE FIT
 $\chi^2 = 46.3 / 69 = 0.67$ per point




Figure 6: CLAS6 η off the proton — σ_{LT} . This set: $\chi^2 = 217.0/207 = 1.05$ per point.

CLAS6 η structure functions — σ_{TT} — IN THE FIT
 $\chi^2 = 69.8 / 69 = 1.01$ per point




Figure 7: CLAS6 η off the proton — σ_{TT} . This set: $\chi^2 = 217.0/207 = 1.05$ per point.

CLAS6 η structure functions — $\sigma_U = \sigma_T + \epsilon\sigma_L$ — IN THE FIT
 $\chi^2 = 100.9 / 69 = 1.46$ per point




Figure 8: CLAS6 η off the proton — $\sigma_U = \sigma_T + \epsilon\sigma_L$. This set: $\chi^2 = 217.0/207 = 1.05$ per point.

Hall-A neutron, σ_T separated (2017) — σ_{LT} — IN THE FIT
 $\chi^2 = 6.3 / 4 = 1.58$ per point




Figure 9: Hall A 2017, π^0 off the NEUTRON — σ_{LT} . This set: $\chi^2 = 20.7/12 = 1.72$ per point.

Hall-A neutron, σ_T separated (2017) — σ_T — IN THE FIT
 $\chi^2 = 0.6 / 4 = 0.14$ per point




Figure 10: Hall A 2017, π^0 off the NEUTRON — σ_T . This set: $\chi^2 = 20.7/12 = 1.72$ per point.

Hall-A neutron, σ_T separated (2017) — σ_{TT} — IN THE FIT
 $\chi^2 = 13.8 / 4 = 3.44$ per point




Figure 11: Hall A 2017, π^0 off the NEUTRON — σ_{TT} . This set: $\chi^2 = 20.7/12 = 1.72$ per point.

Hall-A proton, σ_T separated (2016) — σ_{LT} — IN THE FIT
 $\chi^2 = 33.9 / 7 = 4.85$ per point




Figure 12: Hall A 2016, Rosenbluth separated — σ_{LT} . This set: $\chi^2 = 95.3/21 = 4.54$ per point.

Hall-A proton, σ_T separated (2016) — σ_T — IN THE FIT
 $\chi^2 = 34.1 / 7 = 4.87$ per point




Figure 13: Hall A 2016, Rosenbluth separated — σ_T . This set: $\chi^2 = 95.3/21 = 4.54$ per point.

Hall-A proton, σ_T separated (2016) — σ_{TT} — IN THE FIT
 $\chi^2 = 27.3 / 7 = 3.91$ per point




Figure 14: Hall A 2016, Rosenbluth separated — σ_{TT} . This set: $\chi^2 = 95.3/21 = 4.54$ per point.

Hall-A proton 6 GeV (2011) — σ_{LT} — IN THE FIT
 $\chi^2 = 45.9 / 28 = 1.64$ per point




Figure 15: Hall A 2011 — σ_{LT} . This set: $\chi^2 = 143.3/112 = 1.28$ per point.

Hall-A proton 6 GeV (2011) — $\sigma_{LT'}$ — IN THE FIT
 $\chi^2 = 16.0 / 28 = 0.57$ per point




Figure 16: Hall A 2011 — $\sigma_{LT'}$. This set: $\chi^2 = 143.3/112 = 1.28$ per point.

Hall-A proton 6 GeV (2011) — σ_{TT} — IN THE FIT
 $\chi^2 = 39.4 / 28 = 1.41$ per point




Figure 17: Hall A 2011 — σ_{TT} . This set: $\chi^2 = 143.3/112 = 1.28$ per point.

Hall-A proton 6 GeV (2011) — $\sigma_U = \sigma_T + \epsilon\sigma_L$ — IN THE FIT
 $\chi^2 = 42.0 / 28 = 1.50$ per point




Figure 18: Hall A 2011 — $\sigma_U = \sigma_T + \epsilon\sigma_L$. This set: $\chi^2 = 143.3/112 = 1.28$ per point.

Hall-A proton 12 GeV (2021) — σ_{LT} — IN THE FIT
 $\chi^2 = 91.9 / 36 = 2.55$ per point




Figure 19: Hall A 2021, the only data above $Q^2 = 4.3$ — σ_{LT} . This set: $\chi^2 = 358.2/144 = 2.49$ per point.

Hall-A proton 12 GeV (2021) — $\sigma_{LT'}$ — IN THE FIT
 $\chi^2 = 50.0 / 36 = 1.39$ per point




Figure 20: Hall A 2021, the only data above $Q^2 = 4.3$ — $\sigma_{LT'}$. This set: $\chi^2 = 358.2/144 = 2.49$ per point.

Hall-A proton 12 GeV (2021) — σ_{TT} — IN THE FIT
 $\chi^2 = 102.3 / 36 = 2.84$ per point




Figure 21: Hall A 2021, the only data above $Q^2 = 4.3$ — σ_{TT} . This set: $\chi^2 = 358.2/144 = 2.49$ per point.

Hall-A proton 12 GeV (2021) — $\sigma_U = \sigma_T + \epsilon \sigma_L$ — IN THE FIT
 $\chi^2 = 114.0 / 36 = 3.17$ per point




Figure 22: Hall A 2021, the only data above $Q^2 = 4.3$ — $\sigma_U = \sigma_T + \epsilon \sigma_L$. This set: $\chi^2 = 358.2/144 = 2.49$ per point.

COMPASS 2025, averaged over the grid — σ_{TT} — IN THE FIT
 $\chi^2 = 31.8 / 13 = 2.45$ per point




Figure 23: COMPASS 2025, model averaged over the grid — σ_{TT} . This set: $\chi^2 = 65.4/26 = 2.51$ per point.

COMPASS 2025, averaged over the grid — $\sigma_U = \sigma_T + \epsilon\sigma_L$ — IN THE FIT
 $\chi^2 = 33.5 / 13 = 2.58$ per point




Figure 24: COMPASS 2025, model averaged over the grid — $\sigma_U = \sigma_T + \epsilon\sigma_L$. This set: $\chi^2 = 65.4/26 = 2.51$ per point.

CLAS6 η beam-spin asymmetry — $A_{LU}^{\sin\phi}$ — IN THE FIT
 $\chi^2 = 13.4 / 12 = 1.12$ per point




Figure 25: Zhao, η beam-spin asymmetry — $A_{LU}^{\sin\phi}$. This set: $\chi^2 = 13.4/12 = 1.12$ per point.

CLAS12 $\sigma_{LT'}/\sigma_0$ — $\sigma_{LT'}/\sigma_0$ — IN THE FIT
 $\chi^2 = 45.6 / 30 = 1.52$ per point




Figure 26: CLAS12 $\sigma_{LT'}/\sigma_0$ — $\sigma_{LT'}/\sigma_0$. This set: $\chi^2 = 45.6/30 = 1.52$ per point.

eg1-dvcs, polarised target — A_{LL}^{const} — IN THE FIT
 $\chi^2 = 20.4 / 10 = 2.04$ per point




Figure 27: eg1-dvcs, polarised target — A_{LL}^{const} . This set: $\chi^2 = 71.7/40 = 1.79$ per point.

eg1-dvcs, polarised target — $A_{LL}^{\cos\phi}$ — IN THE FIT
 $\chi^2 = 21.0 / 10 = 2.10$ per point




Figure 28: eg1-dvcs, polarised target — $A_{LL}^{\cos\phi}$. This set: $\chi^2 = 71.7/40 = 1.79$ per point.

eg1-dvcs, polarised target — $A_{UL}^{\sin\phi}$ — IN THE FIT
 $\chi^2 = 21.0 / 10 = 2.10$ per point




Figure 29: eg1-dvcs, polarised target — $A_{UL}^{\sin\phi}$. This set: $\chi^2 = 71.7/40 = 1.79$ per point.

eg1-dvcs, polarised target — $A_{UL}^{\sin 2\phi}$ — IN THE FIT
 $\chi^2 = 9.3 / 10 = 0.93$ per point




Figure 30: eg1-dvcs, polarised target — $A_{UL}^{\sin 2\phi}$. This set: $\chi^2 = 71.7/40 = 1.79$ per point.