




TATA INSTITUTE OF FUNDAMENTAL RESEARCH

Graduate School Admissions (GS-2020)

Participant ID:	GS2020CHMIPH02256
Registration No:	GS2020CHMIPH003305
Participant Name:	SAJID SK
Test Center Name:	Techno India MAC Center
Test Date:	08/12/2019
Test Time:	9:30 AM - 12:30 PM
Subject:	Chemistry

Section : **PART A**

- Q.1** Compound **2** and compound **1** were mixed in a 1:4 molar ratio. The mixture was subjected to an ammonium molybdate test for phosphate estimation. The result of the molybdate test yielded a phosphate concentration of 36.0 mM for the mixture. What are the concentrations of compound **1** and compound **2** in the mixture?



Ans

- ☒ 1. Compound **1**: 30 mM
Compound **2**: 6 mM
- ☒ 2. Compound **1**: 24 mM
Compound **2**: 6 mM
- ☒ 3. Compound **1**: 28.8 mM
Compound **2**: 7.2 mM
- ☒ 4. Compound **1**: 24 mM
Compound **2**: 12 mM

Question Type : **MCQ**
Question ID : **4639241345**
Status : **Answered**
Chosen Option : **2**





- Q.2** The coefficient of compressibility of water at 293 K is $4.9 \times 10^{-6} \text{ atm}^{-1}$ in the range 1 to 25 atm pressure. What will be the value of work involving the compression of 1 mol of liquid water from a pressure of 1 atm to 25 atm at 293 K in a reversible process.

Ans

- ☒ 1. $21.75 \times 10^{-5} \text{ atm dm}^3$
- ☒ 2. $0.2234 \times 10^{-5} \text{ atm dm}^3$
- ☒ 3. $225.223 \times 10^{-5} \text{ atm dm}^3$
- ☒ 4. $2.752 \times 10^{-5} \text{ atm dm}^3$


Question Type : **MCQ**
Question ID : **4639241331**
Status : **Answered**
Chosen Option : **3**

Q.3 The octanol-water partition coefficient ($\log P$) for the following amines are as shown below:


				
Log P	1.36	1.08	0.21	-0.32

Ans


☒ 1.



☒ 2.



☒ 3.



☒ 4.



Question Type : **MCQ**

Question ID : **4639241321**

Status : **Answered**

Chosen Option : **2**

Q.4 The half-life for the decomposition of a substance dissolved in Chloroform is 6 hours at 298K. How much of the substance will be left after a day if the initial weight of the dissolved substance is 160mg.

Ans

☒ 1. 40 mg

☒ 2. 160 mg

☒ 3. 10 mg

☒ 4. 27 mg

Question Type : **MCQ**





Question ID : **4639241333**

Status : **Answered**

Chosen Option : **3**

Q.5 A complex of Chromium (3+) in aqueous hydrochloric acid, was found to exist in two geometric isomeric forms. A white precipitate was formed on addition of equimolar amount of AgNO_3 solution to the complex. The structure of the complex is:

Ans



Question Type : **MCQ**

Question ID : **4639241344**

Status : **Answered**

Chosen Option : **2**

Q.6 In the case of Eigenstates of a particle in a box with infinite walls, which of the following statements is true about the wavefunction ($\psi(r)$), its first ($\psi'(r) = d\psi(r)/dr$) and second derivatives ($\psi''(r) = d^2\psi(r)/dr^2$)?

Ans

 1.

All three quantities are continuous everywhere

 2.

$\psi(r)$, $\psi'(r)$ are continuous but not $\psi''(r)$.

 3.

$\psi(r)$ is continuous everywhere, but discontinuous at the boundaries of the box.

 4.

$\psi(r)$ is continuous everywhere, $\psi'(r)$ and $\psi''(r)$ are continuous everywhere except at the boundaries.


Question Type : **MCQ**

Question ID : **4639241340**

Status : **Answered**


Chosen Option : **4**

Q.7 The major product formed in the following reaction is




Ans


✗ 1.




✗ 2.



✗ 3.



✓ 4.




Question Type : MCQ

Question ID : 4639241357


Status : Answered

Chosen Option : 4


Q.8 Methyl groups in the following compounds are



I



II



III

Ans 1.

diastereotopic (**I**), enantiotopic (**II**), diastereotopic (**III**)

2.

enantiotopic (**I**), enantiotopic (**II**), enantiotopic (**III**)

3.

diastereotopic (**I**), neither enantiotopic nor diastereotopic (**II**), diastereotopic (**III**)

4.

enantiotopic (**I**), neither enantiotopic nor diastereotopic (**II**), enantiotopic (**III**)

Question Type : **MCQ**

Question ID : **4639241346**

Status : **Answered**

Chosen Option : **4**

Q.9 An optically transparent thin crystal shows a quadratic response to visible light, i.e. it scatters some of the light in a waveform that is square of the field of light that falls on it. If light of 500 nm wavelength falls on it, what are the wavelengths of the scattered light?

Ans

1. 500 nm and 750 nm

2. 500 nm and 250 nm

3. 500 nm and 1000 nm

4. 500 nm only

Question Type : **MCQ**

Question ID : **4639241323**

Status : **Answered**

Chosen Option : **2**

Q.10 A scalar coupling ($^2J_{\text{IH-1H}}$) constant for a doublet in compound X is measured as 5.0 Hz on a 500 MHz NMR spectrometer. Which of the following statements is incorrect?

Ans  1.

The difference $\Delta\delta$ for the two components of the doublet depends on the field strength of the spectrometer.

 2.

For this doublet, the coupling constant measured in Hz depends on the field strength of the spectrometer.

 3.

For this doublet, the coupling constant is 5.0 Hz when the spectrum of X is recorded on a 250 MHz NMR spectrometer

 4.

The difference $\Delta\delta$ for the two components of this doublet is 0.05ppm when measured on a 100MHz spectrometer.

Question Type : **MCQ**


Question ID : **4639241353**


Status : **Answered**


Chosen Option : **1**


Q.11 An organic compound is estimated through Dumas method and was found to evolve 6 moles of CO_2 , 4 moles of H_2O and 1 mole of N_2 gas. The formula of the compound is:

Ans

 1. $\text{C}_{12}\text{H}_8\text{N}$

 2. $\text{C}_{12}\text{H}_8\text{N}_2$

 3. $\text{C}_6\text{H}_8\text{N}$

 4. $\text{C}_6\text{H}_8\text{N}_2$

Question Type : **MCQ**

Question ID : **4639241336**

Status : **Answered**

Chosen Option : **4**

Q.12 An ideal gas obeys the gas law $PV = nRT$. However, real gases deviate from this, and their behavior can be described in

$$PV/nRT = 1 + B_P P + \dots$$

where B_P is the first virial coefficient in the expansion. Do you expect the B_P to be temperature dependent? If so, what would you expect the sign of B_P to be at low temperatures? (Near but above the critical point of condensation)

Ans

☐ 1. B_P will be temperature independent

☒ 2. Negative

☐ 3. Positive

☐ 4. Can be either positive or negative

Question Type : MCQ

Question ID : 4639241325

Status : Answered

Chosen Option : 2

Q.13

For an ideal gas, $\left(\frac{\partial H}{\partial P}\right)_T$ is equal to

Ans

☐ 1. V

☒ 2. 0

☐ 3. $\left(\frac{\partial V}{\partial T}\right)_P$

☐ 4. $\left(\frac{\partial T}{\partial P}\right)_H$

Question Type : MCQ

Question ID : 4639241332

Status : Answered

Chosen Option : 2

Q.14 Beer's law for electromagnetic radiation absorption through a medium can be expressed as:

$$I = I_0 \exp(-\alpha L)$$

Where, α is the absorption coefficient, L is the path length through the medium, I_0 is the incident intensity and I is the transmitted intensity.

Consider an absorbing medium with two energy levels spaced by 600 cm^{-1} . On this medium, when monochromatic infrared radiation on resonance is made incident, which of the following statement(s) holds true?

Ans  1.

Light absorption will increase with increasing the temperature of the absorbing medium

 2.

Beer's law is valid only for ultraviolet-visible part of the spectrum and is invalid in the infrared spectral region

 3.

Light absorption will decrease with increasing temperature of the absorbing medium

 4.

Light absorption not change with the temperature of the absorbing medium

Question Type : **MCQ**





Question ID : **4639241339**

Status : **Answered**

Chosen Option : **1**

Q.15 Which reaction has the greatest increase in entropy?

Ans  1.



Question Type : **MCQ**

Question ID : **4639241352**

Status : **Answered**

Chosen Option : **1**

Q.16 A protein sequence in solution at temperature T folds from a denatured extended state to its native state. Under which of the following conditions on changes in enthalpy (ΔH) and entropy (ΔS) should such a transition be NOT favorable?

Ans

✓ 1. $0 > \Delta H > T\Delta S$

✗ 2. $\Delta H < 0 < T\Delta S$

✗ 3. $0 > T\Delta S > \Delta H$

✗ 4. $\Delta H = 0$ and $T\Delta S > 0$

Question Type : **MCQ**

Question ID : **4639241350**

Status : **Answered**

Chosen Option : **1**

Q.17 Iron, Cobalt and Nickel are three known elements that display ferromagnetism. Which of the following statements is true about ferromagnets:

Ans

✗ 1.

In the presence of a magnetic field, the unpaired spins of a ferromagnet all align with the external field. Then in the absence of the external magnetic field, these spins then revert back immediately to their original state.

✗ 2.

The origin of magnetism in a ferromagnet arises from randomly arranged paired spins in a lattice.

✗ 3.

A ferromagnetic material is weaker (in its attraction to an external magnetic field) than a paramagnetic material

✓ 4. **None of the above.**

Question Type : **MCQ**


Question ID : **4639241356**

Status : **Answered**


Chosen Option : **1**


Q.18 In a spherical polar coordinate system, a point A at (x, y, z) in the Cartesian coordinate system can be described by (r, θ , ϕ) where r, θ , and ϕ have their usual meaning. Expression for the volume of an infinitesimally small cube confined by dx, dy, and dz in terms of the spherical coordinate system is given by

Ans

 1. $drd\theta d\phi$

 2. $r\sin\theta drd\theta d\phi$

 3. $r^2\sin^2\theta drd\theta d\phi$

 4. $r^2\sin\theta drd\theta d\phi$


Question Type : **MCQ**

Question ID : **4639241326**

Status : **Answered**


Chosen Option : **4**

Q.19 The nitrogen mustard Chlorambucil is an anti-cancer drug. Predict the product obtained when Chlorambucil interacts with the DNA base pair AT shown below under physiological conditions.




Ans


✓ 1.




✗ 2.



✗ 3.



✗ 4.



Question Type : **MCQ**

Question ID : **4639241341**

Status : **Answered**

Chosen Option : **1**

Q.20 Predict the reagent for the following amine de-protection reaction.



Ans

✓ 1. $\text{H}_3\text{O}^+/\text{H}_2\text{O}$

✗ 2. $\text{NaOH}/\text{H}_2\text{O}$

✗ 3. $\text{NaCl}/\text{H}_2\text{O}$

✗ 4. $\text{Pd}/\text{C}, \text{H}_2$

Question Type : MCQ

Question ID : 4639241320

Status : Answered

Chosen Option : 1

Q.21 Which of the following belongs to the same symmetry group as NH_3 ?

Ans

✗ 1. BF_3

✗ 2. CH_4

✗ 3. CH_3OH

✓ 4. CHCl_3

Question Type : MCQ

Question ID : 4639241337

Status : Answered


Chosen Option : 4


Q.22 A flask contains 1 litre of growth medium (in which E. coli. bacteria are grown) with 180 gm of glucose as the only Carbon source for the bacteria. Provide a rough estimate of the number of E. coli. that can grow upto the final stage given


- a. A bacterium contains $\sim 3 \times 10^6$ protein molecules
- b. An average protein has ~ 300 amino acids
- c. Half of all glucose consumption goes to production

Ans

 1. 10^6

 2. 3×10^{14}

 3. 3×10^{16}

 4. 2.5×10^{13}

Question Type : MCQ


Question ID : 4639241322


Status : Not Answered

Chosen Option : --


Q.23 Rotational energy of a diatomic molecule is given by $E_{rot} = J(J + 1)hB_e$, where E_{rot} is in Joules. If the rotational constant for H_2 molecule is given as $B_e = 1.8324 \times 10^{12}$ Hz, the rotational period of the H_2 molecule in $J = 10$ level will be

Ans

 1. 1.33×10^{-19} sec

 2. 5.0×10^{-15} sec

 3. 5.46×10^{-13} sec

 4. 7.39×10^{-7} sec

Question Type : MCQ


Question ID : 4639241328

Status : Answered

Chosen Option : 2

Q.24 In a face centered arrangement of A and B atoms. Where A atoms are at the corners of the unit cell and B atoms are at the face centers. For each unit cell, one A atom is missing from a corner position and one B atom is missing from one face position. The simplest formula of the resulting compound will be:

Ans



Question Type : **MCQ**



Question ID : **4639241355**

Status : **Answered**

Chosen Option : **2**

Q.25 IR spectra of proteins have a peak in the $1600 - 1700 \text{ cm}^{-1}$ region, known as the "Amide I" region. This comes largely from the backbone $C=O$ stretch vibration. Given the natural abundance of different isotopes, would occasionally find a backbone $C=O$ at a different frequency? If so, what would be the next most common frequency, assuming that the "Amide I" has a peak exactly at 1650 cm^{-1} ?

Ans



No other frequency will be observed

Question Type : **MCQ**

Question ID : **4639241324**

Status : **Answered**

Chosen Option : **2**

Q.26 For a Harmonic Oscillator in its ground state, i.e., $v = 0$ state, the energy is given by $E = \frac{1}{2}h\nu$, where ν is the vibrational frequency. This is due to its

Ans

☒ 1. Kinetic energy

☒ 2. Potential energy

☒ 3.

Sum of Kinetic and Potential energies

☒ 4. Heat of formation

Question Type : **MCQ**

Question ID : **4639241329**

Status : **Answered**

Chosen Option : **3**

Q.27 Among the following, which molecule will have the smallest spacing between vibrational levels? (superscripts denote the nuclear mass of the atoms)

Ans

☒ 1. $^1\text{H}^{35}\text{Cl}$

☒ 2. $^1\text{H}^{37}\text{Cl}$

☒ 3. $^2\text{H}^{37}\text{Cl}$

☒ 4. $^2\text{H}^{35}\text{Cl}$

Question Type : **MCQ**

Question ID : **4639241343**

Status : **Answered**

Chosen Option : **3**

Q.28 The degeneracy of the energy level $12 \frac{h^2}{8ma^2}$ of a particle in a three dimensional cube of length "a" is

Ans

☒ 1. 1

☒ 2. 3

☒ 3. 6

☒ 4. 12

Question Type : **MCQ**

Question ID : **4639241327**

Status : **Answered**

Chosen Option : **1**

Q.29 A particle is thermally diffusing on a 1-dimensional harmonic potential given by $V(x) = \frac{1}{2}kx^2$, A where k is the spring constant and x is the position of the particle. The equilibrium probability $P(x)$ of finding the particle at a position x follows:

Ans

☐ 1. $P(x) \propto x^2$

☐ 2. $P(x) \propto x$

☒ 3.

$P(x) \propto \exp(-Ax^2)$, where A is a constant

☐ 4.

$P(x) \propto \exp(-Ax)$, where A is a constant

Question Type : **MCQ**

Question ID : **4639241351**

Status : **Answered**

Chosen Option : 1

Q.30 What is the last digit of 3^{4798} ?

Ans

☐ 1. 3

☒ 2. 9

☐ 3. 1

☐ 4. 7

Question Type : **MCQ**

Question ID : **4639241354**

Status : **Answered**

Chosen Option : 2

Q.31 How many normal modes does the CS_2 molecule have? How many of them can be observed using IR spectroscopy?

Ans

☐ 1. 3 modes, 2 can be observed

☐ 2. 6 modes, 5 can be observed

☒ 3. 4 modes, 3 can be observed

☐ 4. 3 modes, 3 can be observed

Question Type : **MCQ**

Question ID : **4639241348**

Status : **Answered**

Chosen Option : 3

Q.32 Within a circle of radius 'b', four largest possible identical circles of radius 'a' are fit such that they do not cross each other. What is the ratio a/b?

Ans

☐ 1. $\frac{1}{2}$

☒ 2. $\frac{1}{(1+\sqrt{2})}$

☐ 3. $\frac{2}{(1+\sqrt{2})}$

☐ 4. $\frac{1}{2}(1+\sqrt{2})$

Question Type : MCQ

Question ID : 4639241338

Status : Answered

Chosen Option : 2

Q.33 If 1.50 g of $\text{H}_2\text{C}_2\text{O}_4 \cdot 2\text{H}_2\text{O}$ were heated to drive off the water of hydration, how much anhydrous $\text{H}_2\text{C}_2\text{O}_4$ would remain?

Ans

☐ 1. 0.34 g

☐ 2. 0.92 g

☒ 3. 1.07 g

☐ 4. 1.50 g

Question Type : MCQ

Question ID : 4639241334

Status : Answered

Chosen Option : 3

Q.34 Order of field effect of alkyl groups are tert-butyl > iso-propyl > ethyl > methyl. Which molecule among PhCH_3 , PhCH_2CH_3 , $\text{PhCH}(\text{CH}_3)_2$ and $\text{PhC}(\text{CH}_3)_3$ has the highest dipole moment in gas phase.

Ans



Question Type : MCQ

Question ID : 4639241318

Status : Answered

Chosen Option : 4

Q.35 The Standard reaction Gibbs energy (ΔG°) for the ATP hydrolysis



In a typical bacterial cell, the concentration of ATP, ADP and P_i are 8 mmol L^{-1} , 1 mmol L^{-1} and 8 mmol L^{-1} , respectively. What is the reaction Gibbs energy under this condition?

Ans

☐ 1. -31 kJ mol^{-1}

☒ 2. -49 kJ mol^{-1}

☐ 3. -18 kJ mol^{-1}

☐ 4. -13 kJ mol^{-1}

Question Type : **MCQ**

Question ID : **4639241330**

Status : **Answered**

Chosen Option : **1**

Q.36 Light excitation of a molecule promotes an electron from a state $\phi_a(x, y, z)$ to $\phi_b(x, y, z)$, where x, y, z are spatial coordinates with respect to the molecular centre of mass. The change in the spatial position x of the electron can be calculated as:

Ans ☒ 1.

$$\int_{-\infty}^{\infty} \phi_b^*(x, y, z) x \phi_b(x, y, z) dx dy dz - \int_{-\infty}^{\infty} \phi_a^*(x, y, z) x \phi_a(x, y, z) dx dy dz$$

☐ 2.

$$\int_{-\infty}^{\infty} \phi_b^*(x, y, z) x \phi_b(x, y, z) dx - \int_{-\infty}^{\infty} \phi_a^*(x, y, z) x \phi_a(x, y, z) dx$$

☐ 3.

$$\int_{-\infty}^{\infty} \phi_b^*(x, y, z) \phi_b(x, y, z) dx - \int_{-\infty}^{\infty} \phi_a^*(x, y, z) x \phi_a(x, y, z) dx$$

☐ 4.

$$\int_{-\infty}^{\infty} \phi_b^*(x, y, z) \phi_b(x, y, z) dx dy dz - \int_{-\infty}^{\infty} \phi_a^*(x, y, z) \phi_a(x, y, z) dx$$


Question Type : **MCQ**

Question ID : **4639241349**


Status : **Not Answered**

Chosen Option : **--**


Q.37 An isolated water molecule has C_{2v} symmetry, however water clusters can have symmetries other than C_{2v} . What are the symmetries of the following three water clusters, which are in the plane of the paper?



(i)



(ii)



(iii)

Ans

☒ 1. All of them have C_{2v} symmetry

☒ 2. i) C_{2v} ii) D_{2h} iii) C_s

☒ 3. i) C_s ii) D_{2h} iii) C_s

☒ 4. i) C_{2v} ii) D_{2h} iii) C_{2v}


Question Type : MCQ

Question ID : 4639241342

Status : Answered

Chosen Option : 1

Q.38 How many absorption peaks will the following compound have in its ^{13}C -NMR spectrum?



Ans

☒ 1. 3

☒ 2. 4

☒ 3. 5

☒ 4. 6

Question Type : MCQ


Question ID : 4639241335

Status : Answered





Chosen Option : 4

Q.39

Predict the product of the following reaction.



Ans



Question Type : MCQ

Question ID : 4639241319


Status : Answered

Chosen Option : 2


Q.40 Which one of the following compounds has largest dipole moment?



I



II



III

Ans

✗ 1.

II

✓ 2.

III

✗ 3.

I

✗ 4.

All of them have equal dipole moment

Question Type : MCQ

Question ID : 4639241347

Status : Answered

Chosen Option : 2