

MEST - Assignment 2

Due date: 04/03/2022

(For each solution, show your work through a set of important steps. Use Dirac Notation for all questions)

Linear algebra and Dirac notation

1. Write the following expressions in the Dirac formalism (1+1+2 pts):

(a) $f(x) = \sum_n \phi_n(x) \int \phi_n^*(x') f(x') dx'$

(b) $\frac{\partial}{\partial x} f(x) = m(x) \int m^*(x') g(x') d(x')$

(c)

$$E_0^{MP2} = \sum_{ij}^{occ} \sum_{ab}^{virt} \frac{|\int dx_1 dx_2 \chi_i^*(x_1) \chi_j^*(x_2) r_{12}^{-1} (1 - \hat{P}_{12}) \chi_a(x_1) \chi_b(x_2)|^2}{\epsilon_i + \epsilon_j - \epsilon_a - \epsilon_b}$$

(\hat{P}_{12} is the permutation operator)

2. Prove that unitary transformation preserves the :

(a) Hermiticity of a matrix (2 pts)

(b) Trace of a matrix (2 pts)

3. In quantum computing a qubit is a two-state quantum mechanical system that corresponds to the basic unit of quantum information. Qubit states are generally represented by a superposition of two orthonormal basis states (say in basis X) denoted as $|\alpha\rangle = \begin{pmatrix} 1 \\ 0 \end{pmatrix}$, $|\beta\rangle = \begin{pmatrix} 0 \\ 1 \end{pmatrix}$. Suppose we have qubit states in basis Y represented by $|+\rangle = \frac{1}{\sqrt{2}}(|0\rangle + |1\rangle)$, $|-\rangle = \frac{1}{\sqrt{2}}(|0\rangle - |1\rangle)$. (1 + 2 + 1 + 2 pts)

(a) Write the qubit state $|\psi\rangle = a|-\rangle + b|+\rangle$ as linear combination of basis states of basis X ($|\alpha\rangle, |\beta\rangle$). What are the conditions on a and b for $|\psi\rangle$ to be normalized state vector

(b) Find the unitary transformation matrix H that map the basis X ($|\alpha\rangle, |\beta\rangle$) to basis Y ($|+\rangle, |-\rangle$)

(c) Write out the $H|\psi\rangle$ in terms of basis X and basis Y

(d) What is the probability of obtaining the $|\alpha\rangle$ and $|\beta\rangle$ states when measuring the $H|\psi\rangle$

Slater determinant, perturbation and Variation principle

1. Consider a system (for example, a molecule) consisting of three electrons with spins α, α, β occupying three different MOs ψ_1, ψ_2 and ψ_3 that are spatially different. (2 + 3 + 5 pts)
 - a) List the possible Slater determinants (SDs) for this case.
 - b) Are these SDs suitable approximate wave functions of such a system [Hint: check if they are pure spin states?]
 - c) Construct suitable approximate wavefunctions that are pure spin states [Hint: try to diagonalize the S^2 matrix in this SD basis].
2. Consider a charged particle in one-dimensional harmonic oscillator potential. Suppose we turn on a weak electric field (E) so that the potential is shifted by an amount $V' = -qEx$ (q is the charge of the particle). (3 + 3 pts)
 - (a) Show that there is no first-order change in the energy levels and calculate the second order correction.
 - (b) The Schrodinger equation can be solved exactly in this case by a change of variables. Find the exact energies and show that they are consistent with the perturbation theory approximation.
3. In relativistic theory the kinetic energy is given by the expression $T = \sqrt{p^2c^2 + m^2c^4} - mc^2$. This relativistic equation for kinetic energy reduces to classical result in nonrelativistic limit $p \ll mc$; expanding in powers of small number (p/mc) we have the lowest-order relativistic contribution to the Hamiltonian is $V' = -\frac{\hat{p}^4}{8m^3c^2}$ where p is momentum operator. This term is responsible for orbital contraction in elements (e.g Lanthanide contraction is partially attributed to relativistic effects) and shift the energy levels. Prove that the first order relativistic energy correction is given by:

$$E_r^{(1)} = -\frac{E_n^2}{2mc^2} \left[\frac{4n}{l + \frac{1}{2}} - 3 \right]$$

(Use: $\langle \frac{1}{\hat{r}} \rangle = \frac{1}{an^2}$ and $\langle \frac{1}{\hat{r}^2} \rangle = \frac{1}{a^2n^3(l+\frac{1}{2})}$) (5 pts)

4. Using a wavefunction based on hydrogenic orbitals the helium atom ground state energy differs by ≈ 0.15 a.u. from experimental values. This can be improved by treating the the nuclear charge, Z , as a variational parameter. Derive the general expression for energy expectation value in terms of Z for the ground-state wavefunction given by

$$\Psi(\bar{r}_1, \bar{r}_2) = \phi_{1s}(r_1)\phi_{1s}(r_2)\frac{1}{\sqrt{2}} [\alpha(\omega_1)\beta(\omega_2) - \alpha(\omega_2)\beta(\omega_1)] \quad (1)$$

To simplify the evaluation, use the spherical harmonic expansion of interelectronic repulsion term

$$\frac{1}{r_{12}} = \frac{1}{r_{>}} \sum_{l=0}^{\infty} \left(\frac{r_{<}}{r_{>}} \right)^l \frac{4\pi}{2l+1} \sum_{m=-l}^l Y_l^m(\theta_1, \phi_1) Y_l^m(\theta_2, \phi_2) \quad (2)$$

Here $r_> = \max(r_1, r_2)$ and $r_< = \min(r_1, r_2)$. Find the optimal Z value and the corresponding ground-state energy. (Hint: for the evaluation of this integral, split the integral into two parts corresponding to $(r_1 > r_2)$ and $(r_2 > r_1)$; and note that only $l=0$ contributes in the case of $1s$ orbitals). (10pts and you have taken the first steps in becoming a quantum chemist)

Total angular momentum, Term symbol and Spin-orbit coupling

1. Compute the following commutators given that $J(1)$ and $J(2)$ are individual angular momentum operators and J is total angular momentum operator (5 pts):

(a) $[J(1) \cdot J(2), J]$

(b) $[J(1) \cdot J(2), J^2]$

(c) $[J^2, J(1)^2]$

(d) $[J^2, J(1)_z]$

(e) $[J^2, J_z]$

2. The spin-orbit coupling term is given by the expression $H_{SO} = \xi L \cdot S$ where, ξ is spin-orbit constant and L and S are orbital and spin angular momentum. In class we have seen that emission line of atomic spectra of sodium split into closely spaced doublet having wavelength 5889.96 and 5895.93 angstrom. Calculate the spin orbit constant ξ and energy of 2P level. (2 pts)
3. The shift in energy levels due to spin-orbit coupling (for $l > 0$) is given by expression:

$$\Delta E = \frac{(E_n)^2}{mc^2} \left(\frac{n[j(j+1) - l(l+1) - s(s+1)]}{l(l+1/2)(l+1)} \right)$$

where, E_n is nth bohr orbital energy (2 + 2 + 2 + 2 pts)

- (a) Derive general expression for difference between the $j = l + 1/2$ and $j = l - 1/2$ levels
- (b) Calculate the energy difference due to spin-orbit coupling between $3p \ ^2P_{1/2}$ and $^2P_{3/2}$ states of a hydrogen atom.
- (c) Go online to the NIST Atomic Spectra Database (Energy Levels), retrieve the experimental energies for the two states (enter the keyword "H I" to get the energy levels for the neutral hydrogen atom), and compare the experimental difference to your calculated difference and comment on it.
- (d) Calculate the wavelength of the transition $3p \ (^2P_{1/2}) \rightarrow 1s \ (^2S_{1/2})$ and $3p \ (^2P_{3/2}) \rightarrow 1s \ (^2S_{1/2})$. Briefly discuss their location in electromagnetic spectrum and their separation.