AMATH/PMATH 331: Applied Real Analysis Fall 2026

Assignment 4; due Tuesday, November 10, 2026

---

[1] Let $A \subseteq \mathbb{R}^n$ be a non-empty subset, and let $\mathrm{Map}(A, \mathbb{R}^m)$ be the space of maps from A to $\mathbb{R}^m$.

[a] Let (f_k) be a sequence in $\mathrm{Map}(A, \mathbb{R}^m)$ which converges uniformly to a *bounded* function f, so $f \in B(A, \mathbb{R}^m)$. Prove that the f_k are *eventually bounded*. That is, prove that there exists $N \in \mathbb{N}$, such that $k \geq N$ implies $f_k \in B(A, \mathbb{R}^m)$.

[b] Suppose (f_k) and (g_k) are two sequences in $\mathrm{Map}(A, \mathbb{R}^n)$ which converge uniformly to *bounded* functions $f, g \in B(A, \mathbb{R}^m)$. Prove that the sequence $(\langle f_k, g_k \rangle)_{k \in \mathbb{N}}$ in $\mathrm{Map}(A, \mathbb{R})$ converges uniformly to $f \cdot g \in \mathrm{Map}(A, \mathbb{R})$, where $\langle \cdot, \cdot \rangle$ is the dot product in $\mathbb{R}^m$.

[c] Show that the assumption of boundedness of the limits in part [b] is essential as follows. Show that the sequence (f_k) in $M(\mathbb{R}, \mathbb{R})$ given by $f_k(x) = x + \frac{1}{k}$ converges uniformly on $\mathbb{R}$ to the unbounded function $f(x) = x$, but that $f_k^2 \in M(\mathbb{R}, \mathbb{R})$ does not converge uniformly to f^2.

---

[2] Let $f \in \mathrm{Map}(\mathbb{R}^n, \mathbb{R}^m)$ be a map, and suppose that f is *continuous* at $\mathbf{x}_0 \in \mathbb{R}^n$.

Define a sequence (f_k) in $\mathrm{Map}(\mathbb{R}^n, \mathbb{R}^m)$ by $f_k(\mathbf{x}) = f(\mathbf{x}_0 + \frac{1}{k}\mathbf{x})$. Let $g \in \mathrm{Map}(\mathbb{R}^n, \mathbb{R}^m)$ be the *constant map* $g(\mathbf{x}) = f(\mathbf{x}_0)$ for all $\mathbf{x} \in \mathbb{R}^n$.

Prove that, for any $R > 0$, the sequence (f_k) converges uniformly to g on the set $B_R(\mathbf{0})$.

---

[3] For $k \in \mathbb{N}$, define $f_k \colon [-1, 1] \to \mathbb{R}$ by $f_k(x) = (1 - x^2)^k$.

[a] For any $\delta \in (0, 1)$, prove that the sequence of functions (f_k) *converge uniformly* on the subset $[-1, -\delta] \cup [\delta, 1]$ of $\mathbb{R}$.

[b] Prove that the sequence of functions (f_k) *does not converge uniformly* on the subset $[-1, 1]$ of $\mathbb{R}$.

---

[4] Let $A \subseteq \mathbb{R}^n$ be non-empty. Let (f_k) be a sequence in $\mathrm{Map}(A, \mathbb{R}^m)$ which converges *uniformly* to a limit $f \in \mathrm{Map}(A, \mathbb{R}^m)$. We proved in class that if every f_k is continuous on A, then f is also continuous on A.

[a] Adapt that proof to show that if every f_k is *uniformly* continuous on A, then f is also *uniformly* continuous on A.

[b] Explain why we do not need to do any extra work if A is compact. That is, why does part [a] follow immediately if A is compact?

---

[5] COMING SOON

---

[6] COMING SOON

---

[7] COMING SOON

---

[8] COMING SOON

---

[9] COMING SOON

---

[10] COMING SOON
