

Quantum Amplifiers and Optimal Cloning: Applications in Quantum Metrology

Detailed Analysis and Applications

Contents

1	Introduction	3
1.1	Motivation	3
2	The No-Cloning Theorem	3
2.1	Statement of the Theorem	3
2.2	Mathematical Proof	3
2.3	Physical Implications	4
3	Optimal Quantum Cloning	4
3.1	Universal Quantum Cloning Machine (UQCM)	4
3.1.1	Definition	4
3.1.2	Optimal Fidelity	4
3.2	State-Dependent Cloning	4
3.2.1	Phase-Covariant Cloning	4
4	Quantum Amplifiers as Optimal Cloners	5
4.1	Conceptual Connection	5
4.2	Information-Theoretic Perspective	5
4.3	Quantum Amplification Map	5
5	Quantum Metrology and Amplification	5
5.1	Parameter Estimation Framework	5
5.2	Cramér-Rao Bound	5
5.3	Role of Amplification	6
5.3.1	Signal Enhancement	6
5.3.2	Noise Figure	6
5.4	Cloning-Based Metrology	6
5.4.1	Strategy	6
5.4.2	Fidelity-Precision Trade-off	6
6	Types of Quantum Amplifiers	6
6.1	Phase-Preserving Amplifiers	6
6.2	Phase-Sensitive Amplifiers	7
6.3	Noiseless Linear Amplification	7

7	Fundamental Limits and Trade-offs	7
7.1	Amplification vs Noise	7
7.2	Bandwidth-Gain Product	7
7.3	Quantum-to-Classical Transition	7
8	Applications in Quantum Metrology	7
8.1	Gravitational Wave Detection	7
8.2	Atomic Clocks	8
8.3	Quantum Sensing	8
8.4	Quantum Imaging	8
9	Experimental Implementations	8
9.1	Optical Systems	8
9.1.1	Parametric Down-Conversion	8
9.1.2	Four-Wave Mixing	9
9.2	Superconducting Circuits	9
9.2.1	Josephson Parametric Amplifier	9
9.3	Atomic Systems	9
9.3.1	Atomic Ensemble Amplification	9
10	Theoretical Framework	9
10.1	Quantum Channel Representation	9
10.2	Covariance Matrix Formalism	10
10.3	Quantum Amplifier Inequality	10
11	Recent Developments	10
11.1	Deterministic Amplification	10
11.2	Quantum-Enhanced Amplifiers	10
11.3	Machine Learning for Optimal Cloning	10
12	Challenges and Future Directions	11
12.1	Current Challenges	11
12.2	Future Prospects	11
13	Summary	11
13.1	Key Takeaways	11
13.2	Relationship to Quantum Information	11
14	Multiple Choice Questions	12
14.1	Conceptual Understanding	12
14.2	Mathematical and Technical Questions	14
14.3	Application-Based Questions	16
14.4	Advanced Conceptual Questions	17
15	Answer Key	18
16	Further Reading	18
16.1	Foundational Papers	18
16.2	Recent Reviews	18

1 Introduction

Quantum amplifiers represent a fundamental class of quantum information processing devices that enhance the distinguishability of quantum states while respecting the constraints imposed by quantum mechanics. Unlike classical amplifiers, quantum amplifiers must contend with the no-cloning theorem and the uncertainty principle, leading to fundamental trade-offs between amplification and fidelity.

1.1 Motivation

The concept of quantum amplification is crucial in several contexts:

- **Quantum Metrology:** Amplifying weak quantum signals for precise measurements
- **Quantum Communication:** Boosting quantum signals over long distances
- **Quantum Computation:** Enhancing the robustness of quantum information
- **Fundamental Physics:** Understanding the limits of quantum information processing

2 The No-Cloning Theorem

2.1 Statement of the Theorem

Theorem 1 (No-Cloning Theorem). *It is impossible to create an exact copy of an arbitrary unknown quantum state.*

2.2 Mathematical Proof

Consider a cloning operation that acts as:

$$U|\psi\rangle|0\rangle = |\psi\rangle|\psi\rangle \quad (1)$$

For two arbitrary states $|\psi\rangle$ and $|\phi\rangle$:

$$U|\psi\rangle|0\rangle = |\psi\rangle|\psi\rangle \quad (2)$$

$$U|\phi\rangle|0\rangle = |\phi\rangle|\phi\rangle \quad (3)$$

Taking the inner product:

$$\langle\psi|U^\dagger U|\phi\rangle\langle 0|0\rangle = \langle\psi|\phi\rangle = (\langle\psi|\phi\rangle)^2 \quad (4)$$

This implies:

$$\langle\psi|\phi\rangle(1 - \langle\psi|\phi\rangle) = 0 \quad (5)$$

Therefore, either $\langle\psi|\phi\rangle = 0$ or $\langle\psi|\phi\rangle = 1$, meaning perfect cloning is only possible for orthogonal or identical states.

2.3 Physical Implications

The no-cloning theorem has profound consequences:

1. **Quantum Information Security:** Eavesdropping can be detected
2. **Measurement Disturbance:** Measurements inevitably disturb quantum states
3. **Information Trade-offs:** Perfect amplification is impossible

3 Optimal Quantum Cloning

Since perfect cloning is impossible, we seek *optimal* approximate cloning that maximizes fidelity while creating multiple copies.

3.1 Universal Quantum Cloning Machine (UQCM)

3.1.1 Definition

A universal quantum cloning machine produces copies with state-independent fidelity:

$$|\psi\rangle \rightarrow \rho_1 \otimes \rho_2 \quad (6)$$

where ρ_1 and ρ_2 are the reduced density matrices of the two copies.

3.1.2 Optimal Fidelity

For 1-to-2 cloning in dimension d , the optimal fidelity is:

$$F = \frac{2d+1}{3d} \quad (7)$$

For qubits ($d = 2$):

$$F_{qubit} = \frac{5}{6} \approx 0.833 \quad (8)$$

3.2 State-Dependent Cloning

If we have prior knowledge about the states to be cloned (e.g., they come from a known set), better fidelity can be achieved.

3.2.1 Phase-Covariant Cloning

For states on the equator of the Bloch sphere:

$$|\psi_\theta\rangle = \frac{1}{\sqrt{2}}(|0\rangle + e^{i\theta}|1\rangle) \quad (9)$$

The optimal fidelity is:

$$F_{PC} = \frac{1}{2} + \frac{1}{2\sqrt{2}} \approx 0.854 \quad (10)$$

4 Quantum Amplifiers as Optimal Cloners

4.1 Conceptual Connection

An optimal cloner can be viewed as a quantum amplifier because:

1. It increases the number of systems carrying the quantum information
2. The information is distributed across multiple copies
3. This makes the overall information more robust against noise
4. Multiple copies allow for better measurement strategies

4.2 Information-Theoretic Perspective

Consider the mutual information between the original state and measurement outcomes:

$$I(\rho : M) = S(\rho) + S(M) - S(\rho, M) \quad (11)$$

After cloning:

$$I(\rho^{\otimes N} : M) > I(\rho : M) \quad (12)$$

The cloned copies collectively provide more information about the original state.

4.3 Quantum Amplification Map

A quantum amplifier can be described by a completely positive trace-preserving (CPTP) map:

$$\mathcal{A} : \mathcal{H} \rightarrow \mathcal{H}^{\otimes N} \quad (13)$$

For a coherent state $|\alpha\rangle$:

$$\mathcal{A}|\alpha\rangle = |g\alpha\rangle + \text{noise} \quad (14)$$

where g is the gain.

5 Quantum Metrology and Amplification

5.1 Parameter Estimation Framework

In quantum metrology, we aim to estimate an unknown parameter θ encoded in a quantum state:

$$\rho(\theta) = U(\theta)\rho_0U^\dagger(\theta) \quad (15)$$

5.2 Cramér-Rao Bound

The precision of parameter estimation is bounded by:

$$\Delta\theta \geq \frac{1}{\sqrt{NF[\rho(\theta)]}} \quad (16)$$

where $F[\rho(\theta)]$ is the quantum Fisher information:

$$F[\rho(\theta)] = \text{Tr}[\rho(\theta)L_\theta^2] \quad (17)$$

and L_θ is the symmetric logarithmic derivative.

5.3 Role of Amplification

5.3.1 Signal Enhancement

Quantum amplifiers enhance weak signals before measurement:

$$\text{SNR}_{out} = G \cdot \text{SNR}_{in} \quad (18)$$

However, quantum noise adds:

$$\text{Total Noise} = G \cdot \text{Input Noise} + \text{Amplifier Noise} \quad (19)$$

5.3.2 Noise Figure

The noise figure quantifies the degradation of signal-to-noise ratio:

$$NF = \frac{\text{SNR}_{in}}{\text{SNR}_{out}/G} \quad (20)$$

For quantum-limited amplifiers:

$$NF_{quantum} = 2 \quad (3 \text{ dB}) \quad (21)$$

This is a fundamental limit arising from the uncertainty principle.

5.4 Cloning-Based Metrology

5.4.1 Strategy

1. Clone the probe state N times
2. Perform measurements on all copies
3. Combine measurement results

5.4.2 Fidelity-Precision Trade-off

With cloning fidelity F , the effective Fisher information becomes:

$$F_{eff} = N \cdot F \cdot F[\rho(\theta)] \quad (22)$$

The precision scales as:

$$\Delta\theta \sim \frac{1}{\sqrt{NF \cdot F[\rho(\theta)]}} \quad (23)$$

6 Types of Quantum Amplifiers

6.1 Phase-Preserving Amplifiers

These amplifiers preserve the phase of the input signal:

$$\hat{a}_{out} = \sqrt{G}\hat{a}_{in} + \sqrt{G-1}\hat{b}^\dagger \quad (24)$$

where $\hat{b}$ is a vacuum mode.

Examples:

- Optical parametric amplifiers (OPAs)
- Josephson parametric amplifiers (JPAs)

6.2 Phase-Sensitive Amplifiers

These amplifiers amplify one quadrature while de-amplifying the other:

$$\begin{pmatrix} \hat{X}_{out} \\ \hat{P}_{out} \end{pmatrix} = \begin{pmatrix} \sqrt{G} & 0 \\ 0 & 1/\sqrt{G} \end{pmatrix} \begin{pmatrix} \hat{X}_{in} \\ \hat{P}_{in} \end{pmatrix} \quad (25)$$

Example: Squeezed light generation

6.3 Noiseless Linear Amplification

This is a probabilistic amplifier with post-selection:

$$|\alpha\rangle \xrightarrow{P_{success}} |g\alpha\rangle \quad (26)$$

with probability:

$$P_{success} = \exp(-|\alpha|^2(g^2 - 1)) \quad (27)$$

7 Fundamental Limits and Trade-offs

7.1 Amplification vs Noise

The Caves limit for phase-preserving amplifiers:

$$\Delta X_{out}^2 \geq G \cdot \Delta X_{in}^2 + (G - 1) \quad (28)$$

This shows that amplification necessarily adds noise.

7.2 Bandwidth-Gain Product

For physical amplifiers:

$$G \cdot \Delta f \leq G_{max} \cdot \Delta f_{max} \quad (29)$$

High gain comes at the cost of reduced bandwidth.

7.3 Quantum-to-Classical Transition

As the number of clones increases:

$$\lim_{N \rightarrow \infty} F(N) \rightarrow F_{classical} \quad (30)$$

The quantum advantage diminishes with excessive cloning.

8 Applications in Quantum Metrology

8.1 Gravitational Wave Detection

Challenge: Detecting extremely weak signals ($\Delta L/L \sim 10^{-21}$)

Solution: Quantum amplifiers enhance the signal before photodetection

Implementation:

- Squeezed light injection
- Reduces shot noise below standard quantum limit
- Signal-recycling mirrors act as phase-sensitive amplifiers

8.2 Atomic Clocks

Precision: $\Delta f/f \sim 10^{-18}$

Role of Cloning:

- Multiple atomic ensembles probed simultaneously
- Collective enhancement of signal
- Robust against individual atom losses

8.3 Quantum Sensing

Magnetic Field Sensing:

$$B = \frac{\hbar\omega_L}{gJ\mu_B} \quad (31)$$

Amplification Strategy:

1. Prepare entangled probe state
2. Evolution under magnetic field
3. Clone the final state
4. Perform optimal measurements

8.4 Quantum Imaging

Goal: Image objects with minimal photon flux

Cloning-Based Protocol:

- Single photon probes the object
- Clone the reflected photon state
- Multiple measurements for reconstruction
- Reduced damage to sensitive samples

9 Experimental Implementations

9.1 Optical Systems

9.1.1 Parametric Down-Conversion

Process:

$$\omega_p \rightarrow \omega_s + \omega_i \quad (32)$$

Hamiltonian:

$$H = \chi \hat{a}_s^\dagger \hat{a}_i^\dagger \hat{a}_p + h.c. \quad (33)$$

9.1.2 Four-Wave Mixing

In nonlinear optical fibers:

$$\omega_1 + \omega_2 \rightarrow \omega_3 + \omega_4 \quad (34)$$

Gain:

$$G = 1 + \sinh^2(\gamma PL) \quad (35)$$

9.2 Superconducting Circuits

9.2.1 Josephson Parametric Amplifier

Hamiltonian:

$$H = \hbar\omega_r \hat{a}^\dagger \hat{a} - \frac{E_J}{2} \cos(\hat{\phi} + \phi_p(t)) \quad (36)$$

where $\phi_p(t) = \phi_0 \cos(\omega_p t)$ is the pump.

Performance:

- Gain: $G \sim 20$ dB
- Bandwidth: ~ 10 MHz
- Near quantum-limited noise

9.3 Atomic Systems

9.3.1 Atomic Ensemble Amplification

Using collective atomic excitations:

$$|\Psi\rangle = \frac{1}{\sqrt{N}} \sum_{j=1}^N |g_1 \dots e_j \dots g_N\rangle \quad (37)$$

Amplification through stimulated emission:

$$G = \exp(\alpha NL) \quad (38)$$

10 Theoretical Framework

10.1 Quantum Channel Representation

A quantum amplifier is a quantum channel:

$$\mathcal{E}(\rho) = \sum_k E_k \rho E_k^\dagger \quad (39)$$

with Kraus operators E_k satisfying:

$$\sum_k E_k^\dagger E_k = \mathbb{I} \quad (40)$$

10.2 Covariance Matrix Formalism

For Gaussian states, amplification is described by:

$$V_{out} = SV_{in}S^T + N \quad (41)$$

where:

- V is the covariance matrix
- S is the symplectic transformation (amplification)
- N is the noise covariance

10.3 Quantum Amplifier Inequality

For any quantum amplifier with gain G :

$$\Delta X_{out} \Delta P_{out} \geq \frac{G}{2} \quad (42)$$

This generalizes the uncertainty principle to amplified states.

11 Recent Developments

11.1 Deterministic Amplification

Recent protocols achieve deterministic (non-probabilistic) quantum amplification:

- Using continuous measurement and feedback
- Quantum error correction techniques
- Reservoir engineering

11.2 Quantum-Enhanced Amplifiers

Entanglement-Enhanced Amplification:

$$|\Psi\rangle_{amp} = \frac{1}{\sqrt{2}}(|0\rangle_S|0\rangle_A + |1\rangle_S|1\rangle_A) \quad (43)$$

Provides:

- Better noise performance
- Enhanced bandwidth
- Surpasses classical limits

11.3 Machine Learning for Optimal Cloning

Neural networks optimize cloning protocols:

$$F_{optimal} = \max_{\{E_k\}} \langle \psi | \mathcal{E}(|\psi\rangle \langle \psi|) | \psi \rangle \quad (44)$$

12 Challenges and Future Directions

12.1 Current Challenges

1. **Decoherence:** Environmental noise limits amplification
2. **Bandwidth:** Trade-off with gain
3. **Scalability:** Difficult to amplify multi-mode states
4. **Integration:** Combining with existing quantum systems

12.2 Future Prospects

- **Quantum networks:** Long-distance quantum communication
- **Quantum computing:** Error mitigation through amplification
- **Fundamental tests:** Probing quantum-to-classical boundary
- **Quantum sensing:** Ultra-precise measurements

13 Summary

13.1 Key Takeaways

1. Quantum amplifiers are constrained by fundamental quantum limits
2. Optimal cloning provides a framework for understanding amplification
3. Trade-offs exist between fidelity, gain, and noise
4. Applications in metrology leverage multiple copies for enhanced precision
5. Various physical implementations exist across different platforms

13.2 Relationship to Quantum Information

Quantum amplification sits at the intersection of:

- Quantum information theory (no-cloning, entanglement)
- Quantum optics (coherent states, squeezed light)
- Quantum metrology (precision measurements)
- Quantum computing (error correction)

14 Multiple Choice Questions

14.1 Conceptual Understanding

1. **What does the no-cloning theorem state?**

- (A) It is impossible to copy any quantum state
- (B) It is impossible to copy an arbitrary unknown quantum state perfectly
- (C) Quantum states can only be copied with 50% fidelity
- (D) Only orthogonal quantum states cannot be copied

Answer: (B)

2. **For a universal 1-to-2 quantum cloning machine for qubits, what is the optimal fidelity?**

- (A) $F = 1/2$
- (B) $F = 2/3$
- (C) $F = 5/6$
- (D) $F = 1$

Answer: (C)

3. **Why can an optimal cloner be viewed as a quantum amplifier?**

- (A) It increases the amplitude of quantum states
- (B) It increases the energy of quantum states
- (C) It distributes information across multiple copies, making it more robust
- (D) It increases the entanglement of quantum states

Answer: (C)

4. **What is the quantum noise figure for an ideal phase-preserving amplifier?**

- (A) 0 dB
- (B) 3 dB
- (C) 6 dB
- (D) It depends on the gain

Answer: (B)

5. **In quantum metrology, the Cramér-Rao bound states that the precision of parameter estimation:**

- (A) Is unlimited with enough measurements
- (B) Is bounded by the quantum Fisher information
- (C) Improves linearly with the number of measurements
- (D) Is independent of the quantum state used

Answer: (B)

6. What is the main trade-off in quantum amplification?

- (A) Energy vs. momentum
- (B) Time vs. frequency
- (C) Gain vs. noise
- (D) Speed vs. accuracy

Answer: (C)

7. Phase-sensitive amplifiers:

- (A) Amplify all quadratures equally
- (B) Amplify one quadrature while de-amplifying the other
- (C) Only work at specific frequencies
- (D) Cannot be realized physically

Answer: (B)

8. Noiseless linear amplification is:

- (A) Deterministic and adds no noise
- (B) Probabilistic but adds no noise when successful
- (C) Impossible according to quantum mechanics
- (D) Only possible for classical states

Answer: (B)

9. The Caves limit for quantum amplifiers states that:

- (A) Amplification is always noiseless
- (B) Amplification necessarily adds quantum noise
- (C) Only phase information can be amplified
- (D) Classical amplifiers perform better than quantum ones

Answer: (B)

10. In gravitational wave detection, quantum amplifiers are used to:

- (A) Generate the gravitational waves
- (B) Increase the mass of the mirrors
- (C) Reduce shot noise through squeezed light
- (D) Amplify the gravitational waves themselves

Answer: (C)

14.2 Mathematical and Technical Questions

resume **For a quantum amplifier with gain G , the minimum added noise power is:**

- (A) 0
- (B) $(G - 1)\hbar\omega/2$
- (C) $G\hbar\omega/2$
- (D) $G^2\hbar\omega/2$

Answer: (B)

resume **The quantum Fisher information for a pure state $|\psi(\theta)\rangle$ is:**

- (A) $F = |\partial_\theta \psi(\theta)|^2$
- (B) $F = 4(\langle \partial_\theta \psi | \partial_\theta \psi \rangle - |\langle \psi | \partial_\theta \psi \rangle|^2)$
- (C) $F = \text{Tr}[\rho \partial_\theta \rho]$
- (D) $F = |\langle \psi | \partial_\theta \psi \rangle|$

Answer: (B)

resume **In the Heisenberg picture, a phase-preserving amplifier transforms the annihilation operator as:**

- (A) $\hat{a}_{out} = G\hat{a}_{in}$
- (B) $\hat{a}_{out} = \sqrt{G}\hat{a}_{in} + \sqrt{G-1}\hat{b}^\dagger$
- (C) $\hat{a}_{out} = \hat{a}_{in} + \sqrt{G}\hat{b}$
- (D) $\hat{a}_{out} = \exp(G)\hat{a}_{in}$

Answer: (B)

resume **The success probability for noiseless linear amplification of a coherent state $|\alpha\rangle$ with gain g is:**

- (A) $P = 1$
- (B) $P = \exp(-|\alpha|^2 g^2)$
- (C) $P = \exp(-|\alpha|^2 (g^2 - 1))$
- (D) $P = 1/g^2$

Answer: (C)

resume **For optimal universal cloning in dimension d , the fidelity is:**

- (A) $F = 1/d$
- (B) $F = (d + 1)/(2d)$
- (C) $F = (2d + 1)/(3d)$
- (D) $F = (d + 1)/d$

Answer: (C)

resume **A Josephson parametric amplifier typically achieves:**

- (A) Gain ~ 10 dB, far from quantum-limited noise
- (B) Gain ~ 20 dB, near quantum-limited noise
- (C) Gain ~ 60 dB, classical noise
- (D) Infinite gain with zero noise

Answer: (B)

resume **The bandwidth-gain product for a physical amplifier:**

- (A) Is unlimited
- (B) Is fundamentally limited by quantum mechanics
- (C) Depends only on the material used
- (D) Can be arbitrarily increased with more power

Answer: (B)

resume **In four-wave mixing, the gain is given by:**

- (A) $G = \gamma PL$
- (B) $G = 1 + \sinh^2(\gamma PL)$
- (C) $G = \exp(\gamma PL)$
- (D) $G = \cosh(\gamma PL)$

Answer: (B)

resume **For cloning-based metrology with N copies and cloning fidelity F , the precision scales as:**

- (A) $\Delta\theta \sim 1/N$
- (B) $\Delta\theta \sim 1/\sqrt{NF}$
- (C) $\Delta\theta \sim 1/F$
- (D) $\Delta\theta \sim \sqrt{F/N}$

Answer: (B)

resume **The covariance matrix after amplification with gain matrix S and noise N is:**

- (A) $V_{out} = SV_{in} + N$
- (B) $V_{out} = SV_{in}S^T + N$
- (C) $V_{out} = V_{in}S + N$
- (D) $V_{out} = S(V_{in} + N)S^T$

Answer: (B)

14.3 Application-Based Questions

resume **In atomic clocks, cloning helps by:**

- (A) Increasing the clock frequency
- (B) Making the atoms move faster
- (C) Allowing simultaneous probing of multiple atomic ensembles
- (D) Reducing the energy consumption

Answer: (C)

resume **Squeezed light in gravitational wave detectors:**

- (A) Increases the laser power
- (B) Reduces noise in one quadrature below the standard quantum limit
- (C) Makes the mirrors heavier
- (D) Has no real benefit

Answer: (B)

resume **The main advantage of quantum imaging with cloning is:**

- (A) Higher resolution than classical imaging
- (B) Reduced photon flux and sample damage
- (C) Faster image acquisition
- (D) Ability to see through opaque objects

Answer: (B)

resume **In quantum communication, quantum amplifiers:**

- (A) Allow perfect transmission over any distance
- (B) Boost signals while adding minimal quantum noise
- (C) Remove all noise from the channel
- (D) Are not useful due to the no-cloning theorem

Answer: (B)

resume **Parametric down-conversion in quantum optics:**

- (A) Converts high-energy photons to low-energy photons
- (B) Splits one photon into two correlated photons
- (C) Combines two photons into one
- (D) Has nothing to do with amplification

Answer: (B)

14.4 Advanced Conceptual Questions

resume **The quantum-to-classical transition in cloning means:**

- (A) Quantum effects disappear completely
- (B) As the number of clones increases, the quantum advantage diminishes
- (C) Classical cloning becomes possible
- (D) The no-cloning theorem no longer applies

Answer: (B)

resume **Entanglement-enhanced amplification:**

- (A) Uses entanglement between signal and ancilla to improve performance
- (B) Is impossible according to quantum mechanics
- (C) Only works for classical states
- (D) Removes the need for quantum amplifiers

Answer: (A)

resume **The primary challenge in scaling quantum amplifiers is:**

- (A) Cost of materials
- (B) Decoherence and environmental noise
- (C) Lack of theoretical understanding
- (D) Patent restrictions

Answer: (B)

resume **Machine learning for optimal cloning:**

- (A) Cannot improve beyond theoretical limits
- (B) Can help find optimal protocols within theoretical bounds
- (C) Makes the no-cloning theorem irrelevant
- (D) Only works for classical data

Answer: (B)

resume **The relationship between quantum amplifiers and quantum error correction is:**

- (A) They are unrelated fields
- (B) Amplification can be viewed as a form of redundancy encoding
- (C) Error correction makes amplification unnecessary
- (D) Amplification always increases errors

Answer: (B)

Q	Ans	Q	Ans	Q	Ans	Q	Ans
1	B	6	C	11	B	16	B
2	C	7	B	12	B	17	B
3	C	8	B	13	B	18	B
4	B	9	B	14	C	19	B
5	B	10	C	15	C	20	B
21	C	23	B	25	B	27	B
22	B	24	B	26	A	28	B
						29	B
						30	B

Table 1: Complete Answer Key for All MCQs

15 Answer Key

16 Further Reading

16.1 Foundational Papers

- Wootters, W. K., & Zurek, W. H. (1982). A single quantum cannot be cloned. *Nature*, 299(5886), 802-803.
- Bužek, V., & Hillery, M. (1996). Quantum copying: Beyond the no-cloning theorem. *Physical Review A*, 54(3), 1844.
- Caves, C. M. (1982). Quantum limits on noise in linear amplifiers. *Physical Review D*, 26(8), 1817.

16.2 Recent Reviews

- Scarani, V., et al. (2005). Quantum cloning. *Reviews of Modern Physics*, 77(4), 1225.
- Clerk, A. A., et al. (2010). Introduction to quantum noise, measurement, and amplification. *Reviews of Modern Physics*, 82(2), 1155.